

Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Foods Eaten Away from Home, by Race/Ethnicity and Age, in the United States, 2005-2006

Race/ethnicity and age (years)	Percent reporting ³ % (SE)	Food energy % (SE)	Protein % (SE)	Carbo- hydrate % (SE)	Total sugars % (SE)	Dietary fiber % (SE)	Total fat % (SE)	Saturated fat % (SE)	Mono- unsaturated fat % (SE)	Poly- unsaturated fat % (SE)
Non-Hispanic White:										
2 - 5.....	66 (5.1)	30 (2.7)	26 (2.0)	31 (3.2)	31 (3.0)	27 (3.5)	31 (2.5)	29 (2.2)	32 (2.8)	34 (2.8)
6 - 11.....	82 (3.1)	41 (2.1)	38 (2.3)	41 (2.1)	43 (2.4)	37 (2.3)	42 (2.4)	41 (2.2)	42 (2.7)	43 (2.3)
12 - 19.....	75 (2.8)	38 (2.2)	35 (2.4)	38 (2.4)	39 (2.7)	36 (2.1)	39 (2.0)	37 (2.0)	40 (2.0)	39 (2.4)
20 and over...	72 (0.8)	38 (1.2)	36 (1.2)	37 (1.1)	37 (1.0)	33 (1.1)	38 (1.3)	37 (1.3)	39 (1.3)	40 (1.2)
2 and over...	72 (1.0)	38 (0.9)	36 (1.0)	37 (0.8)	37 (0.7)	33 (0.9)	38 (1.0)	37 (1.0)	39 (1.0)	40 (0.9)
Non-Hispanic Black:										
2 - 5.....	63 (6.1)	36 (3.6)	35 (3.6)	35 (3.5)	36 (3.8)	36 (4.0)	38 (3.7)	38 (3.9)	38 (3.5)	37 (3.6)
6 - 11.....	76 (4.3)	43 (3.5)	41 (3.7)	44 (3.6)	48 (3.5)	44 (3.6)	42 (3.6)	43 (3.5)	42 (3.5)	42 (4.2)
12 - 19.....	70 (3.0)	35 (2.1)	34 (2.1)	35 (2.2)	36 (2.2)	34 (2.0)	36 (2.2)	35 (2.0)	36 (2.2)	35 (2.8)
20 and over...	63 (1.3)	31 (0.9)	28 (1.0)	33 (1.1)	34 (1.1)	28 (1.1)	31 (1.0)	30 (1.1)	32 (1.0)	33 (1.3)
2 and over...	65 (1.3)	33 (1.0)	31 (1.1)	34 (1.1)	36 (1.2)	31 (0.7)	33 (1.1)	33 (1.1)	34 (1.2)	34 (1.2)
Mexican American:										
2 - 5.....	58 (3.2)	23 (2.4)	23 (2.2)	23 (2.3)	23 (2.2)	19 (2.7)	24 (3.0)	23 (2.7)	26 (3.2)	25 (3.8)
6 - 11.....	81 (3.4)	38 (3.6)	38 (3.4)	38 (3.5)	40 (3.4)	33 (4.0)	39 (4.0)	39 (4.0)	39 (3.6)	36 (4.6)
12 - 19.....	72 (2.6)	35 (2.2)	34 (1.6)	34 (2.6)	34 (2.9)	33 (2.9)	36 (2.1)	35 (2.2)	36 (2.1)	37 (2.6)
20 and over...	69 (2.0)	40 (1.5)	39 (1.9)	39 (1.4)	39 (1.5)	34 (1.7)	43 (1.5)	42 (1.4)	45 (1.6)	45 (1.9)
2 and over...	70 (1.7)	38 (1.1)	37 (1.4)	37 (1.1)	37 (1.3)	33 (1.3)	40 (1.1)	39 (0.9)	41 (1.2)	41 (1.4)
All Individuals⁴:										
2 - 5.....	64 (2.9)	29 (1.9)	26 (1.5)	29 (2.2)	29 (2.0)	27 (2.3)	30 (2.0)	29 (1.8)	32 (2.1)	32 (2.3)
6 - 11.....	80 (2.6)	39 (1.8)	37 (2.0)	40 (1.7)	42 (1.8)	36 (1.6)	40 (2.1)	40 (2.0)	40 (2.2)	41 (2.1)
12 - 19.....	74 (1.9)	37 (1.5)	35 (1.7)	37 (1.6)	39 (1.8)	35 (1.4)	38 (1.4)	37 (1.4)	39 (1.4)	38 (1.7)
20 and over...	70 (0.8)	37 (0.9)	35 (1.0)	36 (0.9)	36 (0.8)	33 (0.9)	38 (1.0)	37 (1.0)	39 (1.0)	40 (1.0)
2 and over...	71 (0.9)	37 (0.7)	35 (0.8)	36 (0.7)	37 (0.6)	33 (0.7)	38 (0.8)	37 (0.8)	39 (0.8)	39 (0.8)

Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Foods Eaten Away from Home, by Race/Ethnicity and Age, in the United States, 2005-2006 (continued)

Race/ethnicity and age (years)	Choles- terol % (SE)	Vitamin A (RAE) % (SE)	Beta- carotene % (SE)	Lycopene % (SE)	Thiamin % (SE)	Ribo- flavin % (SE)	Niacin % (SE)	Vitamin B6 % (SE)	Folate (DFE) % (SE)
Non-Hispanic White:									
2 - 5.....	27 (1.7)	21 (2.9)	35 (7.6)	30 (5.0)	25 (4.0)	24 (2.7)	26 (2.8)	22 (3.3)	23 (3.8)
6 - 11.....	39 (3.4)	30 (2.5)	28 (6.0)	42 (4.5)	34 (2.7)	34 (2.2)	36 (2.4)	33 (2.3)	31 (3.4)
12 - 19.....	36 (2.4)	26 (2.6)	39 (7.7)	40 (4.3)	31 (2.2)	28 (2.0)	34 (2.5)	32 (2.0)	28 (1.7)
20 and over...	38 (1.7)	27 (1.2)	35 (1.8)	37 (2.7)	33 (1.5)	31 (1.3)	36 (1.3)	33 (1.6)	30 (1.2)
2 and over...	38 (1.5)	27 (1.0)	35 (1.5)	37 (2.2)	33 (1.2)	30 (1.1)	36 (1.0)	33 (1.3)	30 (1.0)
Non-Hispanic Black:									
2 - 5.....	32 (5.0)	31 (4.2)	39 (7.1)	38 (9.7)	31 (3.6)	33 (3.4)	30 (3.5)	28 (4.1)	27 (2.1)
6 - 11.....	35 (4.5)	45 (5.1)	46 (7.8)	40 (4.6)	42 (3.9)	45 (4.4)	40 (3.9)	40 (4.2)	41 (3.1)
12 - 19.....	34 (3.3)	29 (2.5)	32 (4.7)	40 (3.3)	30 (1.8)	32 (1.9)	32 (2.0)	31 (1.8)	26 (2.0)
20 and over...	28 (1.4)	24 (1.6)	26 (2.5)	25 (3.8)	27 (1.3)	27 (1.1)	29 (1.0)	27 (1.0)	27 (1.5)
2 and over...	30 (1.8)	27 (1.5)	29 (2.4)	30 (2.4)	29 (1.0)	30 (1.1)	30 (0.8)	29 (0.9)	28 (1.0)
Mexican American:									
2 - 5.....	18 (2.9)	17 (2.0)	22 (5.3)	20 (3.6)	19 (1.9)	18 (1.9)	22 (2.0)	17 (1.6)	17 (1.7)
6 - 11.....	38 (3.7)	38 (4.9)	45 (12.0)	37 (6.0)	35 (3.6)	38 (3.3)	35 (3.6)	34 (4.1)	34 (4.3)
12 - 19.....	33 (1.5)	25 (2.9)	33 (6.2)	37 (5.7)	30 (2.3)	28 (2.0)	33 (2.1)	29 (2.2)	27 (3.1)
20 and over...	37 (2.2)	30 (1.5)	36 (3.7)	40 (3.3)	36 (1.7)	35 (1.5)	41 (1.7)	36 (2.2)	35 (2.1)
2 and over...	35 (1.7)	29 (1.6)	36 (4.0)	38 (2.8)	34 (1.0)	33 (1.2)	38 (1.2)	34 (1.4)	32 (1.1)
All Individuals⁴:									
2 - 5.....	24 (1.5)	21 (1.7)	29 (4.6)	30 (3.5)	24 (2.5)	23 (1.8)	26 (1.8)	22 (1.9)	23 (2.3)
6 - 11.....	36 (3.1)	32 (2.1)	33 (3.1)	39 (3.3)	34 (2.1)	35 (1.8)	36 (2.1)	33 (2.0)	32 (2.4)
12 - 19.....	35 (1.6)	26 (1.9)	37 (6.0)	39 (3.3)	31 (1.5)	29 (1.4)	34 (1.6)	31 (1.4)	28 (1.2)
20 and over...	37 (1.4)	27 (1.0)	34 (1.4)	36 (1.9)	33 (1.1)	31 (1.0)	35 (1.0)	33 (1.2)	30 (0.9)
2 and over...	36 (1.3)	27 (0.8)	34 (1.2)	36 (1.6)	32 (0.9)	31 (0.8)	35 (0.8)	32 (1.0)	30 (0.8)

Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Foods Eaten Away from Home, by Race/Ethnicity and Age, in the United States, 2005-2006 (continued)

Race/ethnicity and age (years)	Choline % (SE)	Vitamin B12 % (SE)	Vitamin C % (SE)	Vitamin E (alpha- tocopherol) % (SE)	Vitamin K % (SE)	Calcium % (SE)	Phosphorus % (SE)	Magnesium % (SE)
Non-Hispanic White:								
2 - 5.....	25 (1.8)	20 (2.5)	32 (4.4)	31 (3.1)	31 (2.2)	23 (2.0)	25 (2.1)	26 (2.7)
6 - 11.....	37 (2.4)	32 (3.3)	38 (2.9)	40 (2.3)	38 (2.7)	35 (2.2)	38 (2.0)	38 (2.4)
12 - 19.....	34 (1.9)	33 (2.4)	26 (2.8)	35 (2.1)	36 (3.2)	29 (2.2)	34 (2.2)	34 (1.9)
20 and over...	35 (1.4)	31 (2.6)	30 (1.8)	36 (1.7)	38 (1.7)	31 (1.0)	34 (1.2)	33 (1.1)
2 and over...	35 (1.2)	31 (2.1)	30 (1.4)	36 (1.4)	38 (1.5)	31 (0.9)	34 (1.0)	33 (1.0)
Non-Hispanic Black:								
2 - 5.....	34 (3.9)	32 (4.3)	30 (4.8)	39 (5.0)	41 (6.4)	38 (3.8)	37 (3.4)	35 (3.6)
6 - 11.....	40 (4.1)	43 (5.6)	42 (3.6)	42 (4.1)	43 (4.6)	48 (3.8)	46 (4.1)	43 (3.2)
12 - 19.....	35 (2.6)	31 (2.1)	32 (2.6)	36 (2.9)	34 (5.0)	33 (1.9)	35 (1.9)	34 (1.9)
20 and over...	27 (1.0)	27 (2.1)	28 (2.5)	30 (1.4)	24 (2.0)	27 (1.2)	29 (0.9)	29 (1.0)
2 and over...	30 (1.2)	30 (1.7)	30 (1.5)	32 (0.8)	27 (1.8)	31 (1.0)	32 (1.0)	31 (0.9)
Mexican American:								
2 - 5.....	19 (2.3)	18 (2.3)	17 (1.9)	23 (3.1)	26 (3.8)	17 (2.2)	20 (2.3)	19 (2.3)
6 - 11.....	38 (3.4)	37 (3.2)	35 (5.0)	36 (4.9)	34 (5.9)	40 (3.1)	38 (3.1)	35 (3.5)
12 - 19.....	32 (1.4)	31 (3.0)	24 (4.0)	35 (2.3)	37 (2.8)	30 (1.6)	32 (1.7)	31 (1.9)
20 and over...	36 (2.0)	32 (3.6)	28 (2.0)	41 (2.1)	43 (1.2)	32 (1.0)	36 (1.7)	34 (1.8)
2 and over...	34 (1.6)	31 (2.6)	27 (1.7)	38 (1.7)	41 (0.9)	32 (0.8)	34 (1.2)	33 (1.4)
All Individuals⁴:								
2 - 5.....	25 (1.3)	21 (1.5)	27 (2.7)	30 (2.1)	30 (1.9)	23 (1.5)	25 (1.5)	25 (1.9)
6 - 11.....	36 (2.1)	33 (2.8)	36 (1.9)	38 (2.0)	36 (1.9)	37 (1.8)	38 (1.7)	37 (1.9)
12 - 19.....	34 (1.3)	32 (1.8)	26 (1.9)	35 (1.4)	36 (2.4)	29 (1.6)	34 (1.5)	34 (1.3)
20 and over...	35 (1.2)	31 (2.0)	29 (1.1)	35 (1.3)	37 (1.7)	31 (0.8)	33 (0.9)	33 (0.9)
2 and over...	34 (1.0)	30 (1.6)	29 (0.9)	35 (1.1)	36 (1.5)	31 (0.7)	34 (0.8)	33 (0.8)

Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Foods Eaten Away from Home, by Race/Ethnicity and Age, in the United States, 2005-2006 (continued)

Race/ethnicity and age (years)	Iron % (SE)	Zinc % (SE)	Copper % (SE)	Selenium % (SE)	Potassium % (SE)	Sodium [†] (adjusted) % (SE)	Caffeine % (SE)	Alcohol ^s % (SE)
Non-Hispanic White:								
2 - 5.....	23 (3.7)	21 (2.0)	29 (3.2)	27 (3.0)	26 (2.7)	29 (2.7)	36 (7.3)	-- --
6 - 11.....	34 (3.0)	35 (2.9)	41 (2.2)	37 (3.0)	38 (1.9)	41 (1.7)	49 (4.5)	-- --
12 - 19.....	31 (1.9)	33 (2.1)	36 (2.1)	35 (2.6)	34 (1.8)	38 (2.9)	48 (4.3)	-- --
20 and over...	32 (1.3)	34 (1.7)	34 (1.4)	36 (1.3)	33 (1.2)	39 (1.2)	36 (2.0)	43 (3.1)
2 and over...	32 (1.1)	33 (1.5)	34 (1.3)	36 (1.0)	33 (1.0)	39 (0.9)	37 (1.9)	-- --
Non-Hispanic Black:								
2 - 5.....	29 (3.4)	32 (3.0)	34 (4.3)	34 (3.5)	36 (4.0)	36 (3.8)	36 (10.3)	-- --
6 - 11.....	40 (3.7)	41 (2.8)	43 (2.7)	39 (3.8)	45 (3.7)	42 (3.9)	37 (3.5)	-- --
12 - 19.....	30 (1.6)	32 (1.7)	35 (1.8)	33 (2.3)	35 (2.0)	35 (2.3)	41 (6.5)	-- --
20 and over...	28 (1.2)	27 (1.4)	29 (1.1)	29 (1.1)	28 (1.0)	31 (1.1)	40 (1.9)	24 (3.7)
2 and over...	29 (1.0)	29 (1.2)	31 (1.0)	31 (1.1)	31 (1.0)	33 (1.1)	40 (1.8)	-- --
Mexican American:								
2 - 5.....	19 (1.9)	21 (2.1)	21 (2.4)	22 (2.0)	19 (2.2)	25 (2.5)	40 (4.6)	-- --
6 - 11.....	34 (3.7)	37 (3.4)	37 (3.7)	37 (3.8)	38 (3.6)	39 (3.8)	47 (5.9)	-- --
12 - 19.....	30 (2.6)	32 (2.1)	33 (2.0)	35 (1.6)	31 (2.1)	36 (1.8)	41 (3.1)	-- --
20 and over...	36 (1.6)	37 (2.0)	36 (2.0)	39 (2.1)	35 (1.7)	43 (1.4)	47 (3.6)	36 (6.2)
2 and over...	34 (0.9)	35 (1.4)	35 (1.7)	37 (1.6)	33 (1.2)	40 (1.0)	46 (3.2)	-- --
All Individuals⁴:								
2 - 5.....	24 (2.3)	23 (1.3)	27 (2.1)	26 (2.0)	26 (1.8)	29 (2.1)	35 (4.8)	-- --
6 - 11.....	34 (2.3)	35 (2.3)	39 (1.8)	36 (2.3)	38 (1.6)	39 (1.8)	47 (3.4)	-- --
12 - 19.....	31 (1.3)	33 (1.5)	35 (1.3)	35 (1.8)	33 (1.2)	37 (2.0)	48 (3.4)	-- --
20 and over...	32 (1.0)	34 (1.3)	34 (1.1)	35 (1.0)	32 (0.9)	38 (1.0)	38 (1.5)	42 (2.5)
2 and over...	31 (0.9)	33 (1.2)	34 (1.0)	35 (0.8)	33 (0.8)	38 (0.8)	38 (1.4)	-- --

Symbol Legend

† Sodium estimate adjusted for salt used in food preparation. Details available at: www.ars.usda.gov/ba/bhnrc/fsrg.

Footnotes

¹ Away from home includes any location other than home. Home is defined as an individual's dwelling unit and the surrounding areas that are used solely by the occupants of that dwelling unit.

² Percentages are estimated as a ratio of total nutrients from foods eaten away from home for all individuals to total daily nutrient intakes for all individuals. Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection. Total daily nutrient intakes are available from: www.ars.usda.gov/ba/bhnrc/fsrg. See Table 2. Nutrient Intakes from Food: Mean Amounts Consumed per Individual, by Race/Ethnicity and Age, in the United States, 2005-2006.

³ The percentage of respondents in the race/ethnicity/age group who reported consuming at least one item away from home.

⁴ Includes persons of all races and Hispanic origins, not just those presented separately.

⁵ Alcohol estimates are shown only for 20 years and over age groups. Although the data are collected for all individuals, estimates are suppressed due to extreme variability and/or inadequate sample size.

Abbreviations

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

Notes Applicable to All Tables in Series: *What We Eat in America, NHANES 2005-2006*

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2005-2006. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intake information was coded using the USDA Food and Nutrient Database for Dietary Studies 3.0 www.ars.usda.gov/ba/bhnrc/fsrg which is based on nutrient values in the USDA National Nutrient Database for Standard Reference, Release 20 (Agricultural Research Service, Nutrient Data Laboratory, 2008).

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Suggested Citation

U.S. Department of Agriculture, Agricultural Research Service. 2010. Away from Home: Percentages of Selected Nutrients Contributed by Foods Eaten Away from Home, by Race/Ethnicity and Age, *What We Eat in America*, NHANES 2005-2006. Available: www.ars.usda.gov/ba/bhnrc/fsrg.