Table 26. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Race/Ethnicity and Age, in the United States, 2013-2014

Race/ethnicity	Percent			Carbo-	Total	Dietary	Total	Saturated	Mono- unsaturated	Poly- unsaturated
and age	reporting ³	Energy	Protein	hydrate	sugars	fiber	fat	fat	fat	fat
(years)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)
Non-Hispanic White:										
2 - 5	99* (0.5)	29 (1.1)	19 (1.2)	34 (1.4)	40 (1.7)	24 (2.0)	28 (1.1)	28 (1.4)	27 (1.1)	30 (2.5)
6 - 11	99* (0.4)	25 (1.3)	14 (1.5)	29 (1.2)	36 (1.4)	23 (1.7)	24 (1.7)	25 (2.3)	23 (2.0)	23 (1.9)
12 - 19	93 (1.8)	22 (1.3)	12 (0.9)	27 (1.9)	36 (2.7)	19 (1.4)	20 (1.3)	21 (1.2)	19 (1.4)	19 (2.5)
20 and over	96 (0.6)	24 (0.7)	13 (0.7)	28 (0.7)	39 (1.1)	20 (0.8)	21 (0.9)	23 (0.9)	22 (0.9)	20 (1.2)
2 and over	96 (0.5)	24 (0.5)	13 (0.5)	28 (0.5)	38 (0.9)	20 (0.7)	21 (0.8)	23 (0.9)	21 (0.7)	20 (0.9)
Non-Hispanic Black:										
2 - 5	93* (3.3)	26 (2.1)	15 (1.9)	30 (1.8)	36 (2.5)	25 (1.7)	26 (3.1)	25 (3.0)	25 (2.7)	28 (4.1)
6 - 11	90 (1.7)	23 (2.3)	15 (2.1)	27 (2.4)	32 (2.2)	18 (1.7)	22 (2.6)	22 (2.5)	21 (2.4)	25 (3.3)
12 - 19	89 (1.5)	29 (1.9)	16 (1.1)	33 (2.4)	41 (3.1)	26 (2.1)	28 (1.7)	28 (2.2)	27 (1.5)	29 (1.6)
20 and over	92 (0.8)	28 (1.6)	15 (1.0)	32 (1.6)	40 (1.5)	24 (1.3)	25 (1.4)	25 (1.4)	24 (1.5)	25 (1.5)
2 and over	91 (0.7)	28 (1.4)	15 (0.9)	31 (1.4)	40 (1.5)	24 (1.2)	25 (1.4)	25 (1.5)	24 (1.4)	25 (1.5)
Non-Hispanic Asian4:										
2 - 5	97* (2.1)	31* (3.8)	19* (3.0)	35* (4.1)	48* (5.6)	30* (3.3)	32* (4.2)	36* (5.2)	28* (2.8)	28* (3.9)
6 - 11	97* (2.7)	26 (2.0)	16* (1.8)	28 (2.1)	41 (3.6)	24 (2.6)	28 (3.0)	33 (4.5)	26 (2.7)	22* (3.0)
12 - 19	93* (2.7)	24 (2.4)	14 (2.8)	26 (2.8)	36 (5.1)	21 (1.1)	26 (3.3)	29 (3.5)	25 (3.8)	22 (3.5)
20 and over	94 (1.3)	19 (1.0)	10 (0.8)	21 (1.1)	35 (1.5)	17 (0.9)	18 (1.2)	19 (1.4)	19 (1.6)	16 (1.1)
2 and over	94 (1.2)	20 (0.8)	11 (0.8)	23 (0.9)	36 (1.4)	18 (0.9)	20 (1.0)	22 (1.0)	21 (1.3)	17 (1.1)
Hispanic:										
2 - 5	99* (0.5)	34 (1.3)	25 (1.4)	39 (1.6)	49 (1.8)	30 (1.5)	33 (1.4)	35 (1.6)	31 (1.4)	31 (2.0)
6 - 11	95* (1.1)	24 (1.7)	15 (1.7)	28 (1.4)	34 (1.5)	22 (1.6)	24 (2.2)	24 (2.4)	23 (2.1)	25 (2.1)
12 - 19	91 (1.4)	23 (1.0)	13 (0.7)	27 (1.3)	37 (2.4)	20 (1.2)	22 (1.1)	23 (1.2)	22 (1.3)	22 (1.7)
20 and over	94 (1.3)	22 (0.6)	12 (0.5)	24 (0.9)	34 (1.3)	18 (0.8)	20 (1.0)	20 (1.1)	20 (1.2)	18 (0.8)
2 and over	94 (1.0)	23 (0.6)	13 (0.4)	26 (0.7)	35 (0.9)	19 (0.7)	21 (0.9)	22 (1.0)	21 (1.1)	20 (0.9)

Table 26. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Race/Ethnicity and Age, in the United States, 2013-2014 (continued)

Race/ethnicity and age (years)	Choles- terol % (SE)	Vitamin A (RAE) % (SE)	Beta- carotene % (SE)	Lycopene % (SE)	Thiamin % (SE)	Ribo- flavin % (SE)	Niacin % (SE)	Vitamin B6 % (SE)	Folate (DFE) % (SE)
Non-Hispanic White:									
2 - 5	16 (1.7)	24 (2.0)	20* (7.4)	5* (2.1)	20 (1.5)	25 (1.2)	17 (1.6)	19 (1.4)	18 (2.2)
6 - 11	12 (1.5)	18 (1.9)	17 (5.2)	6* (1.7)	17 (1.5)	21 (2.7)	14 (1.9)	16 (2.2)	17 (1.6)
12 - 19	10 (1.0)	22 (2.1)	25* (8.6)	5* (2.3)	17 (2.6)	22 (1.5)	16 (1.7)	19 (2.9)	15 (2.0)
20 and over	10 (0.6)	17 (1.2)	11 (1.9)	8 (1.5)	15 (0.6)	23 (0.9)	15 (0.6)	17 (1.0)	15 (0.6)
2 and over	10 (0.5)	18 (1.0)	13 (1.7)	7 (1.2)	16 (0.7)	23 (0.7)	15 (0.5)	17 (0.8)	16 (0.5)
Non-Hispanic Black:									
2 - 5	11 (1.9)	18 (1.9)	7* (1.6)	2* (0.5)	19 (1.5)	21 (1.6)	16 (1.4)	18 (1.7)	17 (2.0)
6 - 11	12 (1.8)	14 (1.8)	12 (3.1)	8* (4.0)	18 (3.1)	17 (2.1)	17 (3.4)	18 (4.2)	18 (3.2)
12 - 19	12 (1.1)	22 (2.6)	16 (4.4)	12* (5.5)	21 (2.2)	23 (2.2)	19 (1.7)	21 (2.6)	25 (4.7)
20 and over	11 (0.9)	16 (1.0)	13 (2.2)	11 (2.4)	18 (1.1)	22 (0.9)	18 (1.0)	19 (1.1)	19 (1.4)
2 and over	11 (0.8)	17 (1.0)	13 (2.1)	11 (2.2)	18 (1.1)	22 (0.9)	18 (0.9)	19 (1.0)	19 (1.6)
Non-Hispanic Asian4:									
2 - 5	15* (1.9)	22* (2.3)	9* (3.6)	1* (1.1)	22* (2.3)	28* (2.5)	18* (3.4)	21* (3.8)	17* (3.3)
6 - 11	13* (1.2)	21* (2.8)	15* (4.2)	17* (7.2)	16* (1.4)	22* (1.9)	12* (1.4)	15* (2.1)	12* (1.1)
12 - 19	11* (2.4)	21 (2.9)	6* (2.5)	8* (4.7)	17 (2.6)	24 (3.2)	15 (3.4)	16 (2.9)	17 (2.6)
20 and over	7 (0.9)	15 (2.3)	11 (3.1)	17 (4.6)	13 (0.8)	21 (1.4)	11 (0.9)	13 (1.0)	12 (1.0)
2 and over	8 (0.9)	16 (1.9)	11 (2.7)	15 (3.6)	14 (0.6)	21 (1.2)	12 (0.8)	14 (0.9)	13 (0.9)
Hispanic:									
2 - 5	18 (1.3)	31 (1.6)	26 (3.9)	25* (9.9)	27 (1.4)	34 (1.7)	22 (1.5)	26 (2.1)	25 (2.0)
6 - 11	12 (1.5)	17 (1.4)	18 (4.0)	9 (0.9)	17 (1.1)	18 (1.3)	16 (0.9)	16 (1.0)	17 (1.2)
12 - 19	10 (0.8)	17 (1.0)	12 (1.7)	7 (1.4)	15 (0.9)	18 (0.9)	14 (0.9)	15 (1.1)	16 (1.2)
20 and over	9 (0.6)	17 (1.1)	13 (2.5)	11 (1.8)	15 (0.5)	23 (1.0)	18 (1.1)	20 (1.4)	16 (0.7)
2 and over	10 (0.5)	18 (0.6)	14 (2.1)	11 (1.1)	16 (0.4)	22 (0.6)	17 (0.8)	19 (1.0)	17 (0.5)

Table 26. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Race/Ethnicity and Age, in the United States, 2013-2014 (continued)

Race/ethnicity					Vitamin E (alpha-				
and age	Choline	Vitamin B12	Vitamin C	Vitamin D	tocopherol)	Vitamin K	Calcium	Phosphorus	Magnesium
(years)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)
Non-Hispanic White:									
2 - 5	21 (1.8)	20 (1.5)	37 (3.1)	24 (2.1)	28 (2.6)	27 (2.6)	27 (1.8)	24 (1.5)	27 (1.4)
6 - 11	15 (1.4)	15 (2.2)	27 (2.3)	16 (2.4)	28 (2.3)	15 (2.2)	20 (1.9)	18 (1.5)	25 (1.7)
12 - 19	14 (1.2)	21 (2.9)	23 (4.3)	20 (2.9)	21 (2.1)	11 (1.7)	23 (1.1)	17 (1.0)	24 (1.4)
20 and over	16 (0.8)	15 (0.7)	22 (1.1)	16 (1.4)	24 (1.7)	9 (0.8)	24 (1.0)	19 (0.9)	27 (0.9)
2 and over	16 (0.6)	16 (0.8)	23 (1.1)	17 (1.2)	24 (1.4)	10 (0.7)	24 (0.8)	19 (0.7)	27 (0.7)
Non-Hispanic Black:									
2 - 5	15 (1.5)	15 (1.5)	27 (2.6)	22 (3.0)	30 (2.8)	16 (2.4)	23 (2.0)	19 (1.8)	25 (2.2)
6 - 11	14 (1.5)	13 (2.3)	27 (4.5)	13 (2.3)	25 (2.5)	16 (1.8)	18 (2.4)	17 (2.1)	21 (1.9)
12 - 19	16 (1.1)	19 (3.3)	34 (4.1)	22 (3.9)	35 (2.5)	20 (1.9)	25 (2.0)	21 (1.5)	29 (1.7)
20 and over	17 (1.5)	14 (1.0)	27 (1.4)	16 (2.2)	27 (1.6)	15 (2.9)	26 (1.1)	20 (1.3)	29 (1.2)
2 and over	17 (1.2)	15 (0.8)	28 (1.3)	17 (1.8)	28 (1.5)	16 (2.5)	25 (1.2)	20 (1.1)	28 (1.1)
Non-Hispanic Asian4:									
2 - 5	19* (1.9)	21* (3.0)	44* (7.6)	22* (3.8)	30* (4.0)	25* (6.8)	31* (3.3)	25* (2.9)	28* (2.8)
6 - 11	15* (2.1)	18* (1.8)	29 (4.2)	19* (4.6)	26 (2.7)	29 (7.2)	28 (2.6)	23 (2.2)	26 (2.1)
12 - 19	15 (1.9)	21 (4.6)	25* (8.5)	25 (3.6)	27 (5.8)	13* (4.1)	31 (3.2)	20 (2.9)	26 (2.3)
20 and over	10 (0.8)	12 (1.4)	26 (1.9)	14 (2.6)	23 (1.9)	5 (0.8)	25 (1.9)	14 (1.0)	22 (1.2)
2 and over	12 (0.9)	14 (1.5)	26 (1.9)	16 (2.4)	24 (2.0)	7 (1.1)	26 (1.6)	16 (1.1)	23 (1.1)
Hispanic:									
2 - 5	24 (1.3)	30 (1.9)	44 (4.0)	38 (2.2)	33 (1.7)	25 (2.3)	39 (2.0)	31 (1.6)	34 (1.5)
6 - 11	14 (1.5)	13 (1.1)	29 (2.7)	14 (1.6)	27 (2.0)	22 (2.8)	21 (2.2)	18 (1.8)	22 (1.3)
12 - 19	13 (0.8)	15 (1.1)	26 (2.4)	18 (1.5)	28 (3.2)	19 (2.6)	23 (1.4)	17 (0.9)	24 (1.0)
20 and over	14 (0.6)	18 (1.5)	25 (1.0)	18 (1.0)	23 (1.2)	11 (1.0)	25 (1.0)	17 (0.5)	25 (0.8)
2 and over	15 (0.5)	18 (0.9)	27 (1.0)	19 (0.8)	24 (1.2)	14 (1.0)	25 (0.9)	18 (0.5)	25 (0.7)

Table 26. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Race/Ethnicity and Age, in the United States, 2013-2014 (continued)

Race/ethnicity								
and age	Iron	Zinc	Copper	Selenium	Potassium	Sodium	Caffeine	Alcohol
(years)	% (SE)	% (SE)						
Non-Hispanic White:								
2 - 5	20 (1.4)	20 (1.5)	29 (1.4)	15 (1.1)	27 (1.0)	19 (1.6)	49 (4.5)	
6 - 11	19 (1.9)	17 (1.8)	28 (2.1)	11 (1.1)	20 (1.7)	14 (1.2)	47 (7.8)	
12 - 19	15 (1.6)	15 (1.2)	26 (1.6)	10 (0.9)	18 (0.9)	15 (1.0)	40 (4.4)	
20 and over	16 (0.7)	16 (0.8)	29 (1.0)	10 (0.6)	21 (0.8)	13 (0.4)	40 (2.1)	62 (3.4)
2 and over	16 (0.6)	16 (0.7)	28 (0.9)	10 (0.5)	21 (0.6)	13 (0.4)	40 (2.0)	
Non-Hispanic Black:								
2 - 5	21 (1.6)	18 (1.6)	27 (2.9)	13 (1.8)	22 (2.5)	17 (2.0)	42*(12.7)	
6 - 11	18 (3.0)	16 (1.8)	22 (1.8)	15 (2.9)	18 (1.4)	17 (2.3)	48 (7.5)	
12 - 19	24 (3.3)	20 (2.5)	28 (1.6)	15 (1.6)	23 (1.2)	19 (1.4)	43 (7.4)	
20 and over	19 (1.1)	16 (0.8)	28 (1.1)	12 (1.0)	23 (0.9)	16 (1.2)	32 (3.3)	72 (4.6)
2 and over	20 (1.2)	17 (0.8)	28 (0.9)	13 (0.9)	23 (0.7)	17 (1.0)	33 (3.2)	
Non-Hispanic Asian4:								
2 - 5	20* (3.7)	19* (2.6)	26* (3.9)	14* (2.5)	31* (2.3)	19* (3.1)	56*(14.3)	
6 - 11	17* (1.5)	16* (1.7)	26 (2.5)	15* (2.8)	24 (2.3)	16* (1.9)	63*(11.0)	
12 - 19	16 (2.8)	18 (3.3)	25 (2.9)	11* (2.3)	21 (3.0)	16 (2.7)	80 (5.5)	
20 and over	13 (1.0)	13 (1.1)	24 (1.4)	9 (0.9)	17 (0.9)	8 (1.0)	32 (3.1)	52 (6.1)
2 and over	14 (0.8)	14 (1.0)	24 (1.4)	10 (0.6)	19 (0.9)	10 (0.9)	34 (3.1)	
Hispanic:								
2 - 5	25 (2.0)	28 (1.7)	35 (1.5)	21 (1.2)	36 (1.4)	24 (1.3)	60 (6.0)	
6 - 11	18 (1.1)	16 (1.5)	23 (1.6)	13 (1.6)	20 (1.7)	16 (1.8)	43 (5.1)	
12 - 19	18 (1.5)	15 (1.1)	26 (1.1)	11 (0.7)	20 (1.2)	15 (0.8)	37 (3.9)	
20 and over	15 (0.6)	14 (0.5)	26 (0.7)	11 (0.5)	19 (0.5)	13 (0.5)	30 (1.8)	74 (4.9)
2 and over	17 (0.5)	15 (0.5)	26 (0.7)	12 (0.5)	20 (0.5)	14 (0.4)	31 (1.8)	

Symbol Legend

* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages and ratios are as follows:

Percent reporting: An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 1.98.

Nutrient ratios expressed as percentages: An estimated ratio between 25 and 75 percent is flagged when based on a sample size n^* of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect and n^* is the number of individuals in the sample reporting non-zero intake of the respective nutrient. An estimated ratio less than or equal to 25 percent or greater than or equal to 75 percent, is flagged when the smaller of n^*p and n^* (1-p) is less than 8 times the VIF, where p is the percentage expressed as a fraction. Additionally, an estimated ratio is flagged when either the relative standard error or p/(1-p) times the relative standard error is greater than 30 percent. The VIF used in this table is 1.98.

Footnotes

- ¹ Snack occasions were reported as distinct eating occasions during the dietary interview and consisted of one or more food and beverage items, including plain water. Water was the only item reported in approximately 25 percent of the snack occasions. Survey respondents selected the name of all eating occasions from a fixed list that was provided during the interview. All reports of "snack", "drink" or "extended consumption" (items that were consumed over a long period of time) were included as snack occasions. Spanish language interviewers used Spanish language snack occasion names: "merienda", "entre comida", "botana", "botana", "botana", "tentempie", and "bebida".
- ² Percentages are estimated as a ratio of total nutrients from snacks for all individuals to total daily nutrient intakes for all individuals. Total daily nutrient intakes are available from: www.ars.usda.gov/nea/bhnrc/fsrg. See Table 2. Nutrient Intakes from Food and Beverages: Mean Amounts Consumed per Individual, by Race/Ethnicity and Age, in the United States, 2013-2014.
- ³ The percentage of respondents in the race/ethnicity/age group who reported consuming at least one item at an eating occasion designated as snack.
- ⁴ A new sampling methodology was implemented for NHANES 2011-2012 to oversample non-Hispanic non-black Asians in addition to the ongoing oversample of Hispanics, non-Hispanic blacks, low-income whites/others, and whites/others 80+ years.

Abbreviations

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2013-2014

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2013-2014. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2013-2014 www.ars.usda.gov/nea/bhnrc/fsrg.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

Suggested Citation

U.S. Department of Agriculture, Agricultural Research Service. 2016. Snacks: Percentages of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Race/Ethnicity and Age, What We Eat in America. NHANES 2013-2014.