**Table 18. Lunch¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2015-2016

Race/ethnicity	Percent		Carbo- Total Dietar		Dietary	Total	Saturated	Mono- unsaturated	Poly- unsaturated	
and age	reporting <sup>3</sup>	Energy	Protein	hydrate	sugars	fiber	fat	fat	fat	fat
(years)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)
<b>.</b>	70 (82)	,, (22)	,, (22)	70 (82)	70 (82)	, (SZ)	70 (32)	,, (22)	70 (82)	,, (22)
Non-Hispanic White:										
2 - 5	98* (0.6)	26 (1.5)	29 (1.6)	24 (1.6)	18 (1.8)	29 (1.8)	29 (1.6)	27 (1.7)	30 (2.1)	31 (1.4)
6 - 11	94* (2.4)	27 (1.4)	31 (1.1)	26 (1.4)	24 (1.8)	28 (1.5)	28 (1.6)	26 (1.9)	29 (1.6)	30 (1.4)
12 - 19	80 (3.3)	27 (1.6)	32 (2.4)	26 (1.5)	23 (1.5)	30 (1.8)	27 (1.7)	26 (1.7)	27 (1.5)	29 (2.2)
20 and over	81 (1.3)	25 (0.8)	29 (0.8)	23 (0.7)	18 (0.8)	26 (1.0)	27 (0.9)	25 (1.0)	27 (0.9)	29 (0.9)
2 and over	83 (1.2)	25 (0.7)	29 (0.7)	23 (0.7)	19 (0.8)	26 (1.0)	27 (0.8)	26 (0.9)	27 (0.9)	29 (0.8)
Non-Hispanic Black:										
2 - 5	89* (2.6)	26 (1.3)	29 (1.5)	24 (1.1)	20 (1.2)	28 (1.8)	27 (1.8)	26 (1.7)	27 (1.8)	27 (2.8)
6 - 11	88 (2.1)	30 (1.6)	35 (1.3)	28 (1.7)	24 (2.2)	33 (1.9)	31 (1.8)	32 (2.3)	30 (1.7)	30 (2.0)
12 - 19	75 (2.4)	27 (1.7)	29 (1.8)	25 (1.7)	21 (1.6)	29 (2.3)	28 (2.0)	26 (1.7)	28 (1.9)	30 (2.5)
20 and over	71 (2.3)	25 (0.8)	28 (1.1)	23 (0.7)	20 (0.6)	26 (0.8)	26 (1.2)	24 (1.1)	26 (1.2)	28 (1.5)
2 and over	74 (1.7)	25 (0.7)	29 (0.9)	24 (0.7)	21 (0.7)	27 (0.8)	27 (1.0)	25 (1.0)	27 (1.0)	28 (1.3)
Non-Hispanic Asian <sup>4</sup> :										
2 - 5	98* (2.0)	23* (1.6)	23* (1.4)	23* (1.6)	18* (2.2)	31* (3.2)	24* (2.7)	22* (3.3)	24* (3.6)	27* (2.2)
6 - 11	99* (0.7)	25* (1.4)	29* (2.1)	24* (1.3)	17* (1.5)	29* (1.9)	26* (2.2)	22* (3.1)	27* (1.8)	30* (4.0)
12 - 19	91* (2.5)	29 (2.5)	32 (3.2)	28 (2.3)	23 (1.7)	30 (1.9)	30 (2.7)	29 (2.4)	30 (2.7)	33 (3.4)
20 and over	90 (2.1)	27 (0.5)	30 (0.8)	27 (0.6)	18 (1.1)	28 (0.7)	27 (1.0)	26 (1.4)	28 (1.0)	28 (1.0)
2 and over	91 (1.7)	27 (0.6)	30 (0.8)	27 (0.6)	18 (0.8)	29 (0.6)	27 (0.8)	26 (1.1)	28 (0.9)	28 (0.9)
Hispanic:										
2 - 5	85 (2.7)	21 (1.3)	23 (1.8)	20 (1.2)	17 (1.3)	26 (1.8)	21 (1.7)	19 (1.6)	22 (2.1)	22 (2.0)
6 - 11	77 (3.5)	24 (1.3)	27 (1.4)	23 (1.3)	21 (1.5)	26 (2.0)	25 (1.5)	24 (1.7)	25 (1.5)	26 (1.5)
12 - 19	77 (2.5)	26 (1.3)	30 (1.6)	25 (1.2)	22 (1.4)	29 (1.7)	27 (1.6)	26 (1.6)	27 (1.5)	29 (2.0)
20 and over	70 (2.7)	24 (0.8)	27 (1.1)	23 (0.9)	19 (0.9)	25 (1.3)	25 (0.8)	24 (0.9)	26 (0.7)	26 (0.9)
2 and over	73 (1.8)	24 (0.6)	27 (0.9)	23 (0.6)	20 (0.7)	26 (1.0)	25 (0.7)	24 (0.7)	25 (0.6)	26 (0.8)

**Table 18. Lunch¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2015-2016 *(continued)* 

Race/ethnicity and age (years)	Choles- terol % (SE)	Vitamin A (RAE) % (SE)	Beta- carotene % (SE)	Lycopene % (SE)	Thiamin % (SE)	Ribo- flavin % (SE)	Niacin % (SE)	Vitamin B6 % (SE)	Folate (DFE) % (SE)
Non-Hispanic White:									
2 - 5	24 (2.0)	23 (2.9)	35 (9.7)	35 (6.8)	26 (1.9)	22 (1.9)	30 (2.3)	23 (2.2)	24 (2.5)
6 - 11	24 (1.8)	` '	34 (7.2)	25 (7.2)	25 (1.3)	26 (1.6)	28 (1.2)	23 (1.1)	21 (1.4)
12 - 19	27 (3.1)	21 (1.9)	31 (4.5)	31 (4.7)	28 (2.3)	23 (2.1)	29 (2.5)	24 (2.1)	24 (1.7)
20 and over	26 (1.1)	21 (1.1)	29 (2.8)	33 (2.9)	26 (0.6)	20 (0.6)	27 (0.7)	24 (0.6)	23 (0.4)
2 and over	26 (0.9)	21 (1.0)	29 (2.4)	33 (2.5)	26 (0.7)	21 (0.6)	28 (0.7)	23 (0.7)	23 (0.4)
Non-Hispanic Black:									
2 - 5	22 (1.8)	23 (2.8)	40 (9.9)	34 (6.6)	24 (1.5)	22 (1.5)	29 (1.6)	23 (1.8)	21 (1.4)
6 - 11	29 (2.7)	29 (2.9)	41 (6.2)	31 (4.8)	28 (1.3)	29 (1.3)	29 (1.0)	26 (1.3)	24 (2.2)
12 - 19	25 (2.4)	21 (2.2)	31 (7.4)	29 (4.5)	25 (2.1)	23 (2.5)	28 (2.5)	23 (2.4)	22 (2.8)
20 and over	26 (1.1)	20 (2.1)	24 (4.4)	32 (2.2)	25 (0.8)	22 (0.8)	27 (1.1)	24 (1.2)	24 (0.9)
2 and over	26 (0.7)	21 (1.4)	26 (3.7)	32 (2.2)	25 (0.7)	23 (0.7)	27 (1.0)	24 (1.0)	23 (0.7)
Non-Hispanic Asian <sup>4</sup> :									
2 - 5	28* (8.1)	25* (5.1)	64* (8.0)	37* (18.3)	19* (2.2)	15* (1.5)	21* (2.4)	18* (0.9)	24* (4.0)
6 - 11	27* (2.6)	28* (4.1)	43* (8.8)	31*(10.1)	28* (1.3)	23* (1.9)	30* (2.7)	28* (2.3)	26* (2.7)
12 - 19	28 (3.6)	20 (2.2)	27 (4.7)	43 (9.9)	27 (2.1)	23 (1.9)	31 (2.6)	25 (2.3)	24 (2.4)
20 and over	27 (1.2)	29 (2.0)	36 (3.4)	27 (5.3)	29 (0.8)	23 (0.5)	31 (1.2)	29 (0.9)	30 (1.2)
2 and over	27 (1.1)	28 (1.6)	36 (3.1)	29 (5.0)	28 (0.7)	23 (0.5)	31 (1.1)	28 (0.8)	29 (1.0)
Hispanic:									
2 - 5	18 (2.5)	15 (1.7)	41 (7.3)	33 (4.5)	19 (1.5)	16 (1.5)	23 (2.2)	17 (1.6)	18 (1.9)
6 - 11	22 (1.5)	22 (2.5)	35 (7.3)	30 (5.1)	23 (1.5)	21 (1.3)	25 (1.7)	22 (1.4)	20 (1.6)
12 - 19	24 (1.6)	22 (1.8)	38 (4.9)	32 (2.8)	25 (1.7)	23 (1.5)	27 (2.0)	23 (1.9)	22 (1.6)
20 and over	25 (1.4)	21 (1.3)	30 (2.9)	33 (1.9)	24 (0.9)	20 (0.8)	26 (1.3)	23 (1.3)	23 (1.1)
2 and over	24 (1.3)	21 (0.9)	32 (1.9)	32 (1.5)	24 (0.7)	20 (0.7)	26 (0.9)	23 (0.9)	22 (0.8)

**Table 18. Lunch¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2015-2016 *(continued)* 

Race/ethnicity										min E oha-								
and age	Choline		Vitam	in B12	Vita	min C		nin D	tocopherol)		Vitamin K		Cal	cium	Phos	phorus	Magr	nesium
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Non-Hispanic White:																		
2 - 5	25	(1.7)	22	(2.4)	19	(3.1)	19	(2.2)	31	(2.1)	36	(5.6)	23	(1.3)	27	(1.3)	27	(2.2)
6 - 11	27	(1.2)	22	(1.6)	20	(2.6)	27	(1.9)	29	(1.7)	27	(3.0)	29	(2.2)	30	(1.4)	28	(1.4)
12 - 19	27	(2.4)	22	(1.7)	24	(2.1)	20	(2.5)	27	(1.9)	32	(3.4)	28	(2.3)	30	(2.1)	27	(1.8)
20 and over	24	(0.8)	24	(1.6)	22	(0.9)	17	(1.4)	25	(0.9)	30	(1.9)	24	(0.8)	26	(0.6)	22	(0.8)
2 and over	24	(0.7)	24	(1.4)	22	(0.9)	18	(1.4)	25	(0.8)	30	(1.7)	25	(0.8)	27	(0.6)	23	(0.7)
Non-Hispanic Black:																		
2 - 5	25	(1.7)	23	(2.3)	21	(1.9)	21	(1.9)	27	(2.1)	24	(3.8)	25	(1.6)	27	(1.4)	26	(1.7)
6 - 11	30	(2.2)	28	(2.0)	23	(3.1)	33	(2.2)	28	(2.3)	30	(2.7)	36	(2.7)	35	(1.8)	32	(1.7)
12 - 19	25	(2.3)	21	(2.6)	20	(2.7)	15	(3.0)	27	(2.5)	30	(7.2)	26	(2.0)	28	(2.3)	26	(1.8)
20 and over	25	(1.1)	26	(1.6)	25	(1.5)	22	(3.2)	24	(1.4)	24	(2.6)	22	(0.7)	26	(0.9)	22	(0.8)
2 and over	26	(0.8)	25	(1.2)	23	(1.3)	22	(2.1)	25	(1.2)	25	(2.0)	24	(0.8)	27	(0.9)	24	(0.7)
Non-Hispanic Asian <sup>4</sup> :																		
2 - 5	24*	(4.7)	15*	(2.4)	34*	(5.1)	11*	(2.4)	23*	(1.1)	38*	(5.3)	17*	(2.9)	20*	(1.4)	20*	(0.8)
6 - 11	26*	(1.5)	28*	(5.8)	26*	(4.3)	17*	(3.1)	28*	(2.1)	39*	(4.7)	24*	(2.8)	27*	(1.6)	26*	(1.4)
12 - 19	28	(3.4)	27	(3.3)	21	(2.8)	20	(2.7)	27	(3.5)	30	(4.0)	25	(2.0)	29	(2.3)	27	(2.1)
20 and over	27	(0.8)	27	(1.5)	27	(1.0)	22	(2.4)	27	(1.1)	34	(2.3)	24	(1.3)	28	(0.6)	25	(0.4)
2 and over	27	(0.9)	26	(1.4)	26	(0.8)	21	(2.0)	27	(0.8)	34	(2.2)	24	(0.9)	28	(0.5)	25	(0.5)
Hispanic:																		
2 - 5	19	(2.0)	13	(2.4)	26	(3.2)	10	(2.1)	23	(1.7)	30	(6.2)	17	(1.7)	20	(1.6)	21	(1.7)
6 - 11	24	(1.1)	21	(1.7)	23	(2.4)	18	(1.8)	25	(1.5)	28	(3.4)	23	(0.9)	25	(1.2)	25	(1.3)
12 - 19	25	(1.4)	23	(1.5)	22	(2.0)	22	(2.4)	27	(1.7)	36	(3.4)	26	(1.7)	28	(1.4)	27	(1.4)
20 and over	24	(1.2)	22	(1.5)	23	(1.6)	19	(1.7)	24	(1.0)	29	(2.6)	20	(0.6)	25	(0.9)	21	(1.1)
2 and over	24	(1.0)	21	(1.0)	23	(1.3)	18	(1.1)	24	(0.7)	30	(2.0)	21	(0.6)	25	(0.8)	22	(0.9)

**Table 18. Lunch¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, in the United States, 2015-2016 *(continued)* 

Race/ethnicity																
and age	Īι	on	7:	inc	Co	pper	Sele	nium	Pota	ssium	Soc	lium	Caf	feine	Alc	ohol
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Non-Hispanic White:																
2 - 5	24	(1.6)	26	(2.0)	27	(1.9)	30	(1.7)	25	(2.0)	32	(1.9)	32	(7.6)		
6 - 11	21	(1.4)	25	(1.6)	26	(1.7)	31	(1.8)	29	(1.4)	30	(1.2)	19	(2.5)		
12 - 19	24	(1.9)	26	(1.7)	26	(1.7)	31	(2.4)	29	(2.0)	34	(2.4)	26	(4.3)		
20 and over	23	(0.5)	25	(0.7)	21	(0.7)	30	(1.0)	24	(0.7)	31	(0.9)	9	(0.9)	6	(0.8)
2 and over	23	(0.5)	25	(0.6)	22	(0.7)	30	(0.9)	25	(0.7)	31	(0.9)	9	(0.9)		
Non-Hispanic Black:																
2 - 5	22	(1.1)	25	(1.5)	26	(1.7)	30	(2.0)	27	(1.7)	30	(1.2)	15*	(5.7)		
6 - 11	25	(1.5)	30	(1.9)	31	(1.7)	35	(1.7)	32	(2.0)	33	(1.9)	18*	(6.7)		
12 - 19	23	(2.5)	25	(2.2)	24	(1.9)	28	(1.7)	27	(1.9)	29	(1.8)	27	(4.9)		
20 and over	24	(1.2)	28	(1.2)	23	(1.0)	28	(0.9)	25	(1.0)	28	(1.1)	12	(1.4)	4*	(1.2)
2 and over	24	(0.9)	27	(1.0)	24	(0.8)	29	(0.9)	26	(0.9)	29	(1.1)	12	(1.3)		
Non-Hispanic Asian <sup>4</sup> :																
2 - 5	19*	(2.4)	20*	(2.2)	20*	(1.1)	24*	(2.3)	21*	(1.4)	29*	(2.6)	0*	(0.0)		
6 - 11	25*	(3.1)	25*	(2.3)	24*	(2.0)	30*	(1.9)	26*	(0.9)	30*	(1.4)	15*	(4.8)		
12 - 19	26	(2.3)	28	(3.6)	27	(2.2)	33	(3.0)	28	(2.4)	33	(2.9)	7*	(3.0)		
20 and over	29	(0.9)	29	(0.9)	25	(0.7)	30	(0.8)	27	(0.6)	32	(0.9)	11	(1.3)	3*	(1.2)
2 and over	28	(0.8)	29	(1.0)	25	(0.6)	31	(0.9)	27	(0.6)	32	(1.0)	11	(1.3)		
Hispanic:																
2 - 5	17	(1.8)	19	(2.1)	21	(1.6)	23	(1.6)	22	(1.6)	26	(1.2)	25*	(11.0)		
6 - 11	20	(2.0)	24	(1.1)	25	(1.4)	29	(3.0)	26	(1.2)	27	(1.6)	20	(4.0)		
12 - 19	23	(1.6)	27	(1.5)	26	(1.3)	30	(1.7)	28	(1.6)	30	(1.7)	13	(2.4)		
20 and over	23	(0.8)	26	(1.0)	22	(1.1)	27	(1.1)	24	(1.0)	28	(0.9)	11	(1.3)	7*	(2.0)
2 and over	22	(0.7)	25	(0.8)	23	(0.9)	28	(0.9)	25	(0.8)	28	(0.7)	12	(1.3)		

# **Symbol Legend**

\* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages and ratios are as follows:

**Percent reporting:** An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 2.41.

**Nutrient ratios expressed as percentages:** An estimated ratio between 25 and 75 percent is flagged when based on a sample size  $n^*$  of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect and  $n^*$  is the number of individuals in the sample reporting non-zero intake of the respective nutrient. An estimated ratio less than or equal to 25 percent or greater than or equal to 75 percent, is flagged when the smaller of  $n^*p$  and  $n^*$  (1-p) is less than 8 times the VIF, where p is the percentage expressed as a fraction. Additionally, an estimated ratio is flagged when either the relative standard error or p/(1-p) times the relative standard error is greater than 30 percent. The VIF used in this table is 2.41.

#### **Footnotes**

- <sup>1</sup> Lunch includes eating occasions designated by the respondent as "brunch", "lunch" or the Spanish equivalent "comida." Please note these eating occasions include consumption of beverages including water.
- <sup>2</sup> Percentages are estimated as a ratio of total nutrients from lunch for all individuals to total daily nutrient intakes for all individuals. Total daily nutrient intakes are available from: <a href="https://www.ars.usda.gov/nea/bhnrc/fsrg">www.ars.usda.gov/nea/bhnrc/fsrg</a>. See Table 2. Nutrient Intakes from Food and Beverages: Mean Amounts Consumed per Individual, by Race/Ethnicity and Age, in the United States, 2015-2016.
- <sup>3</sup> The percentage of respondents in the race/ethnicity/age group who reported consuming at least one item at an eating occasion designated as lunch.
- <sup>4</sup> A new sampling methodology was implemented for NHANES 2011-2012 to oversample non-Hispanic non-black Asians in addition to the ongoing oversample of Hispanics, non-Hispanic blacks, low-income whites/others, and whites/others 80+ years.

## **Abbreviations**

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

### Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2015-2016

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2015-2016. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2015-2016 <a href="www.ars.usda.gov/nea/bhnrc/fsrg">www.ars.usda.gov/nea/bhnrc/fsrg</a>.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

## **Suggested Citation**

U.S. Department of Agriculture, Agricultural Research Service. 2018. Lunch: Percentages of Selected Nutrients Contributed by Food and Beverages Consumed at Lunch, by Race/Ethnicity and Age, What We Eat in America, NHANES 2015-2016.