**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-2018

Race/ethnicity and age	Perce reporti		Ene	ergy	Pro	otein		rbo- rate		otal gars	Die fil	etary ber		otal at		rated at	unsat	ono- urated at	unsat	oly- urated at
(years)	% (	(SE)	%	(SE)	%	(SE)	<u>%</u>	(SE)	<u></u> %	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Non-Hispanic White:		1																		
2 - 5	98* (	(1.3)	19	(1.4)	18	(1.2)	21	(1.6)	23	(2.2)	19	(1.9)	16	(1.5)	18	(2.2)	14	(1.6)	13	(1.2)
6 - 11	90 (	(2.5)	17	(0.8)	16	(1.2)	19	(1.0)	20	(1.4)	17	(1.0)	15	(0.8)	18	(1.0)	15	(0.8)	12	(0.8)
12 - 19	75 (	(3.2)	17	(1.2)	15	(1.0)	19	(1.4)	22	(1.8)	19	(1.9)	15	(1.3)	16	(1.3)	15	(1.3)	13	(1.6)
20 and over	87 (	(1.0)	16	(0.5)	16	(0.6)	18	(0.8)	19	(1.0)	19	(0.9)	15	(0.4)	16	(0.5)	16	(0.5)	14	(0.6)
2 and over	87 (	(1.0)	16	(0.4)	16	(0.5)	18	(0.6)	20	(0.7)	19	(0.7)	15	(0.4)	16	(0.4)	16	(0.4)	14	(0.5)
Non-Hispanic Black:		}																		
2 - 5	89* (	(4.9)	18	(1.7)	17*	(1.8)	20	(1.0)	21	(0.8)	21	(2.0)	16*	(2.9)	18*	(3.0)	16*	(3.5)	13*	(2.2)
6 - 11	86 (	(4.0)	20	(1.6)	18	(1.5)	22	(1.7)	23	(2.0)	20	(1.7)	18	(1.8)	18	(1.9)	19	(2.0)	15	(2.0)
12 - 19	65 (	(3.6)	14	(1.0)	12	(1.0)	15	(1.1)	17	(1.6)	13	(1.1)	12	(1.0)	14	(1.3)	12	(1.0)	10	(1.2)
20 and over	72 (	(1.9)	17	(0.7)	15	(0.8)	18	(0.5)	20	(0.4)	17	(1.0)	16	(0.9)	17	(1.0)	17	(1.1)	15	(0.9)
2 and over	73 (	(1.9)	16	(0.7)	15	(0.7)	18	(0.5)	20	(0.5)	17	(0.8)	16	(0.9)	17	(1.0)	16	(1.0)	14	(0.9)
Non-Hispanic Asian <sup>4</sup> :																				
2 - 5	97* (	(2.0)	18*	(2.6)	19*	(2.4)	18*	(2.0)	21*	(2.4)	16*	(1.5)	19*	(4.1)	21*	(3.7)	18*	(4.9)	16*	(4.7)
6 - 11	92* (	(3.6)	19*	(1.2)	18*	(1.4)	19*	(1.3)	19*	(1.4)	14*	(1.8)	19*	(1.4)	20*	(1.3)	18*	(1.6)	18*	(2.4)
12 - 19	78 (	(4.1)	20	(1.3)	19	(1.7)	21	(1.6)	22	(3.9)	17*	(1.5)	20	(1.3)	21	(1.5)	20	(1.6)	18	(2.5)
20 and over	83 (	(1.8)	18	(0.5)	18	(0.7)	21	(0.6)	22	(1.2)	21	(0.8)	17	(0.6)	18	(0.5)	16	(0.6)	16	(1.2)
2 and over	84 (	(1.5)	19	(0.4)	18	(0.7)	20	(0.5)	22	(1.2)	20	(0.7)	17	(0.6)	19	(0.5)	16	(0.6)	16	(1.2)
Hispanic:																				
2 - 5	97* (	(2.9)	21	(1.2)	23	(1.6)	21	(1.2)	23	(1.4)	21	(2.0)	21	(1.6)	24	(1.9)	21	(1.9)	16*	(1.9)
6 - 11	91* (	(2.5)	21	(1.7)	23	(1.8)	22	(1.7)	23	(1.7)	22	(2.0)	20	(1.8)	20	(1.6)	19	(1.7)	19	(2.2)
12 - 19	74 (	(2.8)	19	(1.4)	18	(1.5)	20	(1.4)	22	(1.6)	20	(1.8)	17	(1.7)	19	(1.9)	17	(1.7)	15	(1.7)
20 and over	88 (	(1.8)	24	(0.9)	25	(0.9)	25	(0.9)	25	(1.3)	26	(0.9)	25	(1.2)	25	(1.3)	25	(1.1)	24	(1.4)
2 and over	87 (	(1.4)	23	(0.8)	24	(0.8)	24	(0.8)	24	(0.9)	24	(0.8)	23	(1.0)	24	(1.1)	23	(0.9)	22	(1.1)

**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-2018 *(continued)* 

Race/ethnicity and age (years)	Choles terol % (SI	(RAE)	Beta- carotene % (SE)	Lycopene % (SE)	Thiamin % (SE)	Ribo- flavin % (SE)	Niacin % (SE)	Vitamin B6 % (SE)	Folate (DFE) % (SE)
Non-Hispanic White:									
2 - 5	29 (3.9	9) 37 (3.1)	7* (2.8)	1* (0.6)	30 (1.6)	32 (2.0)	23 (1.1)	31 (1.5)	36 (3.4)
6 - 11	23 (3.2	, , ,	3* (0.5)	4* (3.4)	30 (1.7)	31 (1.6)	24 (1.1)	31 (1.5)	37 (2.2)
12 - 19	19 (2.	, , ,	9* (3.2)	6* (4.5)	25 (1.4)	28 (1.9)	17 (1.0)	21 (1.8)	28 (1.7)
20 and over	28 (1		7 (2.0)	5 (1.1)	23 (1.0)	28 (1.0)	17 (0.8)	21 (1.2)	28 (0.9)
2 and over	27 (1.	1) 27 (0.9)	7 (1.8)	5 (1.2)	24 (0.8)	29 (0.9)	18 (0.7)	22 (1.0)	29 (0.9)
Non-Hispanic Black:									
2 - 5	21 (2.4	4) 40 (3.7)	6* (2.7)	12* (7.2)	33 (2.4)	33 (2.3)	26 (1.7)	35 (1.9)	37 (3.7)
6 - 11	24 (3	3) 41 (2.1)	9* (3.3)	8* (5.1)	32 (1.9)	34 (2.4)	25 (1.9)	35 (2.6)	39 (3.9)
12 - 19	23 (3.0	6) 33 (3.0)	4* (1.1)	3* (1.6)	26 (2.6)	27 (2.8)	18 (2.0)	27 (3.1)	27 (3.4)
20 and over	28 (2.2	2) 26 (1.6)	6* (2.0)	6* (1.9)	23 (0.8)	27 (1.0)	18 (0.8)	23 (1.2)	25 (1.0)
2 and over	27 (2.2	2) 28 (1.5)	6 (1.5)	6* (1.9)	25 (0.6)	28 (0.9)	19 (0.6)	25 (1.1)	27 (0.7)
Non-Hispanic Asian <sup>4</sup> :									
2 - 5	27* (4.5	5) 36* (3.7)	6* (2.7)	2* (2.1)	26* (2.1)	30* (2.4)	24* (2.5)	30* (4.7)	33* (2.5)
6 - 11	29* (4.0	0) 33* (3.5)	4* (1.6)	3* (1.5)	24* (1.6)	29* (1.6)	21* (2.1)	27* (2.6)	26* (2.3)
12 - 19	27 (2.9	9) 32 (5.5)	23* (13.3)	3* (1.6)	25 (2.1)	29 (2.8)	19 (1.8)	23 (3.6)	27 (2.6)
20 and over	24 (1.0	6) 27 (2.2)	16 (4.1)	6* (2.1)	23 (1.4)	27 (1.5)	17 (1.1)	20 (1.6)	22 (1.5)
2 and over	25 (1.:	5) 28 (1.7)	16 (3.8)	6* (1.9)	23 (1.2)	27 (1.3)	18 (0.9)	21 (1.4)	23 (1.1)
Hispanic:									
2 - 5	31 (4.	7) 38 (3.3)	11* (3.7)	2* (1.0)	30 (1.9)	33 (2.0)	25 (2.1)	32 (2.7)	35 (3.5)
6 - 11	29 (2.3	8) 37 (1.8)	8* (1.9)	11* (4.2)	30 (1.8)	32 (2.1)	27 (2.2)	33 (2.0)	31 (2.1)
12 - 19	26 (3.0	6) 37 (3.5)	6* (1.3)	9* (3.5)	29 (2.1)	31 (2.8)	22 (2.4)	27 (2.9)	33 (2.6)
20 and over	39 (1	5) 35 (2.0)	22 (3.3)	21 (4.1)	29 (1.1)	33 (1.6)	24 (1.3)	25 (1.6)	30 (1.3)
2 and over	36 (1	3) 36 (1.5)	20 (3.1)	18 (3.3)	29 (0.8)	33 (1.0)	24 (1.0)	27 (1.2)	31 (1.1)

**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-2018 *(continued)* 

Race/ethnicity and age	Che	oline	Vitam	in B12	Vito	min C	Vito	nin D	(alp	min E oha- herol)	Vitor	nin K	Cal	cium	Phos	phorus	Magr	nesium
(years)	%	(SE)	% Italii	(SE)	% Ttal	(SE)	% rtai	(SE)	тосор %	(SE)	% Hai	(SE)	%	(SE)	%	(SE)	Wiagi	(SE)
Non-Hispanic White:																		
2 - 5	27	(2.6)	37	(2.4)	17	(2.7)	39	(3.5)	15	(1.8)	14*	(5.3)	25	(2.4)	24	(1.9)	22	(2.3)
6 - 11	23	(1.9)	37	(2.5)	22	(3.1)	41	(2.8)	19	(4.7)	8	(1.5)	26	(1.6)	21	(1.2)	19	(0.7)
12 - 19	20	(1.6)	28	(2.7)	25	(2.8)	41	(4.5)	18	(2.5)	11	(1.2)	24	(2.1)	20	(1.5)	19	(1.3)
20 and over	24	(0.8)	22	(1.3)	20	(1.0)	36	(1.7)	20	(1.2)	9	(1.1)	22	(1.0)	20	(0.6)	20	(0.6)
2 and over	24	(0.7)	24	(1.2)	20	(1.0)	37	(1.6)	19	(1.0)	9	(1.0)	22	(0.9)	20	(0.5)	20	(0.5)
Non-Hispanic Black:																		
2 - 5	23	(2.0)	41	(2.0)	20	(2.6)	42	(2.6)	15*	(2.0)	14*	(3.9)	29	(2.5)	24	(2.3)	22	(1.6)
6 - 11	25	(2.3)	40	(2.6)	26	(2.3)	48	(3.2)	16	(1.7)	12*	(4.3)	28	(2.1)	24	(1.8)	21	(1.7)
12 - 19	21	(2.7)	29	(3.1)	15	(2.5)	42	(3.9)	12	(1.3)	8*	(1.2)	23	(2.2)	18	(1.4)	14	(0.8)
20 and over	24	(1.6)	25	(1.4)	20	(1.3)	36	(2.1)	17	(0.9)	9	(1.2)	23	(0.9)	19	(0.8)	18	(0.7)
2 and over	24	(1.5)	27	(1.1)	20	(1.0)	38	(2.0)	16	(0.8)	9	(0.9)	23	(1.0)	20	(0.8)	18	(0.6)
Non-Hispanic Asian <sup>4</sup> :																		
2 - 5	24*	(2.7)	40*	(2.8)	18*	(4.9)	37*	(3.6)	15*	(3.2)	10*	(3.9)	27*	(2.5)	23*	(2.7)	20*	(2.0)
6 - 11	26*	(2.7)	33*	(4.2)	12*	(3.8)	37*	(5.9)	15*	(1.5)	11*	(1.9)	25*	(2.4)	24*	(1.6)	18*	(1.0)
12 - 19	25	(2.7)	29	(6.0)	21*	(6.7)	36	(9.6)	19	(1.8)	8*	(2.9)	29	(2.7)	23	(1.9)	20	(1.3)
20 and over	23	(1.0)	25	(3.1)	15	(1.9)	31	(2.1)	17	(1.1)	12	(1.9)	26	(1.2)	21	(0.8)	21	(0.8)
2 and over	23	(1.1)	27	(2.7)	15	(2.1)	32	(1.8)	17	(0.9)	11	(1.6)	26	(1.1)	21	(0.7)	21	(0.7)
Hispanic:																		
2 - 5	29	(3.1)	37	(2.7)	20	(2.5)	41	(4.0)	19	(2.5)	13*	(1.4)	31	(2.5)	27	(1.9)	25	(1.5)
6 - 11	28	(1.7)	37	(2.0)	26	(2.3)	44	(2.0)	20	(2.1)	14	(2.4)	30	(1.9)	26	(1.7)	24	(2.3)
12 - 19	25	(2.4)	32	(3.6)	24	(2.0)	46	(4.6)	18	(2.5)	12	(1.7)	26	(2.1)	23	(1.8)	22	(1.4)
20 and over	33	(1.2)	27	(1.9)	26	(2.0)	42	(2.5)	26	(1.1)	22	(2.3)	29	(1.0)	28	(1.0)	27	(1.0)
2 and over	32	(1.0)	29	(1.3)	25	(1.5)	43	(1.7)	24	(1.0)	20	(2.0)	29	(0.9)	27	(0.8)	26	(0.8)

**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-2018 (continued)

Race/ethnicity and age		on (GE)		inc		pper		nium		ssium		lium		feine		cohol
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	<u>%</u>	(SE)	%	(SE)	%	(SE)	%	(SE)
Non-Hispanic White:																
2 - 5	36	(2.9)	27	(1.6)	18	(2.3)	18	(1.4)	21	(1.9)	15	(0.8)	18	(2.8)		
6 - 11	34	(2.0)	28	(1.1)	15	(0.7)	16	(1.5)	19	(0.9)	14	(0.8)	14	(2.1)		
12 - 19	26	(1.7)	19	(1.6)	16	(1.6)	13	(0.8)	19	(1.2)	13	(0.7)	24	(3.9)		
20 and over	26	(1.3)	19	(1.0)	17	(0.6)	17	(0.7)	20	(0.7)	13	(0.5)	45	(2.0)	1*	(0.5)
2 and over	27	(1.1)	20	(0.9)	16	(0.5)	17	(0.6)	20	(0.6)	13	(0.5)	44	(2.0)		
Non-Hispanic Black:																
2 - 5	36	(2.1)	31	(2.6)	18*	(1.9)	17*	(1.8)	20	(1.8)	16*	(2.1)	6*	(3.6)		
6 - 11	35	(1.9)	28	(2.8)	17	(1.2)	17	(2.3)	21	(1.6)	17	(1.3)	11*	(2.3)		
12 - 19	27	(2.9)	24	(2.7)	12	(1.0)	11	(1.0)	14	(1.3)	13	(0.8)	14*	(1.8)		
20 and over	25	(1.0)	19	(0.7)	16	(0.8)	17	(0.9)	18	(0.7)	15	(0.7)	47	(2.2)	2*	(1.7)
2 and over	27	(0.8)	21	(0.7)	16	(0.7)	16	(0.9)	18	(0.7)	15	(0.6)	45	(2.2)		
Non-Hispanic Asian <sup>4</sup> :																
2 - 5	35*	(3.0)	33*	(2.9)	15*	(2.2)	17*	(2.7)	19*	(1.5)	15*	(2.8)	33*	(12.0)		
6 - 11	28*	(2.4)	23*	(2.0)	14*	(1.8)	18*	(2.0)	18*	(0.8)	17*	(2.1)	22*	(6.5)		
12 - 19	27	(3.5)	21	(2.1)	16*	(0.9)	19	(2.3)	19	(1.8)	19	(2.5)	31	(9.3)		
20 and over	25	(2.0)	20	(1.0)	18	(0.7)	19	(0.9)	20	(0.7)	16	(1.1)	42	(2.4)	0*	(0.0)
2 and over	25	(1.6)	21	(0.9)	18	(0.6)	19	(0.8)	20	(0.6)	16	(0.9)	41	(2.3)		
Hispanic:																
2 - 5	32	(2.2)	31	(2.3)	19	(1.6)	22	(2.0)	24	(1.4)	20	(1.6)	30*	(10.0)		
6 - 11	31	(2.2)	28	(2.2)	20	(2.1)	23	(2.3)	24	(1.7)	21	(1.7)	33*	(13.6)		
12 - 19	29	(1.7)	24	(2.2)	19	(1.3)	18	(1.6)	22	(1.5)	17	(1.6)	31	(5.1)		
20 and over	30	(1.2)	25	(1.2)	25	(1.2)	28	(1.2)	27	(0.9)	24	(1.1)	52	(2.4)	2*	(0.8)
2 and over	30	(0.9)	26	(0.9)	23	(0.9)	26	(1.0)	26	(0.8)	22	(1.0)	50	(2.2)		

# **Symbol Legend**

\* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages and ratios are as follows:

**Percent reporting:** An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 2.57.

**Nutrient ratios expressed as percentages:** An estimated ratio between 25 and 75 percent is flagged when based on a sample size  $n^*$  of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect and  $n^*$  is the number of individuals in the sample reporting non-zero intake of the respective nutrient. An estimated ratio less than or equal to 25 percent or greater than or equal to 75 percent, is flagged when the smaller of  $n^*p$  and  $n^*$  (1-p) is less than 8 times the VIF, where p is the percentage expressed as a fraction. Additionally, an estimated ratio is flagged when either the relative standard error or p/(1-p) times the relative standard error is greater than 30 percent. The VIF used in this table is 2.57.

### **Footnotes**

- <sup>1</sup> Breakfast includes eating occasions designated by the respondent as "breakfast", or the Spanish equivalents "desayuno", and "almuerzo." Please note these eating occasions include consumption of beverages including water.
- <sup>2</sup> Percentages are estimated as a ratio of total nutrients from breakfast for all individuals to total daily nutrient intakes for all individuals. Total daily nutrient intakes are available from: <a href="https://www.ars.usda.gov/nea/bhnrc/fsrg">www.ars.usda.gov/nea/bhnrc/fsrg</a>. See Table 2. Nutrient Intakes from Food and Beverages: Mean Amounts Consumed per Individual, by Race/Ethnicity and Age, in the United States, 2017-2018.
- <sup>3</sup> The percentage of respondents in the race/ethnicity/age group who reported consuming at least one item at an eating occasion designated as breakfast.
- <sup>4</sup> A new sampling methodology was implemented for NHANES 2011-2012 to oversample non-Hispanic non-black Asians in addition to the ongoing oversample of Hispanics, non-Hispanic blacks, low-income whites/others, and whites/others 80+ years.

## **Abbreviations**

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

### Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2017-2018

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2017-2018. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2017-2018 www.ars.usda.gov/nea/bhnrc/fsrg.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

## **Suggested Citation**

U.S. Department of Agriculture, Agricultural Research Service. 2020. Breakfast: Percentages of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, What We Eat in America, NHANES 2017-2018.