**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic

Race/ethnicity and age (years)		Percent reporting <sup>3</sup> % (SE)		ergy (SE)	Pro %	otein (SE)		rbo- rate (SE)	Total sugars % (SE)		Dietary fiber % (SE)		Total fat % (SE)		Saturated fat % (SE)		Mono- unsaturated fat % (SE)		unsat	oly- urated at (SE)
· /	,,,	(~=)	%	(3-)		(~ _)		(32)		(3-)	,,,	(3-)		(3-)		(~=)		(3-)		(3-)
Non-Hispanic White:	07*	(1.0)	10	(0,0)	10	(1.0)	21	(1.0)	22	(1.5)	10	(1.0)	17	(1.1)	10	(1.4)	1.0	(1.0)	1.5	(1.0)
2 - 5	97*	(1.2)	19	(0.9)	19	(1.0)	21	(1.0)	23	(1.5)	18	(1.3)	17	(1.1)	19	(1.4)	16	(1.3)	15	(1.3)
6 - 11 12 - 19	90 73	(2.2)	18 17	(0.8)	16	(0.9)	20	(1.0)	21 22	(1.3)	18 19	(1.1)	16 14	(1.0)	18 16	(1.3)	16 14	(1.0)	13 13	(0.8)
12 - 19 20 and over	73 87	(2.9)		(1.0)	14 16	(0.6)	19 18	(1.1)	20	(1.5)	19	(2.1)		(1.3)	17	(1.3)	14	(1.5)	13	(1.7)
20 and over	0/	(0.8)	16	(0.4)	16	(0.4)	10	(0.5)	20	(0.7)	19	(0.6)	16	(0.4)	1 /	(0.5)	10	(0.4)	14	(0.4)
2 and over	86	(0.8)	17	(0.3)	16	(0.4)	19	(0.4)	20	(0.6)	19	(0.5)	16	(0.4)	17	(0.4)	16	(0.4)	14	(0.4)
Non-Hispanic Black:																				
2 - 5	89	(3.4)	18	(1.3)	18	(1.5)	21	(1.1)	23	(1.5)	20	(1.4)	15	(1.7)	17	(1.8)	14	(1.9)	12	(1.5)
6 - 11	80	(3.4)	18	(1.2)	16	(1.1)	20	(1.2)	22	(1.3)	18	(1.3)	16	(1.5)	17	(1.6)	16	(1.7)	13	(1.7)
12 - 19	66	(3.9)	14	(1.3)	13	(1.1)	16	(1.3)	18	(1.7)	13	(1.1)	13	(1.5)	14	(1.3)	13	(1.6)	12	(1.9)
20 and over	73	(1.5)	17	(0.5)	16	(0.7)	18	(0.5)	20	(0.5)	17	(0.7)	17	(0.7)	18	(0.7)	17	(0.8)	15	(0.7)
2 and over	74	(1.5)	16	(0.5)	16	(0.6)	18	(0.4)	20	(0.4)	17	(0.5)	16	(0.7)	17	(0.7)	16	(0.8)	14	(0.7)
Non-Hispanic Asian:																				
2 - 5	97*	(2.3)	19*	(2.8)	19*	(2.9)	19*	(2.3)	23*	(2.5)	18*	(2.6)	19*	(3.9)	21*	(3.2)	19*	(4.8)	17*	(4.7)
6 - 11	93*	(2.1)	19*	(1.2)	19*	(1.3)	19*	(1.0)	20*	(1.3)	13*	(1.5)	20*	(2.0)	21	(2.0)	19*	(2.4)	17*	(2.2)
12 - 19	78	(3.2)	19	(1.3)	18	(1.3)	20	(1.6)	22	(3.9)	16	(1.3)	19	(1.2)	20	(1.3)	18	(1.3)	17	(1.7)
20 and over	83	(1.5)	18	(0.6)	18	(0.8)	20	(0.8)	22	(1.2)	20	(1.0)	17	(0.6)	18	(0.7)	16	(0.7)	16	(1.0)
2 and over	84	(1.3)	19	(0.5)	18	(0.7)	20	(0.7)	22	(1.2)	20	(0.8)	17	(0.5)	19	(0.6)	17	(0.6)	16	(0.9)
Hispanic:																				
2 - 5	98*	(1.8)	24	(1.3)	26	(1.9)	24	(1.5)	24	(1.1)	24	(1.8)	24	(1.2)	26	(1.2)	24	(1.5)	20	(1.8)
6 - 11	89	(2.7)	22	(1.2)	24	(1.3)	23	(1.2)	24	(1.4)	22	(1.4)	21	(1.3)	21	(1.4)	20	(1.3)	20	(1.7)
12 - 19	71	(2.3)	17	(1.3)	17	(1.6)	19	(1.1)	21	(1.1)	18	(1.3)	16	(1.5)	17	(1.7)	16	(1.6)	15	(1.5)
20 and over	87	(1.2)	24	(0.8)	25	(0.8)	25	(0.8)	25	(1.0)	25	(0.9)	24	(0.9)	25	(0.9)	24	(0.9)	24	(1.1)
2 and over	85	(1.1)	23	(0.8)	24	(0.8)	24	(0.7)	24	(0.8)	24	(0.8)	23	(0.9)	23	(0.9)	23	(0.9)	22	(0.9)

**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic (continued)

Race/ethnicity and age (years)	Chol tero			Vitamin A Beta- (RAE) carotene % (SE) % (SE)		tene	Lyco %	opene (SE)	Thia	amin (SE)		Ribo- flavin % (SE)		Niacin % (SE)		Vitamin B6 % (SE)		olate FE) (SE)
Non-Hispanic White:																		
2 - 5	29	(2.7)	35	(2.9)	5*	(1.8)	1*	(0.6)	29	(1.1)	30	(1.0)	21	(1.2)	29	(1.5)	34	(2.2)
6 - 11		(3.3)	32	(2.1)	3*	(0.7)	3*	(1.8)	29	(1.7)	30	(1.6)	23	(1.2)	29	(1.5)	35	(2.1)
12 - 19		(2.1)	28	(2.8)	8*	(2.4)	4*	(2.4)	24	(0.9)	26	(1.6)	17	(0.9)	23	(2.4)	26	(1.5)
20 and over	30	(1.2)	25	(0.9)	7	(1.4)	6	(1.2)	23	(0.8)	28	(0.8)	18	(0.6)	20	(0.8)	26	(0.8)
2 and over	29	(1.1)	26	(0.9)	7	(1.3)	5	(1.1)	24	(0.7)	28	(0.7)	18	(0.5)	21	(0.7)	27	(0.7)
Non-Hispanic Black:																		
2 - 5	23	(2.3)	36	(2.6)	4*	(1.5)	5*	(3.0)	32	(1.6)	34	(1.7)	23	(1.5)	31	(1.7)	34	(2.6)
6 - 11	23	(2.7)	39	(2.3)	6*	(2.5)	5*	(2.9)	29	(1.4)	31	(1.8)	22	(1.5)	30	(1.9)	32	(2.7)
12 - 19	24	(3.1)	29	(2.1)	5*	(1.1)	6*	(2.0)	22	(2.0)	24	(1.9)	15	(1.4)	22	(1.9)	22	(2.3)
20 and over	29	(1.7)	25	(1.0)	6	(1.0)	5	(1.4)	23	(0.8)	26	(0.9)	17	(0.7)	21	(1.2)	23	(0.8)
2 and over	28	(1.7)	27	(1.0)	6	(0.8)	5	(1.3)	23	(0.6)	27	(0.8)	17	(0.5)	22	(1.0)	24	(0.6)
Non-Hispanic Asian:																		
2 - 5	24*	(3.6)	36*	(3.7)	5*	(1.9)	3*	(1.9)	28*	(2.4)	32*	(2.6)	26*	(3.3)	31*	(4.5)	32*	(2.6)
6 - 11	32	(3.3)	31	(3.3)	5*	(1.3)	3*	(1.4)	24	(1.6)	29	(1.7)	21	(1.7)	27	(2.5)	25	(2.4)
12 - 19	29	(2.5)	31	(4.7)	14*	(7.8)	2*	(1.3)	25	(2.1)	29	(3.0)	18	(1.5)	23	(3.4)	26	(3.0)
20 and over	24	(1.4)	27	(2.3)	15	(3.6)	6	(1.2)	23	(2.1)	27	(2.1)	18	(1.6)	20	(2.1)	22	(1.9)
2 and over	25	(1.4)	28	(1.8)	14	(3.4)	5	(1.2)	23	(1.7)	28	(1.7)	18	(1.3)	21	(1.8)	23	(1.5)
Hispanic:																		
2 - 5	40	(3.9)	39	(2.7)	27	(7.6)	5*	(1.8)	32	(2.1)	33	(1.6)	26	(2.4)	32	(2.4)	36	(2.9)
6 - 11		(2.7)	36	(1.5)	11	(3.0)	11	(2.5)	29	(1.2)	32	(1.6)	26	(1.4)	31	(1.6)	30	(1.5)
12 - 19	25	(3.8)	34	(2.8)	6	(0.9)	9*	(2.8)	28	(2.6)	29	(2.0)	19	(1.7)	25	(2.1)	28	(2.3)
20 and over	40	(1.2)	35	(1.4)	23	(3.2)	18	(3.4)	28	(0.7)	33	(1.2)	24	(1.1)	25	(1.7)	29	(1.0)
2 and over	38	(1.2)	35	(1.1)	21	(2.8)	16	(2.6)	29	(0.6)	32	(0.9)	23	(0.9)	26	(1.3)	29	(0.9)

**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic (continued)

Race/ethnicity and age	Ch	olina	Vitam	nin B12	Vito	min C	Vito	nin D	(alp	min E oha-	Vita	nin K	Cal	cium	Phos	phorus	Magr	nesium
(years)	Choline Vitamin B12 Vitamin C Vitamin D % (SE) % (SE) % (SE) % (SE)		tocopherol) Vita % (SE) %			Vitamin K % (SE)		Calcium % (SE)		% (SE)		(SE)						
Non-Hispanic White:																		
2 - 5	28	(1.7)	35	(2.2)	17	(2.0)	40	(2.4)	18	(2.3)	14	(2.6)	26	(1.5)	24	(1.3)	22	(1.7)
6 - 11	25	(2.0)	35	(2.8)	20	(2.6)	41	(2.7)	20	(4.6)	9	(1.5)	27	(1.5)	22	(1.0)	19	(0.8)
12 - 19	20	(1.3)	26	(3.1)	24	(3.2)	40	(3.7)	15	(2.5)	11	(1.8)	23	(1.6)	19	(1.1)	18	(1.1)
20 and over	26	(0.7)	23	(1.1)	21	(0.8)	37	(1.3)	20	(0.9)	10	(1.0)	22	(0.8)	20	(0.5)	20	(0.5)
2 and over	25	(0.7)	25	(1.0)	21	(0.8)	38	(1.2)	19	(0.7)	10	(0.9)	23	(0.6)	20	(0.4)	20	(0.4)
Non-Hispanic Black:																		
2 - 5	25	(1.7)	37	(2.5)	20	(2.0)	48	(1.6)	15	(1.3)	10	(2.2)	31	(2.0)	24	(1.8)	22	(1.5)
6 - 11	23	(1.8)	34	(2.6)	21	(1.9)	46	(2.6)	14	(1.4)	9	(1.8)	27	(1.8)	22	(1.4)	20	(1.3)
12 - 19	22	(2.5)	24	(2.1)	18	(2.4)	41	(2.7)	14	(2.2)	8	(1.4)	22	(1.7)	18	(1.3)	15	(1.0)
20 and over	25	(1.2)	22	(1.3)	20	(1.2)	37	(1.7)	17	(0.8)	10	(0.9)	23	(1.0)	19	(0.7)	18	(0.6)
2 and over	24	(1.2)	24	(1.1)	20	(1.0)	39	(1.5)	16	(0.8)	9	(0.7)	23	(0.9)	19	(0.7)	18	(0.5)
Non-Hispanic Asian:																		
2 - 5	25*	(2.3)	41*	(3.1)	18*	(4.6)	41*	(5.0)	20*	(4.8)	10*	(3.7)	30*	(3.8)	25*	(3.4)	22*	(3.3)
6 - 11	28	(2.5)	37	(4.2)	11*	(3.0)	42	(4.9)	16*	(2.7)	10*	(1.6)	27	(2.0)	24	(1.1)	19*	(0.9)
12 - 19	26	(2.4)	31	(6.4)	19	(5.0)	41	(8.3)	18	(1.9)	7*	(1.5)	27	(2.8)	22	(1.7)	19	(1.2)
20 and over	23	(0.9)	26	(3.6)	15	(1.6)	31	(2.7)	17	(1.0)	11	(1.3)	26	(1.7)	21	(0.8)	21	(0.7)
2 and over	24	(0.9)	27	(3.0)	15	(1.6)	32	(2.1)	17	(0.8)	10	(1.1)	26	(1.3)	21	(0.7)	21	(0.6)
Hispanic:																		
2 - 5	35	(2.8)	36	(2.2)	20	(2.3)	41	(3.1)	24	(2.1)	21	(3.5)	31	(2.1)	29	(1.5)	26	(1.6)
6 - 11	30	(1.7)	34	(1.5)	25	(2.3)	44	(1.7)	20	(1.5)	15	(2.1)	30	(1.4)	27	(1.2)	25	(1.4)
12 - 19	25	(2.6)	28	(3.1)	25	(1.7)	40	(4.0)	17	(1.6)	11	(1.6)	25	(1.9)	21	(1.7)	20	(1.2)
20 and over	34	(0.9)	26	(1.6)	26	(1.4)	41	(2.0)	25	(1.0)	23	(1.8)	29	(0.8)	28	(0.9)	27	(0.8)
2 and over	33	(0.9)	28	(1.2)	25	(1.2)	42	(1.4)	23	(0.9)	21	(1.6)	29	(0.7)	27	(0.8)	26	(0.7)

**Table 14. Breakfast¹:** Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic (continued)

Race/ethnicity and age	Ir	on	Zi	inc	Coj	pper	Sele	nium	Potas	ssium	Soc	lium	Caf	feine	Alc	cohol
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Non-Hispanic White:																
2 - 5	33	(1.9)	26	(1.4)	19	(2.1)	18	(1.0)	21	(1.3)	16	(0.7)	15	(2.0)		
6 - 11	34	(2.3)	25	(1.4)	15	(0.7)	17	(1.3)	19	(0.8)	14	(0.8)	18	(2.6)		
12 - 19	26	(1.6)	16	(1.6)	15	(1.7)	13	(0.6)	19	(1.1)	13	(0.5)	27	(5.6)		
20 and over	26	(0.9)	18	(0.8)	16	(0.4)	18	(0.5)	20	(0.5)	13	(0.4)	43	(1.8)	1*	(0.5)
2 and over	27	(0.8)	19	(0.8)	16	(0.4)	17	(0.5)	20	(0.5)	13	(0.3)	42	(1.8)		
Non-Hispanic Black:																
2 - 5	35	(1.5)	26	(2.0)	18	(1.1)	17	(1.5)	21	(1.4)	15	(1.3)	7*	(2.3)		
6 - 11	31	(1.9)	23	(1.8)	16	(0.9)	15	(2.0)	19	(1.1)	15	(1.1)	10	(2.0)		
12 - 19	22	(2.2)	18	(1.6)	12	(0.8)	13	(1.3)	15	(1.2)	13	(1.5)	14	(2.4)		
20 and over	24	(0.8)	19	(0.6)	16	(0.6)	17	(0.8)	18	(0.6)	15	(0.6)	47	(1.7)	1*	(0.6)
2 and over	25	(0.7)	19	(0.6)	15	(0.5)	17	(0.8)	18	(0.5)	15	(0.6)	45	(1.7)		
Non-Hispanic Asian:																
2 - 5	39*	(4.9)	34*	(3.5)	17*	(2.9)	17*	(2.6)	21*	(2.4)	16*	(2.8)	33*	(11.0)		
6 - 11	29	(2.4)	24	(1.8)	15*	(1.1)	19*	(1.6)	19*	(1.1)	18*	(2.2)	25*	(5.7)		
12 - 19	27	(3.9)	20	(1.7)	15	(0.9)	18	(1.9)	19	(1.5)	17	(2.0)	33	(8.9)		
20 and over	25	(2.6)	19	(0.7)	18	(0.6)	19	(0.9)	20	(0.6)	16	(1.1)	43	(2.5)	#	
2 and over	26	(2.0)	20	(0.8)	18	(0.5)	19	(0.8)	20	(0.6)	16	(0.9)	43	(2.2)		
Hispanic:																
2 - 5	35	(2.7)	31	(1.9)	23	(1.8)	27	(2.4)	26	(1.3)	23	(1.7)	24*	(9.0)		
6 - 11	30	(1.5)	27	(1.6)	21	(1.4)	24	(1.6)	25	(1.3)	21	(1.2)	22*	(8.4)		
12 - 19	27	(1.7)	21	(2.0)	18	(1.4)	16	(1.8)	20	(1.4)	15	(1.4)	33	(4.4)		
20 and over	29	(0.9)	24	(1.0)	23	(1.0)	27	(0.9)	27	(0.8)	23	(0.9)	52	(1.6)	1*	(0.6)
2 and over	29	(0.8)	25	(0.9)	22	(0.9)	25	(0.9)	26	(0.8)	22	(0.9)	50	(1.5)		

# **Symbol Legend**

\* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages and ratios are as follows:

**Percent reporting:** An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 2.76.

**Nutrient ratios expressed as percentages:** An estimated ratio between 25 and 75 percent is flagged when based on a sample size  $n^*$  of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect and  $n^*$  is the number of individuals in the sample reporting non-zero intake of the respective nutrient. An estimated ratio less than or equal to 25 percent or greater than or equal to 75 percent, is flagged when the smaller of  $n^*p$  and  $n^*$  (1-p) is less than 8 times the VIF, where p is the percentage expressed as a fraction. Additionally, an estimated ratio is flagged when either the relative standard error or p/(1-p) times the relative standard error is greater than 30 percent. The VIF used in this table is 2.76.

# Indicates a non-zero value too small to report.

### **Footnotes**

- <sup>1</sup> Breakfast includes eating occasions designated by the respondent as "breakfast", or the Spanish equivalents "desayuno", and "almuerzo." Please note these eating occasions include consumption of beverages including water.
- <sup>2</sup> Percentages are estimated as a ratio of total nutrients from breakfast for all individuals to total daily nutrient intakes for all individuals. Total daily nutrient intakes are available from: <a href="https://www.ars.usda.gov/nea/bhnrc/fsrg">www.ars.usda.gov/nea/bhnrc/fsrg</a>. See Table 2. Nutrient Intakes from Food and Beverages: Mean Amounts Consumed per Individual, by Race/Ethnicity and Age, in the United States, 2017-March 2020 Prepandemic.
- <sup>3</sup> The percentage of respondents in the race/ethnicity/age group who reported consuming at least one item at an eating occasion designated as breakfast.

### **Abbreviations**

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

## Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2017-March 2020 Prepandemic

The 2017–March 2020 files represent a unique public-use data release from NHANES. The coronavirus disease 2019 (COVID-19) pandemic required suspension of data collection in March 2020. As a result, the partially completed NHANES 2019–2020 cycle was not nationally representative. Therefore, the 2019–March 2020 data were combined with the data from the 2017–2018 cycle to create the nationally representative 2017–March 2020 prepandemic data files.

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2017-March 2020 Prepandemic. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2017-2018 and 2019-2020 as applicable <a href="https://www.ars.usda.gov/nea/bhnrc/fsrg">www.ars.usda.gov/nea/bhnrc/fsrg</a>.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

### **Suggested Citation**

U.S. Department of Agriculture, Agricultural Research Service. 2022. Breakfast: Percentages of Selected Nutrients Contributed by Food and Beverages Consumed at Breakfast, by Race/Ethnicity and Age, What We Eat in America. NHANES 2017-March 2020 Prepandemic.