



United States Department of Agriculture  
Research, Education and Economics



Agricultural Research Service  
Date: 1/14/2010

# Natural Resources Research Update

ARS would like your feedback as to the value of this technical update. You can provide feedback about this specific update by responding to this email or send your comments to [NRRU.update@ars.usda.gov](mailto:NRRU.update@ars.usda.gov) and refer to technical update number: 249763

Contributing Scientists: Richard L. Haney

Location: USDA-ARS Grassland, Soil & Water Research Laboratory, Temple, Texas

Title: Use of the Solvita<sup>®</sup> gel system to measure soil carbon dioxide respiration as a measure of soil fertility.

The measurement of soil carbon dioxide respiration is a means to gauge biological soil fertility. Test methods for respiration employed in the laboratory vary somewhat, and to date the equipment and labor required have somewhat limited more widespread adoption of such methodologies. A new method to measure soil respiration was tested along with the traditional alkali trap and titration method (1). The new method involves the Solvita<sup>®</sup> gel system which was originally designed for CO<sub>2</sub> respiration from compost, but has been applied in this research to soils with treatments of increasing dairy manure compost. The objective of this research was to: (1) examine the relationship between the CO<sub>2</sub> release after one day of incubation from soils amended with dairy manure compost that have been dried and rewetted as determined using the titration method and the Solvita<sup>®</sup> gel system; and (2) compare water-soluble organic N, as well as C, N, and P mineralization after 28 days of incubation with 1-d CO<sub>2</sub> release from the titration method and Solvita<sup>®</sup> gel system. Both 1-day CO<sub>2</sub> from titration and the Solvita<sup>®</sup> gel system were highly correlated with cumulative 28 day CO<sub>2</sub> as well as the basal rate from 7-28 days of incubation. Both methods were also highly correlated with 28 day N and P mineralization as well as the initial water extractable organic N and C concentration. The data suggest that the Solvita<sup>®</sup> gel system for soil CO<sub>2</sub> analysis could be a simple and easily used method to quantify soil microbial activity and possibly provide an estimate of potential mineralizable N and P.

In another study (2) soil CO<sub>2</sub> respiration was measured using three methods: (1) titration method; (2) Infrared gas analysis (IRGA); and (3) the Solvita<sup>®</sup> gel system for soil CO<sub>2</sub> analysis. We acquired 36 soil samples from across the USA for comparison which ranged in pH from 4.5-8.5, organic C from 0.8-4.6% and the clay content from 6-62%. All three methods were highly correlated with each other after 24 hr. of incubation (titration and Solvita<sup>®</sup>  $r^2=0.82$ , respirometer and Solvita<sup>®</sup>  $r^2=0.79$  and titration vs. respirometer  $r^2=0.95$ ).

The 24 hr. (1-day) CO<sub>2</sub> release from all three methods was also highly correlated to both basal soil respiration (7-28 d) and cumulative 28 d CO<sub>2</sub> respiration. An additional 24 soil samples were acquired and added to the original 36, for a total of 60 soil samples. These samples were used for calibration of the Solvita<sup>®</sup> gel digital reader results using CO<sub>2</sub>-titration results and regression analysis. Regression analysis resulted in the equation  $y=20.6*(\text{Solvita}^{\text{®}} \text{ number})-16.5$  with an  $r^2$  of 0.83. The data suggest that the Solvita<sup>®</sup> gel system for soil CO<sub>2</sub> analysis could be a simple and easily used method to quantify soil microbial activity. Once standardized soil sampling and laboratory analysis protocols are established, the Solvita<sup>®</sup> method could be easily adapted to commercial soil testing labs as an index of soil microbial activity.

1. Haney, R.L., Brinton, W.H., Evans, E. 2008. Estimating soil carbon, nitrogen, and phosphorus mineralization from short-term CO<sub>2</sub> respiration. *Communications in Soil Science and Plant Analysis*. 39(17&18):2706-2720.
2. Haney, R.L., Brinton, W.F., Evans, E. 2008. Soil CO<sub>2</sub> respiration: Comparison of chemical titration, CO<sub>2</sub> IRGA analysis and the Solvita<sup>®</sup> gel system. *Renewable Agriculture and Food Systems*. 23: 171-176.

As the principal in-house research arm of the U.S. Department of Agriculture, the Agricultural Research Service has a mission to: Conduct research to develop and transfer solutions to agricultural problems of high national priority and provide information access and dissemination to ensure high-quality, safe food, and other agricultural products, assess the nutritional needs of Americans, sustain a competitive agricultural economy, enhance the natural resource base and the environment, and provide economic opportunities for rural citizens, communities, and society as a whole.

The U. S. Department Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance. (Not all prohibited bases apply to all programs) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.