

US007460227B1

(12) United States Patent Kim et al.

(54) METHOD TO DETECT BONE FRAGMENTS DURING THE PROCESSING OF MEAT OR FISH

(75) Inventors: Moon S. Kim, Silver Spring, MD (US);

Alan M. Lefcourt, Elkridge, MD (US); Yud-Ren Chen, Laurel, MD (US)

(73) Assignee: The United States of America as

represented by the Secretary of Agriculture, Washington, DC (US)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 331 days.

(21) Appl. No.: 11/304,283

(22) Filed: Dec. 15, 2005

Related U.S. Application Data

- (60) Provisional application No. 60/640,339, filed on Dec. 30, 2004.
- (51) **Int. Cl. G01J 3/30** (2006.01)

(56) References Cited

U.S. PATENT DOCUMENTS

4,631,413 A * 12/1986 Jensen et al. 250/458.1

(10) **Patent No.:**

US 7,460,227 B1 Dec. 2, 2008

(45) **Date of Patent:**

5,428,657 A *	6/1995	Papanicolopoulos et al 378/86
5,847,382 A *	12/1998	Koch et al 250/223 R
5,902,177 A *	5/1999	Tessier et al 452/156
6,808,448 B1*	10/2004	Kanaya et al 452/2

OTHER PUBLICATIONS

Kim et al., "Multispectral laser-induced fluorescence imaging system for large biological animals", Jul. 1, 2003, Applied Optics, vol. 42, No. 19, pp. 3927-3934.*

Kim, Moon S., et al., "Optimal Fluorescence Excitation and Emission Bands for Detection of Fecal Contamination", Journal of Food Protection, vol. 66, No. 7, 2003, pp. 1198-1207.

* cited by examiner

Primary Examiner—Kara E Geisel (74) Attorney, Agent, or Firm—Randall E. Deck; John Fado; Lesley Shaw

(57) ABSTRACT

Bones, bone fragments, and shell fragments on or near the surface of processed animal flesh may be detected by detecting their fluorescent emissions. The surface of the flesh is illuminated with UV or visible light to elicit fluorescence of animal bones or shells, and the fluorescent light emissions are measured at first and second different wavelengths, which wavelengths are selected to yield a substantial difference between the ratio or subtractive difference for bones or shells and the ratio or subtractive difference for animal flesh. The ratio and/or subtractive difference of the fluorescent light emissions at the first and second wavelengths is calculated, and the presence of bone, bone fragment, or shell fragment is determined therefrom.

22 Claims, 9 Drawing Sheets

