

# Dietary modulation of gut microflora in broiler chickens: a review of the role of six kinds of alternatives to in-feed antibiotics

Y. YANG, P.A. IJI and M. CHOCT\*

School of Environmental and Rural Science, University of New England, Armidale NSW 2351, Australia

\*Corresponding author: [mchoct@poultrycrc.com.au](mailto:mchoct@poultrycrc.com.au)

---

Since in-feed antibiotics (IFAs) are being taken out of broiler diets around the world, beginning in Sweden in the year 1986, the search for alternatives to replace IFAs has gained increasing interest in animal nutrition. Gut microflora appears to be the target for IFAs and alternatives to exert health benefits and some growth-promoting effects. In this review the effect of six kinds of alternatives to IFAs on gut microflora is discussed and their working mechanisms and growth-promoting effects are reviewed. This review focuses on mannanoligosaccharides (MOS) as alternative to antibiotics using research results obtained by various authors in recent years.

---

**Keywords:** alternatives to in-feed antibiotics; fibre-degrading enzymes; prebiotics; probiotics; mannanoligosaccharides; symbiotics; phytobiotics; gut microflora

## Introduction

Antibiotics have been added to poultry and pig diets to maintain health and production efficiency in the last few decades (Rosen, 1995). However, because of the development of resistance by pathogenic bacteria, which can impact on public health, antibiotics are being taken out of poultry and pig diets around the world, beginning in Sweden in the year 1986 (Dibner and Richards, 2005). The search for alternatives to replace IFAs has gained increasing interest in animal nutrition in recent years. Bedford (2000) pointed out that the growth-promoting effects of antibiotics in animal diets are clearly related to the gut microflora because they exert no benefits on the performance of germ-free (GF) animals.

Gut microflora has significant effects on host nutrition, health, and growth performance (Barrow, 1992) by interacting with nutrient utilization and the development of gut system of the host. This interaction is very complex and, depending on the composition and

activity of the gut microflora, it can have either positive or negative effects on the health and growth of birds. For example, when pathogens attach to the mucosa, gut integrity and function are severely affected (Droleskey *et al.*, 1994) and immune system threatened (Neish, 2002). Chicks grown in a pathogen-free environment grow 15% faster than those grown under conventional conditions where they are exposed to bacteria and viruses (Klasing, 1987). Furthermore, it is generally agreed that gut microflora is a nutritional “burden” in fast-growing broiler chickens (Dibner and Richards, 2005; Lan *et al.*, 2005) since an active microflora component may have an increased energy requirement for maintenance and a reduced efficiency of nutrient utilization.

The focus of alternative strategies has been to prevent proliferation of pathogenic bacteria and modulation of indigenous bacteria so that the health, immune status and performance are improved (Ravindran, 2006). In this review, we will evaluate dietary modulation of gut microflora through the use of fibre-degrading enzymes, probiotics, prebiotics, MOS, phytobiotics, as well as their mechanisms of action and effectiveness in promoting growth in broiler chickens.

## **Fibre-degrading enzymes**

Enzymes are naturally occurring and are produced by all living organisms for catalyzing chemical reactions. Enzymes were discovered in the later part of the 19<sup>th</sup> century and have been used in industry and food processes since the early 1900s (Clarkson *et al.*, 2001). The majority of enzyme products are fermentation products of basophilic microorganisms (Clarkson *et al.*, 2001). Ferket (1993) defined enzymes as special proteins that catalyse or accelerate the rate of specific chemical reactions in which the enzyme activity may be dependent on the substrate in a random manner or it may be through very specific sites on substrates such as fat, protein, or carbohydrates. In monogastric animal diets, exogenous enzymes are used to improve digestibility of a wide range of feed components such as fibre, phytate, protein, etc. Fibre-degrading enzymes are used to break down specially non-starch polysaccharides (NSP), which are large polymers, to smaller polymers to alleviate their anti-nutritive activities (Choct and Annisson, 1992).

The effects of enzymes on gut microflora were classified by Bedford (2000) into two phases: an ileal phase and a caecal phase. In the ileum, enzymes simply reduce the number of bacteria by increasing the rate of digestion and limiting the amounts of substrates available to the microflora. In the caecal phase, enzymes produce soluble, poorly absorbed sugars which feed beneficial bacteria. The volatile fatty acids (VFAs) produced by such bacteria may be of benefit not only in controlling populations of *Salmonella*, and perhaps, *Campylobacter* species, but also in providing an energy source for the bird (Snel *et al.*, 2002).

However, the effects of enzymes on the gut microflora may be far more than those two phases. The composition of gut microflora in the proximal small intestine as well as those associated with the gut wall was shown to be changed by the addition of xylanase (Vahjen *et al.*, 1998; Danicke *et al.*, 1999; Hubener *et al.*, 2002). The authors correlated those effects of xylanase on the gut microflora with its effects on the viscosity of diet, which is well known as one of the major modes of action of enzymes. Inclusion of cereals rich in NSP increases the viscosity of the digesta, reduces apparent nutrient digestibility, alters bacterial profiles and gut physiology. By adding enzymes into a diet, the viscosity of the content is reduced and nutrient uptake and animal performance are improved (Bedford, 2001). Elimination of cell wall encapsulation is another major mode of action of enzymes (Bedford, 2001). It relies on the fact that the feed manufacturing process of grinding and pelleting does not break

open all the cell walls of the endosperm. The addition of enzymes can remove such “encapsulated” material in the gut and hence improve nutrient utilization by the birds.

Therefore, the responses of birds to enzymes depend mainly on dietary cereal quality and quantity. However, quantity and quality of fat, microbial status, bird age and antimicrobial agents can also modify the effects of enzymes (Bedford, 2001). Furthermore, because of cereal quality (e.g. the complexity of carbohydrate) and the thermolabile characteristic of enzymes, improvements are not always observed (Acamovic, 2001).

Supplementation of enzymes generally can lead to 2-5% improvement in feed/gain ratio and 2-3% improvement in growth rate (Broz and Beardsworth, 2002). Reduced incidence of sticky excreta and improved litter conditions are also the benefits of using enzymes, which makes the problem of NSP and associated increased ingesta viscosity more manageable (Morrow, 2001). The development of enzymes is towards a specially-designed stage to use NSP as energy sources and to deactivate anti-nutrients in feed (Choct, 2006).

## **Prebiotics**

Gibson and Roberfroid (1995) defined a prebiotic as a non-digestible food ingredient which beneficially affects the host by selectively stimulating the growth of and/or activating the metabolism of one or a limited number of health-promoting bacteria in the intestinal tract, thus improving the host's microbial balance. The growth of endogenous microbial population groups such as bifidobacteria and lactobacilli is specifically stimulated and these bacteria species are perceived as beneficial to animal health. Prebiotics have the advantage, compared with probiotics, that bacteria are stimulated which are normally present in the GIT of that individual animal and therefore already adapted to that environment (Snel *et al.*, 2002). The dominant prebiotics are fructo-oligosaccharide products (FOS, oligofructose, inulin) (Patterson and Burkholder, 2003); gluco-oligosaccharides, stachyose, malto-oligosaccharides, and oligochitosan have also been investigated in broiler chickens (Zhang *et al.*, 2003; Gao and Shan, 2004; Jiang *et al.*, 2006; Huang *et al.*, 2007).

Compared to the application of prebiotics in human and pet food, the use of prebiotics in broiler chicken diets does not have a long history. Reports on the effects of prebiotics on the activity of the microflora of broilers are limited and the effects are variable, depending on the type of prebiotic. Fructo-oligosaccharides were shown to support the growth of beneficial bacteria, such as lactobacilli (Xu *et al.*, 2003; Yusrizal and Chen, 2003; Zhan *et al.*, 2003), but failed to stimulate the growth of bifidobacteria (Vidanarachchi *et al.*, 2006). When supplemented into a dextrose-isolated soy protein diet, short-chain fructo-oligosaccharides decreased caecal populations of *C. perfringens* (Biggs *et al.*, 2007). Reduced susceptibility to *Salmonella* colonization was noticed in birds on fructo-oligosaccharide treatments compared to controls (Bailey *et al.*, 1991; Fukata *et al.*, 1999). The addition of isomalto-oligosaccharides and stachyose did not affect crop and/or caecal bacterial populations such as lactobacilli, *E. coli* and total aerobes (Zhang *et al.*, 2003; Jiang *et al.*, 2006). It was shown that prebiotics can bring about “bifidogenic” effects and a shift in microbial metabolism from “proteolytic” to the more favourable “saccharolytic” in mice (Gibson and Roberfroid, 1995). A combination of various substances with different rates of fermentation will be effective in mimicking some of the antibiotic effects in pigs (Williams *et al.*, 2001). However, no similar research has been reported in broiler chickens.

Among the prebiotics examined in chicken, fructo-oligosaccharides were shown to

improve growth performance of birds but positive effects are not always observed (Table 1).

**Table 1** Effects of prebiotics on growth performance of broilers .

Parameter	Prebiotics	Response (%)	Reference	
Weight gain (g/bird)	FOS	+5	Li <i>et al.</i> , 2008	
		-2	Biggs <i>et al.</i> , 2007	
		+6	Zhan <i>et al.</i> , 2003	
		+8	Xu <i>et al.</i> , 2003	
	transgalacto-oligosaccharide	0	Biggs <i>et al.</i> , 2007	
		stachoyse	-3	Jiang <i>et al.</i> , 2006
		chitosan	+2	Huang <i>et al.</i> , 2005
isomalto-oligosaccharide	+5	Zhang <i>et al.</i> , 2003		
Body weight (g/bird)	FOS	+3	Yusrizal and Chen, 2003	
		-1	Waldroup <i>et al.</i> , 1993	
FCR (g/g)	FOS	+2	Li <i>et al.</i> , 2008	
		+3	Biggs <i>et al.</i> , 2007	
		0	Yusrizal and Chen, 2003	
		+6	Zhan <i>et al.</i> , 2003	
		+6	Xu <i>et al.</i> , 2003	
	transgalacto-oligosaccharide	-1	Waldroup <i>et al.</i> , 1993	
		0	Biggs <i>et al.</i> , 2007	
		stachoyse	-1	Jiang <i>et al.</i> , 2006
		chitosan	+3	Huang <i>et al.</i> , 2005
		isomalto-oligosaccharide	+4	Zhang <i>et al.</i> , 2003

FCR: Feed conversion ratio.

Also, dietary supplementation of fructo-oligosaccharide (0.3% dose) or oligochitosan (0.1% dose) showed growth-promoting effects similar to antibiotic treatments based on flavomycin (Huang *et al.*, 2005) or aureomycin (Li *et al.*, 2008). The optimal dose for prebiotics to exert growth-promoting effects is not easy to define; however feeding a higher level (0.8%) of inulin and short-chain fructo-oligosaccharide depressed growth performance, digestibility of amino acids as well as metabolisable energy of birds (Biggs *et al.*, 2007). Some positive changes in digestive enzymes, gut morphology, and immune system were noticed in birds given prebiotic-supplemented feed (Xu *et al.*, 2003; Zhang *et al.*, 2003; Huang *et al.*, 2007).

However, there are many considerations in supplementing prebiotics in animal feed. These include the type of diet (*i.e.*, the content of non-digestible oligosaccharides); the type and inclusion level of the supplements; the animal characteristics (species, age, stage of production); and the hygiene status of the farm (Verdonk *et al.*, 2005). The primary ones are the type and inclusion level of the supplement as high dosage of prebiotics can have negative effects on the gut system and retard the growth rate of birds as observed by Biggs *et al.* (2007). It is reported that rapid fermentation of prebiotics, leading to high concentrations of organic acids, impaired the barrier function, which reduced the ability of rats to resist salmonella infection (Ten Bruggencate *et al.*, 2003). It may also be worthwhile to examine the interaction between prebiotics(s) and bird sex. In the report by Yusrizal and Chen (2003), body weight and feed conversion ratio (FCR) of female birds were improved by 10% and 9%, respectively, on oligofructose treatment but no such effects were observed in males.

## Probiotics

In animal nutrition, probiotics are defined as viable micro-organisms used as feed additives, which lead to beneficial effects for the host by improving its microbial balance (Fuller, 1989) or the properties of the indigenous microflora (Havenaar and Huis In 't Veld, 1992). A variety of microbial species have been used as probiotics, including species of *Bacillus*, *Bifidobacterium*, *Enterococcus*, *Escherichia*, *Lactobacillus*, *Lactococcus*, *Streptococcus*, a variety of yeast species, and undefined mixed cultures. *Lactobacillus* and *Bifidobacterium* species have been used most extensively in humans, whereas species of *Bacillus*, *Enterococcus*, and *Saccharomyces* yeast have been the most common organisms used in livestock (Simon et al., 2001). There has been an increase in research on feeding *Lactobacillus* to broiler chickens (Jin et al., 2000; Kalavathy et al., 2003).

The possible modes of action of probiotics were extensively reviewed (Jin et al., 1997; Simon et al., 2001; Ghadban, 2002; Edens, 2003). Two basic mechanisms by which probiotics act to maintain a beneficial microbial population include “competitive exclusion” and immune modulation. Competitive exclusion involves competition for substrates, production of antimicrobial metabolites that inhibit pathogens, and competition for attachment sites. Based on this mechanism, probiotics have been tested for their efficacy at controlling *Salmonella* colonization in broilers and the results are positive and consistent (Table 2).

**Table 2 Effectiveness of probiotics in the prevention of *Salmonella* colonization in broilers in research station and/or commercial field tests.**

Researcher	Number of Chickens	Reduction (%) in the colonization
Higgins et al. (2007)	840	60
Stern et al. (2001)	210	44
Pascual et al. (1999)	344	50
Stern et al. (1998)	100,000	9
Blankenship et al. (1993)	Six commercial blocks	31
Bolder et al. (1992)	420	20
Hinton et al. (1991)	720	39
Goren et al. (1988)	8 million	31

Further, a product of competitive exclusion specially designed at gut mucosal level showed much stronger effectiveness than a general competitive exclusion product; the number of affected birds was reduced by 50% and 10%, respectively (Stern et al., 2001). Probiotic supplementation, especially with *lactobacillus* species, has also shown beneficial effects on resistance to the other infectious agents such as *Clostridium* population (Decroos et al., 2004) and *Campylobacter* (Stern et al., 2001). Regarding the gut microbiota of normal birds, the results of probiotics supplementation are variable because of the difference in origin, strain as well as species of probiotics. Reduced caecal coliform populations were noticed in chickens given a diet supplemented with lactobacilli strains, isolated from chicken intestine, but the populations of other kinds of bacteria were not affected (Watkins and Kratzer, 1984; Jin et al., 1998a, 1998b). In contrast, Murry et al. (2006) reported that birds supplemented with botanical probiotic containing lactobacilli had higher lactobacilli but lower *C. perfringens* compared to the control birds. When multi-strain and/or multi-species probiotics were applied, no significant change(s) in bacterial populations was noticed (Priyankarage et al., 2003; Mountzouris et al., 2007).

By directly interacting with gut mucosal immune system, probiotics can modulate either innate or acquired immunity, or both (Dugas *et al.*, 1999). Further, specific immune modulatory effects of probiotics are dependent on the strain or species of bacteria included in the probiotics (Edens, 2003; Huang *et al.*, 2004). Lactobacilli, the most studied strain of probiotic in both animal and human, have been implicated to increase the activity of certain innate immune functions, specifically the activity of macrophages and natural killer cells (McCracken and Gaskins, 1999). In accordance with this notion, Koenen *et al.* (2004) reported that feeding *L. paracasei* to broilers enhanced the phagocytic activity of the gut cells (caecum, ileum). However, *L. plantarum*, rather than *L. paracasei*, exerted stronger stimulating effect on antigen-specific titre (Koenen *et al.*, 2004). In general, feeding probiotics could improve antibody titres against Newcastle disease; infectious bursal disease virus and/or sheep red blood cell (SRBC, Panda *et al.*, 2000; Zulkifli *et al.*, 2000; Huang *et al.* 2004) but no responses were also observed by Panda *et al.* (1999). The results from Zulkifli *et al.* (2000) further indicated that these effects of probiotics could be affected by the age and strain of broilers. In addition, *Lactobacillus*-based probiotic may strengthen gut defence function via activation and enhancement of local cell-mediated immunity to against certain enteric pathogen (Dalloul *et al.*, 2003). However the exact mechanisms for probiotics to enhance immune function remain largely unknown.

Probiotics did not consistently improve growth performance and/or mortality rate of birds (Table 3).

**Table 3 Growth performance and/or mortality rate of birds to probiotic supplementation.**

Item	Control	Probiotics	Improvement (%)
<b>Liu <i>et al.</i> (2007)</b>			
BWG (g/bird)	1892	1920	+1
FCR (g/g)	1.75	1.74	0
<b>Mountzouris <i>et al.</i> (2007)</b>			
BWG (g/bird)	2216	2237	+1
FCR (g/g)	1.81	1.78	+2
<b>Murry <i>et al.</i> (2006)</b>			
BWG (g/bird)	2784	2720	- 2
FCR (g/g)	1.62	1.63	0
Mortality (%)	7.02	4.76	+32
<b>Timmerman <i>et al.</i> (2006)</b>			
ADG (g/bird)	49.99	49.65	0
FCR (g/g)	1.93	1.87	+3
Mortality (%)	8.84	7.27	+18
<b>Kalavathy <i>et al.</i> (2003)</b>			
BWG (g/bird)	2151	2251	+5
FCR (g/g)	1.96	1.78	+9
<b>Zulkifli <i>et al.</i> (2000)</b>			
BWG (g/bird)	1379	1545	+12
FCR (g/g)	2.08	2.17	-4
Mortality (%)	1.7	2.2	-29
<b>Jin <i>et al.</i> (1998a)</b>			
BWG (g/bird)	1290	1388	+8
FCR (g/g)	2.27	2.10	+7
Mortality (%)	6.7	5.3	+21
<b>Mohan <i>et al.</i> (1996)</b>			
BWG (g/bird)	1046	1128	+8
FCR (g/g)	2.01	1.84	+8

BWG: body weight gain. FCR: feed conversion ratio. ADG: average daily gain.

The inconsistency may become more complex because of rearing environment. For example, under heat stress condition, lactobacilli probiotic supplementation improved body weight gain (BWG) of female birds by 12% but increased FCR and mortality rate by 4% and 29%, respectively (Zulkifli *et al.*, 2000). However, growth-promoting effects of certain probiotics were reported to be comparable to antibiotic treatments (virginiamycin, Cavazzoni *et al.*, 1998; oxytetracycline, Zulkifli *et al.*, 2000; avilamycin, Mountzouris *et al.*, 2007); a meta-analysis of 35 studies involved in probiotics across Brazil between 1995 and 2005 seems to indicate that probiotics are a technically viable alternative to IFA in broiler feeding (Faria Filho *et al.*, 2006). However, the growth-promoting effects of probiotics are dependent on the specific probiotics, the application level of probiotics, the age of birds as well as the delivery method (*i.e.* via water and/or feed). Moreover, there are many factors from nutrition, environment (sanitary condition), to management that could compromise the effectiveness of probiotics (Edens, 2003). This can probably explain the inconsistent results in the growth performance and gut bacterial responses described above.

## **Mannan oligosaccharides**

Mannan oligosaccharides, derived from yeast cell wall, are more complex than the name suggests; they are components of the outer layer of yeast cell walls and their components include proteins, glucans and phosphate radicals as well as mannose (Klis *et al.*, 2002). The basic composition of the wall consists of mannan (30%), glucan (30%) and protein (12.5%). While the ratio of one component to another remains relatively constant from strain to strain, the degree of mannan phosphorylation and the interaction among the mannan, glucan and protein components vary (Lyons, 1994). Mannan oligosaccharides contain protein which has relatively high proportions of serine, threonine, aspartic and glutamic acids, and a paucity of methionine (Song and Li, 2001).

Hooge (2004a) reviewed pen trials conducted with a commercially available dietary MOS (Bio-MOS, Alltech Inc.) from 1993 to 2003 and the meta-analysis showed that Bio-MOS improved the growth performance of birds compared to the negative control (Table 4).

**Table 4 Effects of MOS on growth performance of broiler chickens (adapted from Hooge, 2004a).**

<b>Parameter</b>	<b>Negative Control</b>	<b>MOS</b>	<b>Relative change (%)</b>	<b>Note</b>
Body weight (kg/bird)	2.231	2.267	+1.61	29 pen trials
FCR (g/g)	1.808	1.772	-1.99	29 pen trials
Mortality (%)	4.494	3.534	-21.4	21 pen trials
<b>Parameter</b>	<b>Antibiotic Control</b>	<b>MOS</b>	<b>Relative change (%)</b>	<b>Note</b>
Body weight (kg/bird)	2.246	2.238	-0.36	21 pen trials
FCR (g/g)	1.822	1.820	-0.11	21 pen trials
Mortality (%)	5.404	4.426	-18.1	16 pen trials

FCR: feed conversion ratio.

Compared to a wide range of antibiotics (including avilamycin, bacitracin, bambarmycin or virginiamycin at prophylactic concentrations), a significant decrease in mortality was observed for Bio-MOS treatment (Table 4). The optimal dose of

Bio-MOS for broiler production appears to be around 2 g/kg, depending on the production stage of birds (Rosen, 2007; Yang *et al.*, 2007b). Three major modes of action by which broiler performance is improved by MOS are proposed: 1) control of pathogenic or potential pathogenic bacteria which possess type-1 fimbriae (mannose-sensitive lectin), 2) immune modulation, and 3) modulation of intestinal morphology and expression of mucin and brush border enzymes (Ferket, 2004).

Mannan oligosaccharides not only prevent those pathogenic bacteria possessing type-1 fimbriae, such as *E. coli*, from attaching to gut wall but also displace them from the gut wall. This reduces sub-clinical or lethal infection. Pathogens possessing type-1 fimbriae are much more virulent than non-fimbriated bacteria. Duguid *et al.* (1976) demonstrated that fimbriation in *S. typhimurium* significantly increases both the number of infections (a 26% increase) and deaths (a 40% increase) in mice inoculated orally compared to non-fimbriated organisms from the same parent strain.

The immune modulatory effects of MOS are based on the following two aspects: 1) its mannan and glucan components have antigenicity characteristics, and 2) MOS prevents colonization of specific pathogens but allow them to be presented to immune cells as attenuated antigens (Ferket, 2004). The immuno-stimulatory effects of MOS were demonstrated by Privulescu (1999) in GF vs. conventional piglets (Table 5). A unique character of MOS in immune modulation is that it enhances the protective antibody response to enhance disease resistance while at the same time suppress the acute phase (fever) response (Ferket *et al.*, 2002). Mannan oligosaccharides have also been shown to enhance macrophage response in different animal species (Spring and Privulescu, 1998).

**Table 5 Effects of MOS (Bio-MOS) on immune responses of conventional and germ-free piglets. (Adapted from Privulescu, 1999)**

Item	Without MOS	With MOS	Change
<b>Cellular immune responses</b>			
Phagocytotic index	41.4	78.2	89% increase
Lymphoblast transformation	1.60	3.09	93% increase
Proportion T lymphocytes	37.6	24.7	34% decrease
<b>Cytokines dose in lymphocyte cultures (relative doses)</b>			
IL-2	1	2	100% increase
IFN- $\gamma$	1	3	200% increase
<b>Humoral immune responses</b>			
Serum IgG (mg/100mL)	200	916	358% increase
Serum IgM (mg/100mL)	Trace	106	>500% increase
Serum IgA (OD by ELISA)	163	364	123% increase
Lysozyme ( $\mu$ g/mL)	1.38	3.22	133% increase

Positive effects of MOS on growth performance and gut physiology are observed; however, the information available is limited or/and variable. A number of studies demonstrated that MOS are effective in reducing *Salmonella* infection of birds (Fernandez *et al.*, 2000; Spring *et al.*, 2000; Fernandez *et al.*, 2002; Kumar *et al.*, 2002). A few studies investigated the effect of MOS on intestinal and faecal microbial populations of broilers (Spring *et al.*, 2000; Song and Li, 2001; Jamroz *et al.*, 2004; Kocher *et al.*, 2004) but the results are inconsistent. Yang *et al.* (2008a, 2008b, 2008c) showed that MOS inhibited the development of lactobacilli and coliforms; in particular the colonization of mucosa-associated coliforms was inhibited by MOS as early as 7 days of age. In addition, MOS reduced the abundance of *C. perfringens* in the caeca of birds at 21 days of age (Yang *et al.*, 2008c), which is in agreement with the report from Jamroz *et al.* (2004) and Kocher *et al.* (2004). The effect of MOS on the

development of mucosa-associated bacteria appears to closely relate to the age of birds and rearing environment but less independent of the type of diet; whereas the effect of MOS on luminal bacteria mainly interacts with the type of diet and the age of birds.

Mannan oligosaccharides, specifically Bio-MOS, were shown to alter mucosal architecture and longer villi were noticed in birds fed MOS-supplemented diets (Iji *et al.*, 2001; Loddi *et al.*, 2002; Yang *et al.*, 2007b). Further, Bio-MOS consistently reduced the crypt depth of the mucosa of the small intestine where its growth-promoting effects were observed (Figure 1). Also, this effect was dependent on the age of birds.




Figure 1 Relative change (%) of jejunal crypt depth and growth performance of birds supplemented with Bio-MOS compared to a negative control.

BWG: Body weight gain (g/bird), FCR: Feed conversion ratio (g/g). BWG and FCR were measured from day 8 to day 21. Crypt depth ( $\mu\text{m}$ ) was measured with 7-day-old birds.

\*The experiment was conducted with day-old broilers under a challenge model of pathogenic *E. coli*. The basal diet was mainly composed of wheat, sorghum and barley (Yang *et al.*, 2008b).

\*\*The experiment was conducted with day-old broilers fed on a wheat basal diet (Yang *et al.*, 2008c).

There is scant information on the effect of MOS on nutrient digestion, availability and retention of birds. Studies by Kumprecht and Zobac (1997) showed that total tract digestibility of fibre was increased by MOS but the digestibility of fat, nitrogen-free extract and crude protein was not affected. Similarly, Yang *et al.* (2008a) did not observe any effect of MOS on total tract digestibility of starch, protein, fat and NSP in birds given a sorghum-wheat based diet but MOS significantly improved apparent metabolisable energy and numerically improved the net energy value of the diet, which agrees with the reports from Samarasinghe *et al.* (2003) that energy utilization was improved by MOS. However, the results on protein utilization are variable; while Shafey *et al.* (2001a) reported that MOS supplementation had no effect on nitrogen utilization of birds, Samarasinghe *et al.* (2003) noticed that MOS increased protein utilization. Yang *et al.* (2008c) also observed that MOS largely increased ileal starch digestibility in birds on a wheat-based diet. This effect of MOS seemed to be related to its modulatory effects on gut microflora but the exact working mechanism(s) is unclear.

The development of brush-border membrane bound enzyme activities and mucin mRNA expression was influenced by MOS (Smirnov *et al.*, 2005; Yang *et al.*, 2007a, 2008c). However, the information is too limited to draw any conclusion(s). The effects of

MOS on the immune organ or response were observed by Kumar *et al.* (2002), Kocher *et al.* (2004) but not Shafey *et al.* (2001b), an area that requires more research.

However, based on the available information, it can be concluded that the responses of birds to MOS supplementation are influenced by the type of diet, rearing conditions as well as the age of birds. Therefore, these factors need to be carefully considered in order to obtain maximal growth-promoting effects of MOS in broiler production.

## **Symbiotics**

A symbiotic is, in its simplest definition, a combination of probiotics and prebiotics (Collins and Gibson, 1999; Schrezenmeir and De Vrese, 2001). This combination could improve the survival of the probiotic organism, because its specific substrate is available for fermentation. This could result in advantages to the host through the availability of the live micro-organism and the prebiotic. Bengmark (2001) regards symbiotics as products of fermentation. Since in mixtures of pre- and probiotics, the prebiotics will be fermented when the appropriate choice of products is used, this definition may also be plausible. Examples of symbiotics are FOS and bifidobacteria, and lactitol and lactobacilli (Collins and Gibson, 1999). Bailey *et al.* (1991) used a combination of FOS and competitive exclusion flora to reduce *Salmonella* colonization in chickens. The combination was more effective in reducing *Salmonella* colonization than FOS or competitive probiotic alone. While applying the combination of FOS and bacillus to a corn-soybean basal diet, Li *et al.* (2008) observed that average daily gain (ADG) and FCR were improved by 6% and 2 %, respectively; diarrhoea and mortality rate were reduced by 58% and 67%, respectively, which were very comparable to aureomycin treatment (the relative changes are 4% for ADG, 2% for FCR, 69% for diarrhoea rate and 33% for mortality rate). To our knowledge, this is the only experiment published regarding the growth-promoting effects of symbiotic in broiler chickens thus far. Therefore more research is warranted on this kind of products in order to achieve the application significance in the industry.

## **Phytobiotics**

Plant products have been used for centuries by humans as food and to treat ailments. Natural medicinal products originating from herbs and spices have also been used as feed additives for farm animals in ancient cultures for the same length of time. To differentiate from the plant products used for veterinary purposes (prophylaxis and therapy of diagnosed health problems), phytobiotics were redefined by Windisch and Kroismayr (2006) as plant-derived products added to the feed in order to improve performance of agricultural livestock. Around the world, phytobiotics have been investigated as natural sources of biologically important chemicals since efforts are being made to ban all types of IFAs in many countries. Compared with synthetic antibiotics or inorganic chemicals, these plant-derived products have proven to be natural, less toxic, residue free, and are thought to be ideal feed additives in food animal production (Wang *et al.*, 1998).

With respect to biological origin, formulation, chemical description and purity, phytobiotics comprise a very wide range of substances and four subgroups may be classified: 1) herbs (product from flowering, non-woody and non-persistent plants), 2) botanicals (entire or processed parts of a plant, *e.g.*, root, leaves, bark), 3) essential oils (hydro distilled extracts of volatile plant compounds), and 4) oleoresins (extracts based

on non-aqueous solvents) (Windisch and Kroismayr, 2006). The active compounds of phytobiotics are secondary plant constituents.

Antimicrobial activity and immune enhancement probably are the two major mechanisms by which phytobiotics exert positive effects on the growth performance and health of animals. Compounds (phytochemicals) in phytobiotics are well known to have antimicrobial ability (Cowan, 1999). Polysaccharide components are considered to be the most important immunoactive components (Xue and Meng, 1996). In diseased chickens (either infected with avian *Mycoplasma gallisepticum* or *Eimeria tenella*), Guo and his colleagues (Guo *et al.*, 2004a, 2004b, 2004c) demonstrated that plants and their extracts could improve the growth performance, reduce the populations of coliforms and/or *C. perfringens*, and enhance both cellular and humoral immune responses of chickens. Some herbal extracts have also been shown to possess a coccidiostatic activity (Allen *et al.*, 1997; Youn and Noh, 2001; Christakia *et al.*, 2004).

A common feature of phytobiotics is that they are a very complex mixture of bioactive components. For example, hawthorn fruit, a common growth-enhancing and digestion modifier, has been shown to contain more than 70 kinds of organic chemicals along with some unidentified factors and active bio-active compounds (Wang *et al.*, 1998). Therefore they may exert multiple functions in the animal body. Increased feed intake and digestive secretions are also observed in animals offered phytobiotic-supplemented feed (Windisch and Kroismayr, 2006). Growth enhancement through the use of phytobiotics is probably the result of the synergistic effects among complex active molecules existing in phytobiotics (Gauthier, 2002). However, the exact growth-promoting mechanisms of phytobiotics in broiler chickens are poorly understood.

Among phytobiotics, essential oils (EO) have been applied into chicken feed in Europe and USA (Hooge, 2004b). However, bird growth responses to EO supplementation are still controversial. No EO effects on growth performance were reported by Botsoglou *et al.* (2002); Zhang *et al.* (2005), Jang *et al.* (2007); whereas improved growth performance were observed at different ages of birds fed certain EO-supplemented diet(s) by Jamroz *et al.* (2003), Hernandez *et al.* (2004), and Cross *et al.* (2007). On the other hand, some EO(s) induced growth improvements similar to or even better than an antibiotic treatment (Table 6).

**Table 6 Effects of essential oils on growth performance in broilers.**

Item	Negative control	Antibiotic control	Essential oil treatment
<b>Garcia <i>et al.</i> (2007)</b>			
ADG (g/bird)	68.9	66.5	68.8
FCR (g/g)	1.92	1.54	1.59
<b>Ertase <i>et al.</i> (2005)</b>			
ADG (g/bird)	61.3	65.8	71.3
FCR (g/g)	1.61	1.50	1.41
<b>Jamroz <i>et al.</i> (2003)</b>			
ADG (g/bird)	48.1	48.9	49.2
FCR (g/g)	1.85	1.81	1.79

ADG: average daily gain. FCR: feed conversion ratio.

While comparing the effects of various herbs and oils on broiler performance, Cross *et al.* (2007) concluded that the quality as well as the quantity of active chemicals in plant extract determines bird response. In addition, the efficacy of dietary EO can be affected by intrinsic and extrinsic factors such as nutritional status of animals, infection, diet

composition and environment (Giannenas *et al.*, 2003; Lee *et al.*, 2004b). Essential oils function mainly as antimicrobials and antioxidants; their antimicrobial ability may modulate the gut ecosystem to affect fat digestibility (Lee *et al.*, 2004a), starch or/and protein digestibility of feeds (Jamroz *et al.*, 2003; Hernandez *et al.*, 2004). A commercial preparation of essential oil components reduced faecal *C. perfringens* counts of broilers in a field study (Mitsch *et al.*, 2002). In addition, dietary supplementation of EO reduced the intestinal populations of *E. coli* (Jamroz *et al.*, 2003; Jang *et al.*, 2007) and increased digestive enzymes in either pancreas and/or intestinal mucosa (Lee *et al.*, 2003; Jang *et al.*, 2007); however intestinal mucosal morphology was not affected by EO supplementation (Garcia *et al.*, 2007).

Four factors may affect the effectiveness of phytobiotic additives: 1) plant parts and their physical properties, 2) source, 3) harvest time, and 4) compatibility with the other ingredient (s) in the feed (Wang *et al.*, 1998), which may also explain why 50% difference in BWG and 63% difference in FCR could happen when different kinds of phytobiotics are used in chicken diet (Xing, 2004). Although phytobiotics are a group of natural additives, research into their mechanisms of action, compatibility with diet, toxicity and safety assessment (based on the fact that some phytobiotic might have harmful substance(s)) needs to be done before they can be applied more extensively in poultry feed.

## Conclusions

The withdrawal of IFAs from poultry feed requires the industry to look for various alternatives to maintain or improve the health and performance of birds. Although the efficacy of antibiotic substitutes needs to be assessed by setting up standards and comprehensive multi-factorial models (Rosen, 2003), it is encouraging to know that most of the alternatives reviewed in the context can exert growth-promoting effects; also some of the effects are comparable to those of IFAs. However the other side of the coin is these growth-promoting effects are (very) variable; under certain circumstances the alternatives can even negatively affect the performance. Further, the effects of the alternatives on gut microflora and/or digestive physiology are often inconsistent. These facts require us to carefully examine the modes of action of the alternatives.

By increasing the growth of beneficial microbes or by reduction and removal of potential pathogens, the alternatives to IFAs possibly can improve the health and performance of birds. However, their effects on gut microflora interact with digestive physiology and thus growth in many complex ways, which can be further influenced or even determined by many other factors such as the compatibility between the diet and the alternative, hygiene standards and animal husbandry practices. There possibly remain many questions to be answered or barriers to be overcome so that the alternatives can be applied (more) successfully in the industry in future.

## References

- ACAMOVIC, T. (2001) Commercial application of enzyme technology for poultry production. *World's Poultry Science Journal* 57: 225-242.
- ALLEN, P.C., LYDON, J. and DANFORTH, H. (1997) Effects of components of *Artemisia annua* on coccidia infections in chickens. *Poultry Science* 76: 1156-1163.
- BAILEY, J.S., BLANKENSHIP, L.C. and COX, N.A. (1991) Effect of fructo-oligosaccharide on *Salmonella* colonization of the chicken intestine. *Poultry Science* 70: 2433-2438.

- BARROW, P.A.** (1992) Probiotics for chickens, in: R. FULLER (Ed.) *Probiotics: The scientific basis*, pp. 255-257 (Chapman and Hall, London).
- BEDFORD, M.** (2000) Removal of antibiotic growth promoters from poultry diets: implications and strategies to minimise subsequent problems. *World's Poultry Science Journal* **56**: 347-365.
- BEDFORD, M.R.** (2001) The role of carbohydrases in feedstuff digestion, in: MCNAB, J. & BOORMAN, K. N. (Eds) *Poultry feedstuffs: Supply, composition, and nutritive value*, pp. 319-336 (CAB international, Edinburgh, UK).
- BENGMARK, S.** (2001) Pre-, pro- and symbiotics. *Current Opinion in Clinical Nutrition and Metabolic Care* **4**: 571-579.
- BIGGS, P., PARSONS, C.M. and FAHEY, G.C.** (2007) Effects of Several Oligosaccharides on Growth Performance, Nutrient Digestibilities, and Caecal Microbial Populations in Young Chicks. *Poultry Science* **86**: 2327-2336.
- BLANKENSHIP, L.C., BAILEY, J.S., COX, N.A., STERN, N.J., BREWER, R. and WILLIAMS, O.** (1993) Two-step mucosal competitive exclusion flora treatment to diminish Salmonella in commercial broiler chickens. *Poultry Science* **72**: 1667-1762.
- BOLDER, N.M., VAN LITH, L.A.J.T., PUTIRULAN, F.F., JACOBS-REITSMA, W.F. and MULDER, R. W.A.W.** (1992) Prevention of colonization by *Salmonella enteritidis* PT4 in broiler chickens. *International journal of food microbiology* **15**: 313-317.
- BOTSOGLOU, N.A., FLOROU-PANER, P., CHRISTAKI, E., FLETOURIS, D.J. and SPAIS, A.B.** (2002) Effect of dietary oregano essential oil on performance of chickens and on iron-induced lipid oxidation of breast, thigh and abdominal fat tissues. *British Poultry Science* **43**: 223-230.
- BROZ, J. and BEARDSWORTH, P.** (2002) Recent trends and future developments in the use of feed enzymes in poultry nutrition, in: MCNAB, J. & BOORMAN, K.N. (Eds) *Poultry feedstuffs: Supply, composition and nutritive value*, pp. 345-361 (CAB international, Edinburgh, UK).
- CAVAZZONI, V., ADAMI, A. and CASTROVILLI, C.** (1998) Performance of broiler chickens supplemented with *Bacillus coagulans* as probiotic. *British Poultry Science* **39**: 526-529.
- CHOCT, M.** (2006) Enzymes for the feed industry: Past, present and future. *World's Poultry Science Journal* **62**: 5-15.
- CHOCT, M. and ANNISON, G.** (1992) The inhibition of nutrient digestion by wheat pentosans. *British Journal of Nutrition* **67**:123.
- CHRISTAKIA, E., FLOROU-PANERIA, P., GIANNENASA, I., PAPAZHARIADOU, M., BOTSOGLOU, N.A. and SPAISA, A.B.** (2004) Effect of a mixture of herbal extracts on broiler chickens infected with *Eimeria tenella*. *Animal Research* **53**: 137-144.
- CLARKSON, K., JONES, B., BOTT, R., BOWER, B., CHOTANI, G. and BECKER, T.** (2001) Enzymes: screening, expression, design and production, in: BEDFORD, M.R. (Ed.) *Enzymes in farm animal nutrition*, pp. 315-352 (CABI, Wiltshire).
- COLLINS, M.D. and GIBSON, G.R.** (1999) Probiotics, prebiotics, and symbiotics: approaches for modulating the microbial ecology of the gut. *American Journal of Clinical Nutrition* **69**: 1052S-1057S.
- COWAN, M.M.** (1999) Plant products as antimicrobial agents. *Clinical Microbiology Reviews* **12**: 564-582.
- CROSS, D.E., MCDEVITT, R.M., HILLMAN, K. and ACAMOVIC, T.** (2007) The effect of herbs and their associated essential oils on performance, dietary digestibility and gut microflora in chickens from 7 to 28 days of age. *British Poultry Science* **48**: 496-506.
- DALLOUL, R.A., LILLEHOJ, H.S., SHELLEM, T.A. and DOERR, J.A.** (2003) Enhanced mucosal immunity against *Eimeria acervulina* in broilers fed a *Lactobacillus*-based probiotic. *Poultry Science* **82**: 62-66.
- DANICKE, S., VAHJEN, W., SIMON, O. and JEROCH, H.** (1999) Effects of dietary fat type and xylanase supplementation to rye-based broiler diets on selected bacterial groups adhering to the intestinal epithelium, on transit time of feed, and on nutrient digestibility. *Poultry Science* **78**: 1292-1299.
- DECROOS, K., VERCAUTEREN, T., WERQUIN, G. and VERSTRAETE, W.** (2004) Repression of Clostridium population in young broiler chickens after administration of a probiotic mixture. *Communications in agricultural and applied biological sciences* **69**: 5-13.
- DIBNER, J.J. and RICHARDS, J.D.** (2005) Antibiotic growth promoters in agriculture: History and mode of action. *Poultry Science* **84**: 634-643.
- DROLESKEY, R.E., OYOFO, B.A., HARGIS, B.M., CORRIER, D.E. and DELOACH, J.R.** (1994) Effect of mannose on *Salmonella typhimurium*-mediated loss of mucosal epithelial integrity in cultured chick intestinal segments. *Avian Diseases* **38**: 275-281.
- DUGAS, B., MERCENIER, A., LENOIR-WIJNKOOP, I., ARNAUD, C., DUGAS, N. and POSTAIRE, E.** (1999) Immunity and prebiotics. *Immunology Today* **20**: 387-390.
- DUGUID, J.P., DAREKAR, M.R. and WHEATER, D.W.** (1976) Fimbriae and infectivity in *Salmonella typhimurium*. *Journal of Medical Microbiology* **9**: 459-473.
- EDENS, F.W.** (2003) An alternative for antibiotic use in poultry: probiotics. *Review. Brazilian Cienc. Aviculture* **5**: 75-97.

- ERTAS, O.N., GÜLER, T., ÇİFTÇİ, M., DALKILIÇ, B. and SIMSEK, Ü.G. (2005) The effect of an essential oil mix derived from oregano, clove and anise on broiler performance. *International Journal of Poultry Science* 4: 879-884.
- FARIA FILHO, D.E., TORRES, K.A.A., FARIA, D.E., CAMPOS, D.M.B. and ROSA, P.S. (2006) Probiotics for broiler chickens in Brazil: Systematic review and meta-analysis. *Brazilian Journal of Poultry Science* 8:89-98.
- FERKET, P.R. (1993) Practical use of feed enzyme for turkeys and broilers. *Journal of Applied Poultry Research* 2: 75-81.
- FERKET, P.R. (2004) Alternatives to antibiotics in poultry production: responses, practical experience and recommendations. *Nutritional biotechnology in the feed and food industries: Proceedings of Alltech's 20<sup>th</sup> Annual Symposium*, Kentucky, USA, 56-67.
- FERKET, P.R., PARKS, C.W. and GRIMES, J.L. (2002) Mannan oligosaccharides versus antibiotics for turkeys. *Nutritional biotechnology in the feed and food industries: Proceedings of Alltech's 18<sup>th</sup> Annual Symposium*, Kentucky, USA, 43-64.
- FERNANDEZ, F., HINTON, M. and VAN GILS, B. (2000) Evaluation of the effect of mannan-oligosaccharides on the competitive exclusion of *Salmonella enteritidis* colonization in broiler chicks. *Avian Pathology* 29: 575-581.
- FERNANDEZ, F., HINTON, M. and VAN GILS, B. (2002) Dietary mannan-oligosaccharides and their effect on chicken caecal microflora in relation to *Salmonella enteritidis* colonization. *Avian Pathology* 31: 49-58.
- FUKATA, T., SASAI, K., MIYAMOTO, T. and BABA, E. (1999) Inhibitory effects of competitive exclusion and fructo-oligosaccharide, singly and in combination, on *Salmonella* colonization of chicks. *Journal of Food Protection* 62: 229-233.
- FULLER, R. (1989) Probiotics in man and animals. *Journal of Applied Bacteriology* 66: 365-378.
- GAO, Y. and SHAN, A.-S. (2004) Effects of different oligosaccharides on performance and availability of nutrients in broilers. *Journal of Northeast Agricultural University* 11: 37-41.
- GARCIA, V., CATALA-GREGORI, P., HERNANDEZ, F., MEGIAS, M.D. and MADRID, J. (2007) Effect of formic acid and plant extracts on growth, nutrient digestibility, intestine mucosa morphology, and meat yield of broilers. *Journal of Applied Poultry Research* 16: 555-562.
- GAUTHIER, R. (2002) (XVIII Congreso Latinoamericano de Avicultura) (2002) Poultry Therapeutics: New alternatives. Accessed in 2006. [http://www.jefo.ca/pdf/ALA2003\\_en.pdf](http://www.jefo.ca/pdf/ALA2003_en.pdf).
- GHADBAN, G.S. (2002) Probiotics in broiler production - a review. *Archiv für Geflügelkunde* 66: 49-58.
- GIANNENAS, I., FLOROU-PANERI, P., PAPAZHARIADOU, M., CHRISTAKI, E., BOTSOGLOU, N. A. and SPAIS, A.B. (2003) Effect of dietary supplementation with oregano essential oil on performance of broilers after experimental infection with *Eimeria tenella*. *Archives of Animal Nutrition* 57: 99-106.
- GIBSON, G.R. and ROBERFROID, M. (1995) Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *Journal of Nutrition* 125: 1401-1412.
- GOREN, E., DE JONG, W.A., DOORNENBAL, P., BOLDER, N.M., MULDER, R.W.A.W. and JANSEN, A. (1988) Reduction of *Salmonella* infection of broilers by spray application of intestinal microflora: a longitudinal study. *The Veterinary Quarterly* 10: 249-255.
- GUO, F.C., KWAKKEL, R.P., WILLIAMS, B.A. ET AL. (2004a) Effects of mushroom and herb polysaccharides, as alternatives for an antibiotic, on growth performance of broilers. *British Poultry Science* 45: 684-694.
- GUO, F.C., KWAKKEL, R.P., WILLIAMS, B.A. ET AL. (2004c) Effects of mushroom and herb polysaccharides on cellular and humoral immune responses of *Eimeria tenella*-infected chickens. *Poultry Science* 83: 1124-1132.
- GUO, F.C., WILLIAMS, B.A., KWAKKEL, R.P., LI, ET AL. (2004b) Effects of mushroom and herb polysaccharides, as alternatives for an antibiotic, on the caecal microbial ecosystem in broiler chickens. *Poultry Science* 83: 175-182.
- HAVENAAR, R. and HUIS IN 'T VELD, M.J.H. (1992) Probiotics: a general view, in: WOOD, B.J.B. (Ed.) *Lactic acid bacteria*, Vol. 1. pp. 151-170 (Elsevier Applied Science Publishers, Amsterdam).
- HERNANDEZ, F., MADRID, J., GARCIA, V., ORENCO, J. and MEGIAS, M.D. (2004) Influence of two plant extracts on broiler performance, digestibility and digestive organ size. *Poultry Science* 83: 169-174.
- HIGGINS, J.P., HIGGINS, S.E., VICENTE, J.L., WOLFENDEN, A.D., TELLEZ, G. and HARGIS, B.M. (2007) Temporal effects of lactic acid bacteria probiotic culture on salmonella in neonatal broilers. *Poultry Science* 86: 1662-1666.
- HINTON, M., MEAD, G.C. and IMPEY, C.S. (1991) Protection of chicks against environmental challenge with *Salmonella enteritidis* by "competitive exclusion" and acid treated feed. *Letters In Applied Microbiology* 12: 69-71.
- HOOGE, D. (2004a) Meta-analysis of broiler chicken pen trials evaluating dietary mannan oligosaccharide, 1993-2003. *International Journal of Poultry Science* 3: 163-174.
- HOOGE, D. (2004b) Dietary alternatives for improving live performance of antibiotic-free poultry. Accessed in 2006. [www.feedinfo.com](http://www.feedinfo.com).

- HUANG, M.K., CHOI, Y.J., HOUDE, R., LEE, J.W., LEE, B. and ZHAO, X. (2004) Effects of *Lactobacilli* and an acidophilic fungus on the production performance and immune responses in broiler chickens. *Poultry Science* **83**: 788-95.
- HUANG, R.L., YIN, Y.L., LI, M. X. *ET AL.* (2007) Dietary oligochitosan supplementation enhances immune status of broilers. *Journal of the Science of Food and Agriculture* **87**: 153-159.
- HUANG, R.L., YIN, Y.L., WU, G.Y., ZHANG, Y.G., LI, T.J., LI, L.L., LI, M.X., TANG, Z.R., ZHANG, J., WANG, B., HE, J.H. and NIE, X.Z. (2005) Effect of dietary oligochitosan supplementation on ileal digestibility of nutrients and performance in broilers. *Poultry Science* **84**: 1383-1388.
- HUBENER, K., VAHJEN, W. and SIMON, O. (2002) Bacterial responses to different dietary cereal types and xylanase supplementation in the intestine of broiler chicken. *Archives of Animal Nutrition* **56**: 167-187.
- IJI, P.A., SAKI, A.A. and TIVEY, D.R. (2001) Intestinal structure and function of broiler chickens on diets supplemented with a mannan oligosaccharide. *Journal of the Science of Food and Agriculture* **81**: 1186-1192.
- JAMROZ, D., ORDA, J., KAMEL, C., WILICZKIEWICZ, A., WERTELECKI, T. and SKORUPINSKA, J. (2003) The influence of phytogetic extracts on performance, nutrient digestibility, carcass characteristics, and gut microbial status in broiler chickens. *Journal of Animal and Feed Sciences* **12**: 583-596.
- JAMROZ, D., WILICZKIEWICZ, A., ORDA, J., WERTELECKI, T. and SKORUPINSKA, J. (2004) (Electronic Journal of Polish Agricultural Universities) Response of broiler chickens to the diets supplemented with feeding antibiotic or mannanoligosaccharides. Accessed in 2006. <http://www.ejpau.media.pl/volume7/issue2/animal/art-06.html>.
- JANG, I.S., KO, Y.H., KANG, S.Y. and LEE, C.Y. (2007) Effect of a commercial essential oil on growth performance, digestive enzyme activity and intestinal microflora population in broiler chickens. *Animal Feed Science and Technology* **134**: 304-315.
- JIANG, H.Q., GONG, L.M., MA, Y.X., HE, Y.H., LI, D.F. and ZHAI, H.X. (2006) Effect of stachyose supplementation on growth performance, nutrient digestibility and caecal fermentation characteristics in broilers. *British Poultry Science* **47**: 516-522.
- JIN, L.Z., HO, Y.W., ABDULLAH, N. and JALALUDIN, S. (1997) Probiotics in poultry: Modes of action. *World's Poultry Science Journal* **53**: 351-368.
- JIN, L.Z., HO, Y.W., ABDULLAH, N. and JALALUDIN, S. (1998b) Growth performance, intestinal microbial populations and serum cholesterol of broilers diets containing *Lactobacillus* cultures. *Poultry Science* **77**: 1259-1265.
- JIN, L.Z., HO, Y.W., ABDULLAH, N. and JALALUDIN, S. (2000) Digestive and bacterial enzyme activities in broilers fed diets supplemented with *Lactobacillus* cultures. *Poultry Science* **79**: 886-891.
- JIN, L.Z., HO, Y.W., ABDULLAH, N., ALI, A.M. and JALALUDIN, S. (1998a) Effects of adherent lactobacillus cultures on growth, weight of organs and intestinal microflora and volatile fatty acids in broilers. *Animal Feed Science and Technology* **70**: 197-209.
- KALAVATHY, R., ABDULLAH, N., JALALUDIN, S. and HO, Y.W. (2003) Effects of *Lactobacillus* cultures on growth performance, abdominal fat deposition, serum lipids and weight of organs of broiler chickens. *British Poultry Science* **44**: 139-144.
- KLASING, K.C., LAURIN, D.E., PENG, R.K. and FRY, M. (1987) Immunologically mediated growth depression in chicks: influence of feed intake, corticosterone and interleukin-1. *Journal of Nutrition* **117**: 1629-1637.
- KLIS, F.M., MOL, P., HELLINGWERF, K. and BRUL, S. (2002) Dynamics of cell wall structure in *Saccharomyces cerevisiae*. *FEMS Microbiology Reviews* **26**: 239-247.
- KOCHER, A., RODGERS, N.J. and CHOCT, M. (2004) Efficacy of alternatives to IFAs in broilers challenged with *Clostridium perfringens*. In: *Proceedings of the Australian Poultry Science Symposium* **16**: 130-133.
- KOENEN, M.E., KRAMER, J., VAN DER HULST, R., HERES, L., JEURISSEN, S.H.M. and BOERSMA, W.J.A. (2004) Immunomodulation by probiotic lactobacilli in layer- and meat-type chickens. *British Poultry Science* **45**: 355 - 366.
- KUMAR, B.S., VIJAYASARATH, S.K., GOWDA, R.N.S. and SATYANARAYANA, M.L. (2002) Probiotics in the prevention of experimental fowl typhoid in broilers - A pathomorphological study. *Indian Journal of Animal Sciences* **72**: 528-531.
- KUMPRECHT, I. and ZOBAC, F. (1997) The effect of mannan-oligosaccharides in feed mixtures on the performance of broilers. *Zivocisna Vyroba* **42**: 117-124.
- LAN, Y., VERSTEGEN, M.W., TAMMINGA, S. and WILLIAMS, B.A. (2005) The role of the commensal gut microbial community in broiler chickens. *World's Poultry Science Journal* **61**: 95-104.
- LEE, K.W., EVERT, H., KAPPERT, H.J., FREHNER, M., LOSA, R. and BEYNEN, A.C. (2003) Effects of dietary essential oil components on growth performance, digestive enzymes and lipid metabolism in female broiler chickens. *British Poultry Science* **44**: 450-457.
- LEE, K.-W., EVERTS, H. and BEYNEN, A.C. (2004a) Essential oils in broiler nutrition. *International Journal of Poultry Science* **3**: 738-752.

- LEE, K.-W., EVERTS, H., KAPPERT, H.J., VAN DER KUILEN, J., LEMMENS, A.G., FREHNER, M. and BEYNEN, A.C. (2004b) Growth performance, intestinal viscosity, fat digestibility and plasma cholesterol in broiler chickens fed a rye-containing diet without or with essential oil components. *International Journal of Poultry Science* **3**: 613-618.
- LI, X., LIU, L.Q. and XU, C.L. (2008) Effects of supplementation of fructo-oligosaccharide and/or *Bacillus Subtilis* to diets on performance and intestinal microflora in broilers. *Archiv für Tierzucht* **51**: 64-70.
- LIU, J.R., LAI, S.F. and YU, B. (2007) Evaluation of an intestinal *Lactobacillus reuteri* strain expressing rumen fungal xylanase as a probiotic for broiler chickens fed on a wheat-based diet. *British Poultry Science* **48**: 507-514.
- LODDI, M.M., MORAES, V.M.B., NAKAGHI, L.S.O., TUCCI, F.M., HANNAS, M.I., ARIKI, J. and BRUNO, L.D.G. (2002) Mannan oligosaccharide and organic acids on intestinal morphology integrity of broilers evaluated by scanning electron microscopy. *Proceeding 11<sup>th</sup> European Poultry Science Conference*, Bremen, Germany, p.121.
- LYONS, T.P. (1994) Biotechnology in the feed industry: 1994 and beyond: A panorama of techniques, processes and products to address animal production problems today and tomorrow. *Biotechnology in the feed industry: Proceedings of Alltech's 10<sup>th</sup> Annual Symposium*, Kentucky, USA, 1-50.
- MCCRACKEN, V.J. and GASKINS, R.T. (1999) Probiotics and the immune system, in: TANNOCK, G.W. (Ed.) *Probiotics: A Critical Review*, pp. 85-111 (Horizon Scientific Press, Wymondham, UK).
- MITSCH, P., KOHLER, B., GABLER, C., LOSA, R. and ZITTERL-EGELSEER, K. (2002) CRINA poultry reduces colonisation and proliferation of *Clostridium perfringens* in the intestine and faeces of broiler chickens. Abstracts from the Eleventh European Poultry Conference, pp. 06-10, Bremen, Germany.
- MOHAN, B., KADIRVEL, R., NATARAJAN, A. and BHASKARAN, M. (1996) Effect of probiotic supplementation on growth, nitrogen utilization and serum cholesterol in broilers. *British Poultry Science* **37**: 395-401.
- MORROW, C. (2001) Solving the problems of necrotic enteritis. *British Poultry Science* **42**: S64-S67.
- MOUNTZOURIS, K.C., TSIRTSIKOS, P., KALAMARA, E., NITSCH, S., SCHATZMAYR, G. and FEGEROS, K. (2007) Evaluation of the efficacy of a probiotic containing *Lactobacillus*, *Bifidobacterium*, *Enterococcus*, and *Pediococcus* Strains in promoting broiler performance and modulating caecal microflora composition and metabolic activities. *Poultry Science* **86**: 309-317.
- MURRY, A.C., HINTON, A.J. and BUHR, R.J. (2006) Effect of botanical probiotic containing Lactobacilli on growth performance and populations of bacteria in the ceca, cloaca, and carcass rinse of broiler chickens. *International Journal of Poultry Science* **5**: 344-350.
- NEISH, A.S. (2002) The gut microflora and intestinal epithelial cells: A continuing dialogue. *Microbes and Infection*, **4**: 309-317.
- PANDA, A.K., RAMARAO, S.V., REDDY, M.R. and PRAHARAJ, N.K. (1999) Effect of dietary inclusion of probiotic on growth, carcass traits and immune response in broilers. *Indian Journal of Poultry Science* **34**: 343-346.
- PANDA, A.K., REDDY, M.R., RAMARAO, S.V., RAJU, M.V.L.N. and PRAHARAJ, N.K. (2000) Growth, carcass characteristics, immunocompetence and response to *Escherichia coli* of broilers fed diets with various levels of probiotic. *Archive für Geflügelkunde* **64**: 152-156.
- PASCUAL, M., HUGAS, M., BADIOLA, J.I., MONFORT, J.M. and GARRIGA, M. (1999) *Lactobacillus salivarius* CTC2197 prevents *Salmonella enteritidis* colonization in chickens. *Applied Environmental Microbiology* **65**: 4981-4986.
- PATTERSON, J.A. and BURKHOLDER, K.M. (2003) Application of prebiotics and probiotics in poultry production. *Poultry Science* **82**: 627-631.
- PRIVULESCU, M. (1999) "Humoral and cellular immune responses to Bio-Mos in piglets." National Institute of Veterinary Medicine, Pasteur Institute, Bucharest, Romania. Report to Alltech 4 pp.
- PRIYANKARAGE, N., SILVA, S.S.P., GUNARATNE, S.P., KOTHALAWALA, H., PALLIYAGURU, M. W.C.D. and GUNAWARDANA, G.A. (2003) Efficacy of probiotics and their effects on performance, carcass characteristics, intestinal microflora and Salmonella incidence in broilers. *British Poultry Science* **44**: S26-S27.
- RAVINDRAN, V. (2006) Broiler nutrition in New Zealand - Challenges and Strategies. Accessed in 2006. [www.feedinfo.com](http://www.feedinfo.com).
- ROSEN, G.D. (1995) Antibacterials in poultry and pig nutrition, in: WALLACE, R.J. & CHESSON, A. (Eds) *Biotechnology in Animal Feeds and Feeding*, pp. 143-172 (Weinheim, Germany).
- ROSEN, G.D. (2003) Setting and meeting standards for the replacement of pronutrient antibiotics in poultry. Accessed in 2008. [http://www.ces.ncsu.edu/depts/poulsci/conference\\_proceedings/nutrition\\_conference/2003/rosen\\_2003.pdf](http://www.ces.ncsu.edu/depts/poulsci/conference_proceedings/nutrition_conference/2003/rosen_2003.pdf).
- ROSEN, G.D. (2007) Holo-analysis of the efficacy of Bio-Mos® in broiler nutrition. *British Poultry Science* **48**: 21-26.

- SAMARASINGHE, K., WENK, C., SILVA, K.F.S.T. and GUNASEKERA, J.M.D.M. (2003) Turmeric (*Curcuma longa*) root powder and mannanoligosaccharides as alternatives to antibiotics in broiler chicken diets. *Asian-Australasian Journal of Animal Sciences* **16**: 1495-1500.
- SCHREZENMEIR, J. and DE VRESE, M. (2001) Probiotics, prebiotics, and symbiotics—approaching a definition. *American Journal of Clinical Nutrition* **73**: 361S-364S.
- SHAFEY, T.M., AL-MUFAREJ, S., SHALABY, M.I. and JARELNABI, A.J. (2001b) Effects of mannan oligosaccharides on antibody response to infectious bronchitis, infectious bursal disease and newcastle disease in chickens. *Journal of Applied Animal Research* **19**: 117-127.
- SHAFEY, T.M., AL-MUFAREJ, S., SHALABY, M.I. and JARELNABI, A.J. (2001a) The effect of feeding mannan-oligosaccharides (Bio-Mos) on the performance of meat chickens under two different vaccination programs. *Asian-Australasian Journal of Animal Sciences* **14**: 559-563.
- SIMON, O., JADAMUS, A. and VAHJEN, W. (2001) Probiotic feed additives—effectiveness and expected modes of action. *Journal of Animal and Feed Sciences* **10**: 51-67.
- SMIRNOV, A., PEREZ, R., AMIT-ROMACH, E., SKLAN, D. and UNI, Z. (2005) Mucin dynamics and microbial populations in chicken small intestine are changed by dietary probiotic and antibiotic growth promoter supplementation. *Journal of Nutrition* **135**: 187-192.
- SNEL, J., HARMSEN, H.J.M., VAN DE WIELEN, P.W.J.J. and WILLIAMS, B.A. (2002) Dietary strategies to influence the gastrointestinal microflora of young animals, and its potential to improve intestinal health, in: BLOK, M.C. (Ed.) *Nutrition and health on the gastrointestinal tract*, pp. 37-69 (Wageningen Academic Publishers, Wageningen, Netherlands).
- SONG, J.-Y. and LI, W.-F. (2001) The preparation of mannan-oligosaccharide from *Saccharomyces cerevisiae* and its effect on intestinal microflora in chicken. *Journal of Zhejiang University (Agric. and Life Sci.)* **27**: 447-450.
- SPRING, P. and PRIVULESCU, M. (1998) Mannanoligosaccharide: its logical role as a natural feed additive for piglets. *Biotechnology in the feed industry: Proceedings of Alltech's 14<sup>th</sup> Annual Symposium*, Kentucky, USA.
- SPRING, P., WENK, C., DAWSON, K.A. and NEWMAN, K.E. (2000) The effects of dietary mannanoligosaccharides on caecal parameters and the concentrations of enteric bacteria in the ceca of *Salmonella*-challenged broiler chicks. *Poultry Science* **79**: 205-211.
- STERN, N.J., BAILEY, J.S. and COX, N.A. (1998) Mucosal starter culture TM as part of an integrated program for *Salmonella* control on processed carcasses. *Proceedings of the 10th EPC*, Jerusalem, **II**:709-715.
- STERN, N.J., COX, N.A., BAILEY, J.S., BERRANG, M.E. and MUSGROVE, M.T. (2001) Comparison of mucosal competitive exclusion and competitive exclusion treatment to reduce *Salmonella* and *Campylobacter* spp. colonization in broiler chickens. *Poultry Science* **80**: 156-160.
- TEN BRUGGENCATE, S.J.M., BOVEE-UDENHOVEN, I.M.J., LETTINK-WISSINK, M.L.G. and VAN DER MEER, R. (2003) Dietary fructo-oligosaccharides dose-dependently increase translocation of salmonella in rats. *Journal of Nutrition* **133**: 2313-2318.
- TIMMERMAN, H.M., VELDMAN, A., VAN DEN ELSEN, E., ROMBOUTS, F.M. and BEYNEN, A.C. (2006) Mortality and growth performance of broilers given drinking water supplemented with chicken-specific probiotics. *Poultry Science* **85**: 1383-1388.
- VAHJEN, W., GLASER, K., SCHAFER, K. and SIMON, O. (1998) Influence of xylanase-supplemented feed on the development of selected bacterial groups in the intestinal tract of broiler chicks. *Journal of Agricultural Science* **130**: 489-500.
- VERDONK, J.M.A.J., SHIM, S.B., VAN LEEUWEN, P. and VERSTEGEN, M.W.A. (2005) Application of inulin-type fructans in animal feed and pet food. *British Journal of Nutrition* **93**: S125-S138.
- VIDANARACHCHI, J.K., MIKKELSEN, L.L., SIMS, I.M., IJI, P.A. and CHOCT, M. (2006) Selected plant extracts modulate the gut microflora in broilers. *Proceedings of Australian Poultry Science Symposium*, Sydney, Australia, **18**:145-148.
- WALDROUP, A.L., SKINNER, J.T., HIERHOLZER, R.E. and WALDROUP, P.W. (1993) An evaluation of fructo-oligosaccharide in diets for broiler chickens and effects on salmonellae contamination of carcasses. *Poultry Science* **72**: 643-650.
- WANG, R., LI, D. and BOURNE, S. (1998) Can 2000 years of herbal medicine history help us solve problems in the year 2000? *Biotechnology in the feed industry: Proceedings of Alltech's 14<sup>th</sup> Annual Symposium*, Kentucky, USA, 273-291.
- WATKINS, B.A. and KRATZER, F.H. (1984) Drinking water treatment with commercial preparation of a concentrated *Lactobacillus* culture for broiler chickens. *Poultry Science* **63**: 1671-1673.
- WILLIAMS, B.A., VERSTEGEN, M.W.A. and TAMMINGA, S. (2001) Fermentation in the large intestine of single-stomached animals and its relationship to animal health. *Nutrition Research Reviews* **14**: 207-227.
- WINDISCH, W. and KROISMAYR, A. (2006) The effects of phytobiotics on performance and gut function in monogastrics. Accessed in 2006. [www.feedinfo.com](http://www.feedinfo.com).
- XING, X.L. (2004) A comparative study on the effects of different Chinese herbal medicinal feed additives in broiler chickens. Accessed in 2006. [www.feedtrade.com.cn](http://www.feedtrade.com.cn).

- XU, Z.R., HU, C.H., XIA, M.S., ZHAN, X.A. and WANG, M.D. (2003) Effects of dietary fructo-oligosaccharides on digestive enzyme activities, intestinal microflora and morphology of male broilers. *Poultry Science* **82**: 1030-1036.
- XUE, M. and MENG, X.S. (1996) Review on research progress and prosperous of immune activities of bio-active polysaccharides. *Journal of Traditional Veterinary Medicine* **3**: 15-18.
- YANG, Y., IJI, P.A., KOCHER, A., MIKKELSEN, L.L. and CHOCT, M. (2007a) Effects of mannanoligosaccharide on growth performance, the development of gut microflora and gut function of broiler chickens raised on new litter. *Journal of Applied Poultry Research* **16**: 280-288.
- YANG, Y., IJI, P.A., KOCHER, A., MIKKELSEN, L.L. and CHOCT, M. (2008c) Effects of mannanoligosaccharides on the growth performance, nutrient digestibility and gut development of young birds given different cereal-based diets. *Animal Physiology and Animal Nutrition* (in press).
- YANG, Y., IJI, P.A., and CHOCT, M. (2007b) Effects of different dietary levels of mannanoligosaccharide on growth performance and gut development of broiler chickens. *Asian-Australasian Journal of Animal Sciences* **20**: 1084-1091.
- YANG, Y., IJI, P.A., KOCHER, A., MIKKELSEN, L.L. and CHOCT, M. (2008b) The response of broiler chickens to mannanoligosaccharide, or fructo-oligosaccharide-supplemented diet against pathogenic *Escherichia coli* challenge. *British Poultry Science* (accepted).
- YANG, Y., IJI, P.A., KOCHER, A., THOMSON, E., MIKKELSEN, L.L., and CHOCT, M. (2008a) Effects of mannanoligosaccharide in broiler chicken diets on growth performance, net energy utilization, nutrient digestibility, and intestinal microflora. *British Poultry Science* **49**: 186-194.
- YOUN, H.J. and NOH, J.W. (2001) Screening of the anticoccidial effects of herb extracts against *Eimeria tenella*. *Veterinary Parasitology* **96**: 257-263.
- YUSRIZAL, C. and CHEN, T.C. (2003) Effect of adding chicory fructans in feed on broiler growth performance, serum cholesterol, and intestinal length. *International Journal of Poultry Science* **3**: 214-219.
- ZHAN, X.-A., HU, C.-H. and XU, Z.-R. (2003) Effects of fructo-oligosaccharide on growth performance and intestinal microflora and morphology of broiler chicks. *Zhong Guo Shou Yi Xue Bao* **32**: 196-198.
- ZHANG, K.Y., YAN, F., KEEN, C.A. and WALDROUP, P.W. (2005) Evaluation of microencapsulated essential oils and organic acids in diets for broiler chickens. *International Journal of Poultry Science* **4**: 612-619.
- ZHANG, W.F., LI, D.F., LU, W.Q. and YI, G.F. (2003) Effects of isomalto-oligosaccharides on broiler performance and intestinal microflora. *Poultry Science* **82**: 657-663.
- ZULKIFLI, I., ABDULLAH, N., AZRIN, N.M. and HO, Y.W. (2000) Growth performance and immune response of two commercial broiler strains fed diets containing *Lactobacillus* cultures and oxytetracycline under heat stress conditions. *British Poultry Science* **41**: 593-597.