Genomic evaluation of crossbred dairy cattle in the United States

G.R. Wiggans,[1] P.M. VanRaden,[2] D.J. Null,[2] E.L. Nicolazzi,[1] G.B. Jansen,[1] and J.H. Megonigal, Jr.[1]

[1] Council on Dairy Cattle Breeding, Bowie, Maryland, USA

[2] Animal Genomics and Improvement Laboratory, Agricultural Research Service, USDA, Beltsville, Maryland, USA

CDCB
COUNCIL ON DAIRY CATTLE BREEDING

USDA
United States Department of Agriculture

Background

- By 2019, >35,000 animals excluded from genomic evaluation
  - Determined to be crossbred based on breed SNPs
  - >$1 million spent in genotyping with no genomic evaluation provided
- Evaluate crossbreds by combining individual-breed (purebred) SNP effects weighted by breed proportions
  [VanRaden et al., 2017, J. Dairy Sci. 100(Supp. 2):409–410]

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Crossbreds excluded before April 2019

- Genomic predictions for ~50 traits for 5 breeds:
  Ayrshire, Brown Swiss, Guernsey, Holstein, Jersey

- Excluded from evaluations based on presence of unlikely alleles in a set of breed-determining SNPs because:
  - Imputation relies on breed-specific haplotype libraries and allele frequencies (Holstein would overwhelm otherwise)
  - Marker effects differ by breed

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Requirements for evaluation of crossbreds

- Breed composition for weighting individual breed contributions
  - Pedigree often not reliable because of missing ancestors
- Evaluations for each breed on same base so that they can be combined
- Method to impute genotypes for crossbreds

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Breed proportions (breed base representation; BBR)

- Reference population of ~36,000 purebred males across 5 breeds
- SNP effects estimated with same 79K SNPs as in genomic evaluation
- Phenotypic values for 5 breed traits: 100% for animal's breed; 0% otherwise
- SNP-effect solutions updated each April
- BBR calculated and released for all animals every month
  - Adjusted to 0–100% range; values of ≤2 redistributed to other breeds
  - Animal BBR updated if any of its breed values differ by ≥4

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Imputation of crossbreds

- **Starts with Haplotype library from BBR reference population**
  - For Holsteins, bulls and cows with progeny, trad evaluations required (only 1/3 of cows with single progeny included)
  - All genotypes from other breeds
- **Parent genotypes from purebred directories added to improve imputation accuracy in crossbred directory**
- **PTAs released weekly; BBRs delayed until monthly so released BBR is from imputation using appropriate reference population**

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Evaluation of crossbreds

- 5 evaluations calculated for each trait of each animal (1 from each breed's SNP-effect solutions)
- Animals with highest BBR of <90% evaluated by weighting individual breed solutions by BBR
- Reference population limited to BBR of ≥94%
- Type, calving, health traits, and feed saved not blended because not comparable or not available for all breeds

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Frequency of crossbred genotypes



CDCB COUNCIL ON DAIRY CATTLE BREEDING

USDA

Evaluation details

- Results from purebred and crossbred directories combined so that animals with same evaluation breed included in same distribution file
- Animals with BBR of ≥85% for different breed not evaluated because of possible misidentification
- Inbreeding of future progeny calculated with relationships from purebred evaluations

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Weekly evaluations

- Approximate BBR calculated as genotypes are loaded; uses only SNPs on chip
- Genotypes with BBR of ≥85% initially placed in purebred directory
- Genotypes based on imputation from purebred directories with BBR of <90% moved to crossbred directory
- Genotypes in crossbred directory with BBR of ≥90% receive single-breed evaluation

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Breed of evaluation

- Initial breed of evaluation derived from preferred ID
- Remains breed of evaluation if BBR >40%; breed of highest BBR becomes breed of evaluation otherwise
- Change in breed of ID required if BBR of >85% for different breed
- Evaluations converted to individual breed bases for reporting

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Recessive conditions

- Not reported for crossbreds
- Most characteristics breed-specific

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Comparisons

- Little effect on purebred evaluations because of requirement for BBR of ≥94% for reference population
- Largest effect for animals with 90% < BBR <94% where own traditional evaluation not included
- Larger changes for animals previously with a single-breed evaluation and now with mixed breed ancestry reflected in evaluation

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Summary

- Genomic evaluations of crossbreds began April 2019
- Single-breed evaluations combined, weighted by breed proportions
- Number of crossbreds being genotyped rapidly increasing
- Routine updating of BBR when significant change
- Extends pedigree validation, provides breed composition, lower reliability than purebreds

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA

Acknowledgments and disclaimers

- Participating dairy producers for supplying pedigree and genomic data

- USDA-ARS project ARS 8042-31000-002-00, “Improving dairy animals by increasing accuracy of genomic prediction, evaluating new traits, and redefining selection goals” supported research and development

- Mention of trade names or commercial products is solely for the purpose of providing specific information and does not imply recommendation or endorsement by CDCB or USDA

- CDCB and USDA are equal opportunity providers and employers

CDCB COUNCIL ON DAIRY CATTLE BREEDING USDA