



APS Journals

The premier source for peer-reviewed plant pathology research since 1911.

Journals Home

APS Home

IS-MPMI Home

My Profile

Subscribe

Search

Advanced Search

Help



Subscribe Free alerts RSS



About the cover for August 2015

ISSN: 0191-2917

SEARCH

Enter Keywords

- Phytopathology
 Plant Disease
 MPMI

search

Advanced Search

Resources

About Plant Disease

First Look

Most Downloaded Articles

Submit a Manuscript

Customer Care

About My Password

Copyright and Permissions

Plagiarism and Ethics

Advertise

e-Xtra

Open Access

plant disease

Editor-in-Chief: Mark L. Gleason

Published by The American Phytopathological Society

Home > Plant Disease > Table of Contents > Full Text HTML

Previous Article Next Article

August 2015, Volume 99, Number 8

Page 1188

<http://dx.doi.org/10.1094/PDIS-02-15-0138-PDN>

DISEASE NOTES

First Report of the Cereal Cyst Nematode *Heterodera filipjevi* on Winter Wheat in Montana

A. T. Dyer, R. Al-Khafaji, and T. Lane, Department of Plant Sciences and Plant Pathology, Montana State University, Bozeman, MT 59717-3150; **T. Paulitz**, USDA-ARS, Pullman, WA 99164; and **Z. A. Handoo, A. M. Skantar, and D. J. Chitwood**, Nematology Laboratory, USDA-ARS, Beltsville, MD 20705.

Citation

Open Access.

Among the cereal cyst nematode complex, three species, *Heterodera avenae*, *H. filipjevi*, and *H. latipons*, are the most destructive to wheat (Smiley and Yan 2015). While *H. avenae* occurs in several U.S. states, *H. filipjevi* has only been reported from Washington in 2008 (Smiley and Yan 2015). *Heterodera latipons* has not been reported in the United States. In Chouteau County, Montana, a winter-wheat field reported to contain *H. avenae* was observed with areas of heavily stunted winter wheat 'Yellowstone'. Roots from affected plants displayed moderate numbers of nematode cysts. Given symptom severity, *H. filipjevi* was suspected (Golinowski et al. 1997). Living nematode cysts from wheat roots collected from the affected field were examined morphologically and by PCR for species identification. Observations of morphological characters critical for identification indicated that the specimens were *H. filipjevi* (Akinbade et al. 2010; Subbotin et al. 2010). Measurements of second-stage juveniles (J2, $n = 17$) included length of body (range = 490 to 615 μm , mean = 565.1 μm), stylet (24.0 to 25.0 μm , 24.3 μm) with anchor-shaped basal knobs, tail (50.0 to 70.0 μm , 61.4 μm), and hyaline tail terminus (35 to 43 μm , 38.5 μm). The lateral field had four lines, the inner two more distinct. Shapes of the tail, tail terminus, and stylet knobs were also consistent with *H. filipjevi*. The lemon-shaped cysts ($n = 10$) were light brown to yellow and had a zigzag pattern and bifenestrate vulval cone. Morphometrics included body length (L) excluding neck (475 to 718 μm , 643.0 μm); body width (W) (336 to 545 μm , 460.3 μm); L/W (1.3 to 1.6 μm , 1.4 μm); neck length (60 to 100 μm , 83.5 μm) and width (50 to 75 μm , 59.3 μm); fenestra length (40 to 55 μm , 48.0 μm) and μm width (20 to 30 μm , 24.8 μm); heavy underbridge (60 to 82.5 μm , 72.8 μm); vulval slit (7.5 to 12.5 μm , 9.4 μm); and many bullae. Molecular analysis of internal

Quick Links

Add to favorites

E-mail to a colleague

Alert me when new articles cite this article

Download to citation manager

Related articles found in APS Journals



Article History

Print: 5 Aug 2015

Ahead of Print: 26 Jun 2015

First Look: 25 Mar 2015

Accepted: 14 Mar 2015

This Journal is brought to you via a subscription from the DigiTop--USDA's Digital Desktop Library



CALL FOR PAPERS

SPECIAL Focus Issue

Noncoding RNA-Mediated Regulation of Plant-Microbe Interactions

Submissions Due: September 30, 2015



transcribed spacer (ITS1 and 2) and 28S rRNA confirmed identity as *H. filipjevi* (Akinbade et al. 2014). Amplicons generated from three J2 (two hatched from one cyst, one from another) were cloned and sequenced (Akinbade et al. 2014). Sequence from the 28S region (GenBank Accession No. KP878489) was >99.9% identical to those of *H. filipjevi* from China (GU083594, GU083592, and GU083597), and distinct from several populations of *H. avenae* (99.0 to 99.2%). The ITS rDNA from this population (KP878490) matched with 99.9% identity to several *H. filipjevi* sequences available from GenBank, including ones from Russia (AF274399), Tajikistan (AY148402), UK (AY148403), United States (GU079654), and China (GU083595). The next closest species match was *H. avenae*, with 96 to 97% identity to the Montana population. This is the first report of *H. filipjevi* from Montana. Eggs of *H. filipjevi* are noted for hatching in the fall during winter wheat planting, and therefore have the potential to affect winter wheat more than *H. avenae*, whose eggs hatch only in the spring after a cold period. Moreover, resistance in wheat against these two nematode species is different and will require additional considerations.

References:

Section:

- Akinbade, S. A.**, et al. 2014. *Plant Dis.* 98:1018. 10.1094/PDIS-09-13-0961-PDN
[\[Abstract\]](#) [\[ISI\]](#) [Open URL](#)
- Golinowski, W.**, et al. 1997. Page 80 in: *Cellular and Molecular Aspects of Plant-Nematode Interactions*. **C. Fenoll**, et al., eds. Kluwer, Dordrecht, the Netherlands.
[\[CrossRef\]](#) [Open URL](#)
- Smiley, R.**, and **Yan, G.** 2015. *Plant Dis.* 99:37610.1094/PDIS-08-14-0789-RE.
[\[Abstract\]](#) [Open URL](#)
- Subbotin, S. A.** 2010. *Systematics of Cyst Nematodes, Vol. 8 Part B*. **Mundo-Ocampo M'** et al.' eds. Leiden' Brill, the Netherlands' [Open URL](#)

[Citation](#)

