

COLORADO WATER RESOURCES RESEARCH INSTITUTE
Information Series No. 105
Joe E. Brummer, Editor
March, 2008



**PROCEEDINGS
HIGH ALTITUDE REVEGETATION
WORKSHOP NO. 18
MARCH 2008**

HIGH ALTITUDE REVEGETATION WORKSHOP

Proceedings

NO. 18

**Colorado State University
Fort Collins, Colorado
March 4-6, 2008**

Edited by

**Joe E. Brummer
Colorado State University
Department of Soil and Crop Sciences
1170 Campus Delivery
Fort Collins, CO 80523**

**Information Series No.105
Colorado Water Resources Research Institute
Colorado State University**

PREFACE

The 18th biennial High Altitude Revegetation Workshop was held at the Hilton Fort Collins in Fort Collins, Colorado on March 4-6, 2008. The Workshop was organized by the High Altitude Revegetation (HAR) Committee in conjunction with the Departments of Soil and Crop Sciences and Forest, Rangeland and Watershed Stewardship at Colorado State University. The Workshop was well attended this year by 241 people from a broad spectrum of universities, government agencies, and private companies. Discussions centered on the revegetation of disturbed lands always seem to be of interest to many people as evidenced by the number in attendance this year at the Workshop. The HAR Workshop is somewhat unique in that it focuses on the practical, on-the-ground application of revegetation techniques. People come away from the Workshop with new information and new ideas that they can take home and apply directly to their specific situations.

This Workshop would never happen without the dedication and contributions from the many people on the HAR Committee. This is an all volunteer organization and everyone that contributed to this years Workshop is to be commended for their efforts.

The Committee tried something different this year and sent out a solicitation for volunteer papers instead of inviting speakers as was done in the past. This approach worked very well and we would like to thank all the people who took time to prepare not only a presentation or poster, but also a paper or abstract for inclusion in these proceedings. The proceedings consist of 21 papers and 3 abstracts grouped into 8 workshop sessions, 6 poster papers, and 3 poster paper abstracts. In addition to the papers and posters presented on March 5 and 6, a special Practical Revegetation Session was held on March 4 at Rocky Mountain National Park. This session was well received and attended by approximately 80 people.

For current information on upcoming High Altitude Revegetation Committee events, visit our website at: www.highaltitudereveget.org.

Joe E. Brummer
Editor



TABLE OF CONTENTS

ii

PREFACE

iii

TABLE OF CONTENTS

KEYNOTE SESSION

1

The Southwest Ecological Restoration Institutes: Filling the Information Gap
Dan Binkley, Director, Colorado Forest Restoration Institute

2

Alternatives to Topsoil: A Five-Year Case Study in Mine Site Revegetation
Steven L. McGeehan, Analytical Sciences Laboratory
University of Idaho, Moscow, ID

OIL, GAS AND MINING RECLAMATION

Chaired by: Mike Ellis
Phil Barnes
Bryce Romig

32

A Test Plot Design to Evaluate Organic Amendments and Planting Densities on Overburden Minesoils at a Limestone Quarry in Tijeras, New Mexico
Kenneth Carlson, Habitat Management Inc., Englewood, CO

49

Evaluating Seeding Techniques and Native Plant Establishment in The Pineedale Anticline, Wyoming
Susan Winslow, Bridger Plant Material Center, USDA Natural Resources Conservation Service
Bridger, MT

74

Burlington Mine VCLUP Case History An Ecological Approach to Mine Site Remediation
Maureen O'Shea-Stone, Walsh Environmental
Boulder, CO

WETLAND AND RIPARIAN AREA CONCERNS

Chaired by: Ray Sperger
Mindy Wheeler

iii



David Johnson, President, Western Ecological Resource
Boulder, CO
90

Restoration of Subalpine Wetlands White River National Forest
Natasha Davis, Forest Rangeland & Watershed Stewardship Dept.
Colorado State University, Fort Collins, CO
135

*Revegetation of Fluvial Mine Tailing Deposits: The Use of Five
Riparian Shrub Species*
Jennifer Patterson, Engineering and Hydrosystems Inc.
Douglas County, Colorado
Littleton, CO
155

Chaired by: Denise Arthur
Carl Mackey

*Stapleton's Northfield Ponds- Landscape Architecture, Ecology and
Engineering*
Jayne Kopperl, EDAW/AECOM
Denver, CO
166

*Science-Based Approach to Revegetation of the Summitville Mine
Superfund Site: From Greenhouse Screening to Site-Wide Reclamation*
Julie Rieder, Restoration Ecology Lab
Colorado State University, Fort Collins, CO
178

*Response of Transplanted Aspen to Drip Irrigation on Reclaimed
Mine Lands*
Roy Karo, Seneca Coal Co.
Hayden, CO
190

BANQUET SPEAKER

When the Grass Stood Stirrup-High
Dave Bradford, Rangeland Management Specialist with the Grand Mesa,
Uncompahgre and Gunnison National Forests
221

DISTURBED SITE RESTORATION I

Chaired by: Jody Nelson
Mark Paschke

Promoting Plant Diversity with Low Broadcast Seeding Rates Applied
 to Amended Overburden Minesoils at a Limestone Quarry in Tijeras,
 New Mexico
Robin Bay, Habitat Management Inc.
 Englewood, CO

310

OIL, GAS AND MINING RECLAMATION II

Mt. Goliath Revegetation Project at the Dos Chappel Nature Center
Mark Fusco, Denver Botanic Gardens
 Denver, CO

303

High Altitude Native Restoration at the Winter Park Resort
Denise Arthur, ESCO Associates
 Boulder, CO

285

Design and Construction of the Shell Trenches Alternative RCRA
 Cover at Rocky Mountain Arsenal, Commerce City, Colorado
Carl Mackey, Washington Group International
 Commerce City, CO

274

Chaired By: Jody Nelson
 Mark Paschke

DISTURBED SITE RESTORATION II

Revegetation of the Rocky Flats Site Colorado
Jody Nelson, DOE Legacy Management/S.M. Stoller Corp.
 Westminster, CO

258

Establishing Native Plants on Abandoned Farmland
Cynthia Brown, Bioagricultural Sciences and Pest Management Dept
 Colorado State University, Fort Collins, CO

232

Evaluation of Gully Erosion Control and Restoration Techniques
 in North Crystal Creek Basin, Pikes Peak, Colorado
Eric Billmeyer, Rocky Mountain Field Institute
 Colorado Springs, CO

222

POSTER PAPERS

- Restoring Sage Grouse Habitat in the Pinedale Anticline Gas Field, Wyoming: Shell/BLM Revegetation Pilot Project**
Richard Carr, C-M Environmental Group
Pinedale, WY
329
- Visual Resource Consideration in the Reclamation Equation**
John McCarty, USDI Bureau of Land Management
Washington, DC
360
- AGENTS OF CHANGE**
Chaired by: Jeff Connor
Steve Spaulding
- Mountain Pine Beetle and Spruce Beetle Outbreaks Changing our High Elevation Forests in Colorado**
Sheryl Costello, Forest Health Protection, USDA Forest Service
Golden, CO
383
- Cooperation: Key to Success in Controlling Dalmatian Toadflax in the Upper South Fork of the Shoshone River, Wyoming**
Bob Parsons, Park County Weed and Pest Control District
Powell, WY
384
- Vegetational Future of Colorado Mountain Vegetation-Recovery Following an Episode of Heavy Infestation by Trees**
David Buckner, ESCO Associates
Boulder, CO
397
- The Application of Ecological Principles to Accelerate Reclamation of Well Pad Sites**
Joshua Eldridge, Department of Forest, Rangeland and Watershed Management, Colorado State University
Fort Collins, CO
403
- Biosolids Use for Reclaiming Fluvial Mine Tailings**
C.L. Freeman, Department of Soil & Crop Sciences, Colorado State University
Fort Collins, CO
417

511	COMMITTEE LIST
509	SUMMER TOURS
493	PARTICIPANT LIST
472	Paonia, CO Michael Ward , Michael Ward Outdoors <i>Mountain Coal Company, West Elk Mine, Dry Fork of Minnesota Creek, Reclamation Photography Project, 2004-2007</i>
471	Questa Mine, NM Chevron Mining Inc. Anne Wagner , PhD, Manager, Environmental and Public Policy, <i>History of Reclamation at the Questa Mine - A Case History</i>
461	Akron, CO Plains Area Maysoon Mikha , USDA-ARS Great Plains Research Station, Northern <i>Soils in the Central Great Plains Region</i> <i>Best Management Practices for Remediation/Restoration of Degraded</i>
460	Jackson, WY District Rachel Markko , Natural Resource Specialist, Teton Conservation <i>Wyoming</i> <i>Karn's Meadow Wetland Rehabilitation Along Flat Creek in Jackson</i>
459	Fort Collins, CO State University Stewardship, Warner College of Natural Resources, Colorado Julie Knudson , Department of Forest, Rangeland and Watershed <i>Canada Thistle Infested Site</i> <i>Soil Fertility Manipulation and Grass Seeding for Restoration of a</i>
451	Gardnerville, NV Ed Kleiner , President, Comstock Seed Co. <i>Parallel Trends in the Reclamation Industry</i>
438	Boulder, CO Scientist and Engineers George Greenwood , Wildfire Mitigation, Walsh Environmental <i>Community Wildfire Protection Project Implementation Case Study</i> <i>The Yegge Road Shaded Fuel Break: A Jefferson County</i>

BEST MANAGEMENT PRACTICES FOR REMEDIATION/RESTORATION OF DEGRADED SOILS IN THE CENTRAL GREAT PLAINS REGION

Maysoon M. Mikhah¹, Merle F. Vigil¹, Joseph G. Benjamin¹, Dave J. Poss¹,
and Francisco J. Calderon¹

¹USDA-ARS, Central Great Plains Research Station, Northern Plains Area,
40335 CR GG, Akron, CO 80720

ABSTRACT

Farmlands in the Central Great Plains Region (CGPR) have lost topsoil through wind and water erosion induced by tillage and poor soil management (Wheat-fallow management). Productivity of degraded/eroded soils can be restored using organic amendment such as manure and improved crop and soil management. Our objectives are to: (i) identify optimal rates of manure to supply nutrients to typical dryland crops in the CGPR; (ii) determine the rate of improvement of soil physical and chemical properties associated with manure amendment/management; and (iii) quantify the difference in restoration of eroded soils using manure as an amendment versus managing those same soils with legume grass mixtures and chemical fertilizer. The experiment, established off on a farmer field near Akron, Colorado has a randomized complete block design with crops/soils managed using manure amendment compared with soils/crops managed with commercial fertilizer. Treatments include a tillage variable (deep plow, shallow sweep, and no-tillage), manure and commercial nitrogen rates (none, low and high). Changes in soil physical, chemical, and biological properties as well as grain yield are evaluated every year. The preliminary data (for one growing season) suggests that manure addition increases the productivity of eroded soils in the Central Great Plain Region. In subsequent years this experiment (after multiple manure applications) could result in changes in soil parameters and increased yield. This report will provide "benchmark" measurements of the treatments being studied and first year grain and biomass yields.

INTRODUCTION

Farmlands in the CGPR have lost topsoil through wind and water erosion induced by tillage and poor soil management (wheat-fallow). These soils are now degraded with low soil quality and productivity. Some of the soil quality parameters that are affected by poor soil management include: soil compaction, infiltration rate, soil water holding capacity, soil nutrient exchange capacity, soil aggregation and aggregate stability, organic carbon build up, soil pH, and soil microbial ecology. In addition to wind and water erosion, numerous studies have indicated that soil degradation is a result of soil organic matter lost through increased soil disturbance and decomposition (Angers et al., 1993; Lal et al., 1995). Productivity and quality of degraded/eroded soils can be restored using manure and improved management. Manure amendment is a management practice that can improve the nutrient status of the soil (Vitosh et al., 1997) and increase soil organic carbon levels (Mikhah and Rice, 2004). Aoyama et al.

Corresponding author (Maysoon.Mikhah@ars.usda.gov)

(1999a) observed an increase in soil organic matter with addition of manure and the subsequent increase the formation of slaking-resistant soil aggregates. Aoyama et al. (1999b) concluded that applying manure contributed to the accumulation of aggregates-protected C and N. Similarly, Mikha and Rice (2004) reported that aggregate-protected labile carbon and nitrogen was significantly greater with manure amendment when compared with chemical fertilizer treatment. The combination of no-till management with manure amendment further increased the formation and stabilization of soil aggregates and increased the physical protection of soil carbon and nitrogen (Mikha and Rice, 2004; Jiao et al., 2006).

Tillage practices can alter soil organic matter and effect soil erosion. Tillage practices can reduce soil organic matter by (i) increasing residue mixing into the soil which increases aeration and enhances residue decomposition, (ii) destroying soil aggregate and exposing previously protected soil organic matter to soil fauna, and (iii) increasing losses due to soil erosion (Blevins and Frye, 1993; Egghall et al., 1994; Lal et al., 1994; Six et al., 1999). Many studies have shown that increased soil organic matter with no-tillage management (Carter, 1992; Beare et al., 1994; Six et al., 1999) improves soil aggregation and aggregate-associated soil organic matter (Mikha and Rice, 2004).

Organic amendment such as manure can have a positive effect on soil quality by improving soil porosity and preventing soil crust formation (Pagliai et al., 2004). Continuous manure application over several years, can have a positive effect by reducing soil bulk density (Miller et al., 2002; McVay et al., 2006). Reduction in soil bulk density and greater soil porosity are clear indicators for reduced soil compaction, improved aeration, greater infiltration and improved conditions for plant root penetration. Frye and Blevins (1997) reported that improving plant productivity is related to enhanced root growth, crop yield, and plant biomass. Also, the addition of organic material as a nitrogen source (manure and/or compost) can mitigate the negative effects of excessive tillage on grain yield and soil organic carbon conservation (Egghall and Power, 1999; Singer et al., 2004; Mando et al., 2005). Similarly Egghall et al. (2004) documented, the residual effect of increased nutrient availability due to multiple years of manure application increase corn grain yield for one growing season and influence soil properties for several years (Egghall et al., 2004).

The impact of manure applications and tillage practices on plant productivity has been studied intensively for more than 30 years. However, the effects of multiple years of beef manure application combined with different tillage systems on improving soil quality parameters and the productivity of eroded soil are not well documented in dryland systems. The objectives of this study are: (i) Identify optimal rates of beef manure to supply nitrogen and phosphorus to typical dryland crops in the CGPR; (ii) determine the rate of improvement of crop yield, soil physical, chemical, and biological properties associated with dryland manure management of eroded soils; and (iii) Quantify the advantages of restoring eroded soils using manure as an amendment versus managing those same soils with chemical fertilizer.

MATERIALS AND METHODS

Site Management

The experimental area consists of 4.8 hectares while the plot area consists of 1.7 hectares. The large remainder is due to requiring large alleys to manage full size farm equipment. Individual experimental units (plots) are 13.7 m wide and 15.2 m long (45 feet wide and 50 feet long). The experimental design is a randomized complete block with four replicates. The replicates are arranged such that three are parallel to each other across the slope. The fourth is split with the first ten plots adjacent to replicate one and the last ten plots adjacent to replicate two.

Cropping Sequence

Crop rotations used are typical to the Central Great Plains Region. The rotation currently being used is corn (2006) – proso Millet (2007) – forage Winter Triticale (2008). The crop rotation in subsequent years will be decided according to weather pattern (temperature and precipitation). The crops are normally planted on all of the plots and alleys ways. In 2006, before manure and tillage treatments were established, corn was planted through all plots in the east west direction (perpendicular to planting in 2007 and 2008) in a plant 2 skip 2 row configuration.

Manure Application

Manure is applied in the fall to allow for winter precipitation to restore moisture lost during tillage operations. Manure is applied using a Meyer spreader. This manure spreader was used at low RPM to obtain a uniform spread width of 2.7 m. Calibration of the manure spreader was performed by driving the manure spreader over a tarp and then weighing the manure collected on the tarp. The rear gate was left open during application and rates were controlled by changing ground speed. Beef manure was obtained from a local feedlot. Samples of the manure “piles” to be used were taken and analyzed for nutrients to determine amount of nutrients applied to the plots.

There is a low and a high manure rate applied for each tillage treatment. The low was determined by estimating the amount of nitrogen required to meet crop needs average over the next six years. The high rate is two times the low rate for fertilizer and three times the low rate for manure. Realizing that this is in excess of crop nutrient needed this high rate was used to ensure an increase in the amount of organic carbon applied. One of our hypotheses is that these higher rates of manure will significantly increase soil organic matter and improve soil physical properties in these plots.

Table 1. Manure Factors relative to treatment and tillage depth.

Tillage Treatment		Frequency		kg/ha ----	
				Low	
				High	
No-tillage	Annual	1X	3X	1X	3X
Sweeps (13 cm depth)	Annual	1X	3X	1X	3X
Deep tillage (36 cm depth)	Bi-Annual	2X	6X	2X	6X
Deep Tillage	One time	6X	18X	6X	18X

Precipitation in 2007 was less than normal (Figure 1) especially during the critical period of millet germination. Crop soil water used during the growing season was calculated by adding the amount of precipitation (from planting (Jun 22nd) to harvesting (September 26th)) to the differences between soil water content at planting and harvesting. Throughout the growing season, crop soil water use ranged between 18.1 cm to 27.7 cm integrated to 120 cm deep soil profile (Table 3). Soil water contents at planting (Jun 22nd, 2007) ranged between 15%-19% in the surface 15 cm (data not shown). Similarly, soil water contents in the whole 120 cm profile ranged from 15% to 19.1% (data not shown). In general, soil water contents at field capacity ranged 17% to 20% depending specific soil type. This indicates that soil water contents were adequate for planting, but a lack of precipitation during June and July may have reduced final

RESULTS AND DISCUSSIONS

Plot Operations
Manure was applied November 16th, 2006 for the 2007 proso millet crop. The tillage treatments for the manure plots were performed the same week just after manure application. The tillage treatments for the fertilizer and control plots were performed the following week because of time constraints. The fertilizer for these plots was not applied until the spring, immediately prior to planting (Jun 22nd, 2007). The deep tilled plots were packed in the spring to firm the soil to ensure good seed to soil contact at planting time. Spray operations were performed at and prior to planting proso millet to control weeds. No herbicide applications were done after the proso millet was sown. The proso millet was sown with a plot harvester after 95% of the head had changed from green to mature. The rest of the plots were sown soon after. The proso was picked up with a plot combine two week later after it had sufficiently dried.

Crop	Fertilizer Application		Manure Application	
	High	Low	1X	18X
Date	22-Jun-07	16-Oct-07	20-Nov-07	16-Oct-07
kg N/ha	60	30	65	0
kg N/ha	60	30	83	160
kg N/ha	260	458	260	458
kg N/ha	1342	1342	0	0

Table 2. Fertilizer and manure application rates by treatment.

Five levels of manure were applied to the plots in fall 2006 depending on the treatment (Table 1). A 1X rate is applied to plots for which there is an annual application on a low fertilizer/manure treatment. The rate factor applied was based on frequency of application and the rate. For example the rate for the Deep tillage, one time treatment was determined by taking the 1X rate times six (since we are estimating fertility needs for the next six years) for the low manure nitrogen rate, while the high rate of the same treatment would have an 18X rate. The amount of manure (that meet the nitrogen requirement) for different treatment combination is presented in Table 2.

Table 3: Soil inorganic nitrogen (kg N ha^{-1}) and plant total soil water used (cm) during 2007 growing season integrated to 120 cm depth.

Table 3: Soil inorganic nitrogen (kg N ha⁻¹) and plant total soil water used (cm) during 2007 growing season integrated to 120 cm depth.

Tillage		Nitrogen type		Preplant -----		--- Growing Season ---	
		$\text{NH}_4^+\text{-N}$	$\text{NO}_3^-\text{-N}$	----- kg N ha ⁻¹ -----		----- cm -----	
No tillage	3X [†] manure	11.0	27.5	26.0	27.7	25.6	24.3
	High [†] fertilizer	10.8	21.8	27.7	25.6	25.6	24.3
	1X ^{††} manure	10.9	25.2	25.6	25.6	25.6	24.3
	Low ^{††} fertilizer	10.9	21.5	24.3	24.3	24.3	24.3
	Control [†]	11.1	15.8	24.1	24.1	24.1	24.1
	3X manure	11.8	46.1	25.5	25.5	25.5	25.5
	High fertilizer	10.7	18.1	25.6	25.6	25.6	25.6
	1X manure	10.5	25.3	26.1	26.1	26.1	26.1
	Low fertilizer	10.7	11.4	25.9	25.9	25.9	25.9
	Control	10.7	17.8	26.3	26.3	26.3	26.3
Deep tillage (2y) [#]	6X manure	12.0	72.7	23.3	23.3	23.3	23.3
	2X manure	12.5	53.0	22.4	22.4	22.4	22.4
	18X manure	11.8	128.5	22.1	22.1	22.1	22.1
	6X manure	12.1	108.1	23.2	23.2	23.2	23.2
	High fertilizer	10.8	22.1	18.2	18.2	18.2	18.2
Deep tillage ^{&}	High fertilizer	10.8	22.1	18.2	18.2	18.2	18.2
	Low fertilizer	11.1	44.0	18.1	18.1	18.1	18.1

no-tillage and manure amendment increased millet yield by 27% (at low nitrogen rate) and by 20% (at high nitrogen rate) compared with the combination of sweep-tillage and manure




Figure 1. Monthly precipitation for 2006, 2007, and 98 years average at Akron, Colorado.

treatment (Figure 2). Millet grain yield increased as manure nitrogen rate increased by 17% when managed with no-tillage and 23% when managed with sweep tillage. The greater grain yields associated with no-tillage combined with manure amendment could be due to soil water conservation during the early growth stage of millet. Deep-tillage showed a reduction in millet





Figure 2. Proso millet yield (Mg ha^{-1}) as affected by nitrogen rate and different tillage practices. (NT): no-tillage; (Deep-2y): manure application once and equivalent to 2 years N rate and tillage at 36 cm depth; (Deep-6y): manure application once and equivalent to 6 years N rate and tillage at 36 cm depth; and (Deep-fertilizer): nitrogen application equivalent to 1 year N rate and tillage at 36 cm depth.

Proso millet biomass production exhibited a similar trend as the grain yield especially with the combination of no-till and manure amendment and sweep-tillage manure (Figure 3). High N rate translated to high plant biomass production. High crop water use translated to high yield except with sweep-tillage manure and deep-2y manure, where high yield were associated with low water usage (Figure 4). There were no changes in crop water use in relation to high and low grain yields associated with deep-fertilizer treatment.

Figure 3. Proso millet biomass (Mg ha^{-1}) as affected by nitrogen rate and different tillage practices. (NT): no-tillage treatment; (Deep-2y): manure application once and equivalent application equivalent to 6 years N rate and tillage at 36 cm depth; (Deep-6y): manure application once and equivalent application equivalent to 1 year N rate and tillage at 36 cm depth.



yield as the manure nitrogen rate increases. There was 10% and 29% reduction in millet yield as manure nitrogen rate increased from 83 to 260 (kg N ha^{-1}) and from 458 to 1342 (kg N ha^{-1}), respectively. However, across tillage (sweep and deep tillage) a maximum yield (with manure treatment) was observed (represented by dash line in Figure 2) around 260 (kg N ha^{-1}) where the yield was reduced as manure nitrogen increase thereafter (Figure 2). The excessive amount of manure nitrogen that was applied with deep-2y and deep-6y in combination with low precipitation could have reduced millet growth and grain yield. Grain yield increased with the combination of no-till and fertilizer (by 4%) and deep-tillage (by 5%), while a reduction was observed with sweep-tillage (by 7%) as the rate of inorganic fertilizer increased from 30 to 60 kg N ha^{-1} (Figure 2). Because this is just the first year of production for this experiment, no explanation can be given for the reduction of grain yield for sweep-tillage with high fertilizer rate compared with low fertilizer rate. Overall, grain yields were 44% greater with no-tillage combined with manure amendment than with no-tillage and fertilizer.


Angers, D.A. A. N'Dayegamiye, and D. Cote. 1993. Tillage-induced differences in organic matter of particle-size fractions and microbial biomass. *Soil Sci. Soc. Am. J.* 57:512-516.

REFERENCES

We would like to thanks all the technicians at the Central Great Plains Research Station: Karen Couch, Brandon Peterson, Linda Hardesty, Donna Fritzler, Stacey Poland, Gene Uhler, Delbert Koch, Michael Pappas, and Paul Campbell for their help in manure application, planting, harvesting, and collecting and analyzing soil and plant samples.

ACKNOWLEDGMENTS

Figure 4. Relationship between proso millet yield (Mg ha^{-1}) and total soil water used (cm) throughout the 2007 growing season as affected by nitrogen rate and different tillage practices. (*) represents high grain yield associated with high water usage and (†) represents high grain yield associated with low water usage.



In summary, the preliminary data suggests that manure additions could improve the productivity of unproductive (eroded) soils in the CGPR. No specific explanations can be given for treatment differences since we have only one growing season and the environmental factors (temperature, precipitation, and evapotranspiration) could have more effect on final grain yields than our imposed treatments. In subsequent years it will be important to determine the improvement in different soil parameters and to document yield effects from different management practices. Several additional "benchmark" measurements (physical, chemical and biological) are being made on the soils in these plots and these measurements will be repeated periodically throughout the duration of the experiment.

- Aoyama, M., D.A. Anger, and A. N'Dayegamiye. 1999a. Particulate and mineral-Associated organic matter in water-stable aggregates as affected by mineral fertilizer and manure applications. *Can. J. Soil Sci.* 79:295-302.
- Aoyama, M., D.A. Angers, A. N'Dayegamiye, and N. Bissonnette. 1999b. Protected organic matter in water-stable aggregates as affected by mineral fertilizer and manure applications. *Can. J. Soil Sci.* 79:419-425.
- Beare, M.H., P.F. Hendrix, and D.C. Coleman. 1994. Water-stable aggregates and organic matter fractions in conventional and no-tillage soils. *Soil Sci. Soc. Am. J.* 58:777-786.
- Blevins, R.L., and W.W. Frye. 1993. Conservation tillage: An ecological approach to soil management. p. 33-78. *In: D.L. Sparks (ed), Adv. in Agro.*, v. 51. Botner, P. 1985. Response of microbial biomass to alternate moist and dry conditions in a soil incubated with ^{14}C - and ^{15}N -labelled plant material. *Soil Biol. Biochem.* 17:329-337.
- Carter, M.R. 1992. Influence of reduced tillage systems on organic matter, microbial biomass, macro-aggregate distribution and structural stability of the surface soil in humid climate. *Soil Tillage Res.* 23:361-372.
- Egball, B., D. Ginting, and J. E. Gilley. 2004. Residual effects of manure and compost applications on corn production and soil properties. *Agron. J.* 96: 442-447.
- Egball, B., N. L.N. Mielke, D.L. McCallister, and J.W. Doran. 1994. Distribution of organic carbon and nitrogen in a soil under various tillage and crop sequences. *J. Soil Water Cons.* 49:201-205.
- Egball, B. and J. F. Power. 1999. Composted and noncomposted manure application to conventional and no-tillage system: corn yield and nitrogen uptake. *Agron. J.* 91: 819-825.
- Frye, W.W., and R.L. Blevins. 1997. Soil organic matter under long-term no-tillage and conventional tillage corn production in Kentucky. p. 227-234. *In* Paul, E.A., K. Paustian, E.A. Elliott, C.V. Cole (eds.), *Soil organic matter in temperate agroecosystems: Long-term experiments in North America*, CRC Press, Boca Raton, FL.
- Jiao, You, J.K. Whalen, and W.H. Hendershot. 2006. No-tillage and manure applications increase aggregation and improve nutrient retention in a sandy-loam soil. *Geoderma* 134: 24-33.
- Kladivko, E.J. 2001. Tillage systems and soil ecology. *Soil Till. Res.* 61:61-76.
- Lal, R., J. Kimble, and B.A. Stewart. 1995. World soil as a source or sink for radiatively-active gases. p. 1-8 *In: R. Lal et al., J. Kimble, E. Levine, and B.A. Stewart (eds.), Soil management and greenhouse effect*. Lewis Publ., Boca Raton, FL.
- Lal, R., A.A. Mahboubi, and N.R. Fausey. 1994. Long-term effect on Mollic Ochraqualf in northwestern Ohio. III. Soil nutrient profile. *Soil Tillage Res.* 15:371-382.

- Mando, A., B. Ouattara, M. Sedogo, L. Stroosnijder, K. Ouattara, L. Brussaard, and B. Vanlauwe. 2005. Long-term effect of tillage and manure application on soil organic fractions and crop performance under Sudano-Sahelian conditions. *Soil & Tillage Research* 80:95-101.
- McVay, K.A., J.A. Budde, K. Fabrizzi, M.M. Mikha, C.W. Rice, A.J. Schlegel, D.E. Peterson, D.W. Sweeney, and C. Thompson. 2006. Management effects on soil physical properties in long-term tillage studies in Kansas. *Soil Sci. Soc. Am. J.* 70:434-438.
- Mikha, M.M. and C.W. Rice. 2004. Tillage and manure effects on soil and aggregate-associated carbon and nitrogen. *Soil Sci. Soc. Am. J.* 68:809-816.
- Miller, J. J., N. J. Sweetland, and C. Chang. 2002. Soil physical properties of a Chernozemic clay loam after 24 years of beef cattle manure application. *Can. J. Soil Sci.* 82: 287-296.
- Pagliai, M., N. Vignozzi, and S. Pellegrini. 2004. Soil structure and the effect of management practices. *Soil & Tillage Research* 79:131-143.
- Paustian, K., H.P. Collins, and E.A. Paul. 1997. Management controls on soil carbon. p.15-49. *In* Paul, E.A., K. Paustian, E.A. Elliott, and C.V. Cole (eds.), *Soil organic matter in temperate agroecosystems: Long-term experiments in North America*, CRC Press, Boca Raton, FL.
- Singer, J.W., K.A. Kohler, M. Liebman, T.L. Richard, C.A. Cambardeila, and D.D. Buhler. 2004. Tillage and Compost Affect Yield of Corn, Soybean, and Wheat and Soil Fertility. *Agron. J.* 96:531-537.
- Six, J., E.T. Elliott, and K. Paustian. 1999. Aggregate and soil organic matter dynamic under conventional and no-tillage systems. *Soil Sci. Soc. Am. J.* 63:1350-1358.
- Tisdall, J.M. 1996. Formation of soil aggregates and accumulation of soil organic matter. p. 57-96. *In*: M.R. Carter and B.A. Stewart (eds.), *Structure and soil organic matter storage in agricultural soils*. Adv. Soil Sci., Lewis Publ., Boca Raton, FL.
- Vorish, M.L., R.E. Lucas, and G.H. Silva. 1997. Long-term effect of fertilizer and manure on corn yield, soil carbon, and other soil chemical properties in Michigan. p. 129-139. *In* Paul, E.A., K. Paustian, E.A. Elliott, C.V. Cole (eds.), *Soil organic matter in temperate agroecosystems: Long-term experiments in North America*, CRC Press, Boca Raton, FL.