

CENTRAL PLAINS EXPERIMENTAL RANGE (CPER)





Location and Climate

The Central Plains Experimental Range (CPER) is located on the northern end of the semi-arid shortgrass steppe in the western Great Plains. At 5,400 feet above sea-level, CPER is located 7.5 miles northeast of Nunn, Colorado, within the <u>USDA</u> <u>Northern Plains Climate Hub</u> region.

Historic Temperature & Precipitation

Historic average annual temperature in Weld County, CO, is 47°F (1979-2019). Long-term average annual precipitation at CPER is 13 inches (1940-2019), but has ranged from 4" (in 1964) to 23" (in 1967). Precipitation in April, May, and June is most important for vegetation growth on the short-grass prairie, but rainfall in late-July and August may stimulate a second pulse of green-up, especially if the spring season was dry. Precipitation across CPER's 15,500 acres is highly variable, through space and time, including both within and across years (Augustine 2010).

Growing Season at CPER

The growing season, roughly May-Sept, is ~133 days. Rangelands at CPER produce an average of 658 lbs/ acre per year (1983-2018), ranging from a 35-year low of 98 lbs/ac (in the 2002 drought) to a high of 1415 lbs/ac (in the wet year of 2009).

USDA is an equal opportunity provider and employer.

LTAR Network and <u>USDA Climate Hubs</u> are working to develop knowledge and technology for sound resource management via research and collaboration with stakeholders. The goal is to ensure sustained crop and livestock production and ecosystem services from agroecosystems, and to forecast and verify the effects of environmental changes, public policies, and emerging technologies.



Monthly average precipitation and temperature (min & max) at CPER (1981-2010).(credit: <u>Climate Toolbox</u>).



Scientists and stakeholders tour the CPER in early November (Photo credit: Dannele Peck)

Long-Term Data Collection



Data has been collected at CPER since 1939. ARS scientists use

this long-term data to evaluate relationships among climate, livestock performance, grassland species diversity, and more (<u>Davis et al. 2020, Derner et al.</u> <u>2020, Raynor et al. 2020</u>).



Stocker cattle grazing the Central Plains Experimental Range near Nunn, CO (Photo credit: David Augustine)

Extreme Weather: Droughts & Deluges

The semi-arid CPER has short, mild summers and long, cold and dry winters. During the summer, convective thunderstorms form regularly over nearby mountains, creating the potential for heavy downpours and severe hail. Drought is also common in the region, as shown in the time-series (2000-2019) below of drought severity in a local watershed.











-10% -5% 0 +5% +10% +15% +20% Projected precipitation changes (%) for Colorado's Northeastern Plains by 2035-2065 (vs. 1971–2000; RCP 4.5) (Source: Lukas et al. 2014)

Climate and Climate Change

The figure on the left shows 3 variables across the wet growing season of 2009, which produced 1415 lbs/ac of vegetation (215% of the 35-year average, 1983-2018). The 3 variables are: [top panel] NDVI, a measure of greenness and proxy for how well vegetation is growing; [middle panel] soil moisture at various depths; and [bottom panel] precipitation.

The bottom-left figure shows the same 3 variables for 2012, the 4th driest growing season in 35 years (1983-2018), which produced 269 lbs/ac (41% of the average).

Implications for Rangelands

- Precipitation in Colorado's Northeastern Plains is expected to increase in winter, but decrease in summer (see figure above, adapted from <u>Lukas et al. 2014</u>).
- Warming and elevated CO2 may cause plants to green up, peak, and decline earlier in the season, yet persist longer during dry summer and autumn seasons (Blumenthal et al. 2018).
- Elevated CO2 may increase forage quantity in this region, but also decrease forage quality, especially in drier years (Augustine et al. 2018).
- In a warmer climate, invasive plant species (e.g., cheatgrass) may become more competitive (Blumenthal et al. 2016).
- Rangeland livestock managers will need to become more flexible and adaptive (<u>Reeves et al. 2017, Derner et al. 2018</u>).



Visit: https://ltar.ars.usda.gov/sites/cper/ CPER Site Lead: justin.derner@usda.gov CPER Ecohydrologist: david.hoover@usda.gov NP Climate Hub: dannele.peck@usda.gov

Date Modified: December 4, 2020 10:37 AM

Websites Referenced

USDA Climate Hubs: <u>https://www.climatehubs.usda.gov/</u> USDA Northern Plains Climate Hub: <u>https://www.climatehubs.usda.gov/hubs/northern-plains</u> Annual Precipitation at CPER: <u>https://www.ars.usda.gov/plains-area/fort-collins-co/center-for-agricultural-resourcesresearch/rangeland-resources-systems-research/docs/range/misc-pages/cper-precipitation/</u> Drought Monitor time-series: <u>https://droughtmonitor.unl.edu/Data/Timeseries.aspx</u> Climate Toolbox: <u>https://climatetoolbox.org/tool/Historical-Climograph</u> USDA ARS LTAR CPER (Central Plains Experimental Range): <u>https://ltar.ars.usda.gov/sites/cper/</u>

Literature Cited & Additional References

Augustine, D.J. 2010. Spatial versus temporal variation in precipitation in a semiarid ecosystem. Landscape Ecology 25:913-925. https://doi.org/10.1007/s10980-010-9469-y

Augustine, D.J., Blumenthal, D.M., Springer, T.L., LeCain, D.R., Gunter, S.A., Derner, J.D. 2018. *Elevated CO2 induces* substantial and persistent declines in forage quality irrespective of warming in mixedgrass prairie. *Ecological Applications* 28:721-735. https://doi.org/10.1002/eap.1680

Blumenthal, D.M., Kray, J.A., Ortmans, W., Ziska, L.H., Pendall, E. 2016. *Cheatgrass is favored by warming but not CO2 enrichment in a semi-arid grassland. Global Change Biology* 22:3026-3038. *https://doi.org/10.1111/gcb.13278*

Blumenthal, D.M., Mueller, K.E., Kray, J.A., LeCain, D.R., Pendall, E., Duke, S., Zelikova, T.J., Dijkstra, F.A., Williams, D.G., Morgan, J.A. 2018. *Warming and elevated CO2 interact to alter seasonality and reduce variability of soil water in a semiarid grass-land. Ecosystems* 21:1533-1544. https://doi.org/10.1007/s10021-018-0237-1

Davis, K.P., Augustine, D.J., Monroe, A.P., Derner, J.D., Aldridge, C.L. 2020. Adaptive rangeland management benefits grassland birds utilizing opposing vegetation structure in the shortgrass steppe. Ecological Applications 30(1):e02020. https://doi.org/10.1002/eap.2020

Derner, J., Briske, D., Reeves, M., Brown-Brandl, T., Meehan, M., Blumenthal, D., Travis, W., Augustine, D., Wilmer, H., Scasta, et al. 2018. Vulnerability of grazing and confined livestock in the Northern Great Plains to projected mid- and latetwenty-first centu-ry climate. Climatic Change 146:19-32. https://doi.org/10.1007/s10584-017-2029-6

Derner, J.D., Raynor, E.J., Reeves, J.L., Augustine, D.J., Milchunas, D.G. 2020. *Climatic and management determinants of large herbivore production in semiarid grassland. Agriculture, Ecosystems & Environment 290:106761. https://www.sciencedirect.com/science/article/pii/S0167880919303779*

Edwards, B.L., Webb, N.P., Brown, D.P., Elias, E., Peck, D.E., Pierson, F.B., Williams, C.J., Herrick, J.E. 2019. *Climate change impacts on wind and water erosion on US rangelands. Journal of Soil and Water Conservation* 74(4):405-418. https://www.jswconline.org/content/74/4/405/tab-article-info

Frankson, R., Kunkel, K., Stevens, L., Easterling, D. 2017. Colorado State Climate Summary. NOAA Technical Report NESDIS 149-CO, 4 pp. https://statesummaries.ncics.org/chapter/co/

Jencso, K., Parker, B., Downey, M., Hadwen, T., Howell, A., Rattling Leaf, J., Edwards, L., Akyuz, A., Kluck, D., Peck, D., et al. 2019. Flash Drought: Lessons Learned from the 2017 Drought Across the U.S. Northern Plains and Canadian Prairies. NOAA Na-tional Integrated Drought Information System. https://www.drought.gov/drought/documents/flash-drought-lessons-learned-2017-drought-across-us-northern-plains-and-canadian-0

Lukas, J., Barsugli, J., Doesken, N., Rangwala, I., Wolter, K. 2014. Climate Change in Colorado: A Synthesis to Support Water Re-sources Management and Adaptation. Western Water Assessment, Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado Boulder. https://wwa.colorado.edu/climate/co2014report/

Raynor, E.J., Derner, J.D., Hoover, D.L., Parton, W.J., Augustine, D.J. 2020. Large-scale and local climatic controls on large herbi-vore productivity: implications for adaptive rangeland management. Ecological Applications 30(3):02053. https://doi. org/10.1002/eap.2053

Reeves, M.C., Bagne, K.E., Tanaka, J. 2017. *Potential climate change impacts on four biophysical indicators of cattle production from western US rangelands. Rangeland Ecology & Management 70:529-539. https://www.fs.fed.us/rm/pubs_journals/2017/rmrs_2017_reeves_m001.pdf*

Reynolds, A.Q., Derner, J.D., Augustine, D.J., Porensky, L.M., Wilmer, H., Jorns, T., Briske, D.D., Scasta, J.D., Fernández-Giménez, M.E., 2019. *Ecological sites: can they be managed to promote livestock production? Rangelands* 41(6):239-243. https://doi.org/10.1016/j.rala.2019.07.003

Shrum, T.R., Travis, W.R., Williams, T.M., Lih, E. 2018. *Managing climate risks on the ranch with limited drought information. Climate Risk Management 20:11-26. https://www.sciencedirect.com/science/article/pii/S2212096317301407*