

# THE REVERSIBLE INHIBITION OF MYOBLAST PROLIFERATION BY $\gamma$ -HEXACHLOROCYCLOHEXANE<sup>1</sup>

M. Koohmaraie<sup>2</sup>, J. E. Schollmeyer<sup>3</sup> and T. R. Dutson<sup>4</sup>

U.S. Department of Agriculture<sup>5</sup>  
Clay Center, NE 68933

PURCHASED BY THE UNITED STATES  
DEPARTMENT OF AGRICULTURE FOR  
OFFICIAL USE.

## ABSTRACT

The effects of treating non-fusing myoblast variants, fu-1 and M<sub>3</sub>A, with two levels ( $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M) of  $\gamma$ -hexachlorocyclohexane, an inhibitor of phosphatidylinositol synthesis, on myoblast proliferation were evaluated by measuring myoblast proliferation (counting cells) and visual inspection via phase microscopy. In the presence of  $\gamma$ -hexachlorocyclohexane, these cells were arrested, presumably in G<sub>1</sub>. The inability of these cells to replicate did not appear to be due to a toxic effect of  $\gamma$ -hexachlorocyclohexane, because these cells were capable of resuming proliferation once they were transferred to media lacking  $\gamma$ -hexachlorocyclohexane. Cells were grown in media containing myo-[2-<sup>3</sup>H]inositol and the radioactive content of water-soluble metabolites, the end product of phosphatidylinositides hydrolysis, was quantitated. Cells that were grown in the presence of  $\gamma$ -hexachlorocyclohexane, in addition to the loss of proliferative ability, also contained significantly less water-soluble metabolites. Therefore, it appears that there is a direct relationship between phosphatidylinositol metabolism and cell proliferation in the cell lines studied. (Key Words: Cell Culture, Cell Growth, Phosphatidylinositols.)

## Introduction

Since the initial isolation of phosphatidylinositol from the brain (Folch, 1949), a great deal has been learned about these unique and minor constituents of the cellular phospholipids (for review, see Berridge, 1984). It is generally accepted that there are three species of phosphatidylinositols (phosphatidylinositol, phosphatidylinositol 4-phosphate, phosphatidylinositol 4,5-bisphosphate), and that these species are interconvertible through phosphorylation (Berridge, 1984; Berridge et al., 1984). It is also well-documented that the hydrolysis of these phospholipids leads to the mobilization of

intracellular Ca<sup>++</sup> (Rebecchi et al., 1983; Berridge et al., 1984; Biden et al., 1984; Burgess et al., 1984a,b; Irvine et al., 1984; Joseph et al., 1984a,b; Kolesnick and Gershengorn, 1984; Aub and Putney, 1985). Among the different species of inositol phosphates generated through hydrolysis of phosphatidylinositols (Berridge, 1984), inositol 1,4,5-triphosphate seems to serve as the secondary messenger for the mobilization of Ca<sup>++</sup> from the intracellular stores, particularly those associated with the endoplasmic reticulum (Streb et al., 1983; Burgess et al., 1984a,b; Dawson and Irvine, 1984; Joseph et al., 1984a,b; Prentki et al., 1984).

The function(s) of the secondary messenger, i.e., the mobilized Ca<sup>++</sup>, generated through this pathway remains unclear. Michell (1979) proposed that a change in cytosol Ca<sup>++</sup> concentration can exert an appreciable effect on the progress of cell proliferation, and that proliferation was initiated by a cytosol Ca<sup>++</sup> concentration in excess of a certain critical level (Kishimoto et al., 1980). Other studies have shown that rate of phosphatidylinositol hydrolysis is directly proportional to rate of chicken ventricular cell proliferation (Ishima et al., 1982), and that turnover of phosphatidylinositol in rapidly dividing lens epithelial cells (t<sub>1/2</sub> approximately 5 h) was much greater than

<sup>1</sup> Michigan Agr. Exp. Sta. Journal Article No. 11910.

<sup>2</sup> Postdoctoral Fellow, Dept. of Food Sci. and Human Nutr., Michigan State Univ., stationed at Roman L. Hruska U.S. Meat Anim. Res. Center, Clay Center, NE 68933.

<sup>3</sup> Roman L. Hruska U.S. Meat Anim. Res. Center, Clay Center, NE 68933.

<sup>4</sup> Dept. of Food Sci. and Human Nutr., Michigan State Univ., East Lansing, MI 48824.

<sup>5</sup> Mention of a trade name, proprietary products or specific equipment does not constitute a guarantee or warranty of the product by the USDA and does not imply its approval to the exclusion of other products that may also be suitable.

Received October 21, 1985.

Accepted January 30, 1986.

in non-proliferating and differentiating fiber cells ( $t_{1/2}$  approximately 63 h; Zelenka, 1980). Moreover, Ristow et al. (1973) observed enhanced incorporation of  $^{32}\text{P}$  and  $^3\text{H}$ -inositol into phosphatidylinositol when  $G_1$ -arrested embryonic rat fibroblasts were stimulated to grow by addition of serum. It was also observed that up to 15 to 20% of the total radioactivity of embryonic rat fibroblasts prelabelled with  $^3\text{H}$ -inositol was eliminated from the phospholipid during 20 to 40 min after the addition of serum (Ristow et al., 1975).

Because other studies suggested that enhanced phosphatidylinositol turnover is related to cell proliferation, we investigated the relationship between cell growth and phosphatidylinositol hydrolysis utilizing  $\gamma$ -hexachlorocyclohexane, a drug with the same configuration as myo-inositol and known to inhibit the synthesis of phosphatidylinositol (Hokin and Brown, 1969; Fisher and Mueller, 1971; Ristow et al., 1973; Hoffmann et al., 1980; Vu et al., 1983; Zelenka and Vu, 1984). Hoffman et al. (1980) reported that  $\gamma$ -hexachlorocyclohexane inhibits phosphatidylinositol synthesis as well as DNA synthesis and mitosis. Our experiments using non-fusing myoblast variants indicate these cells lose the ability to divide when phosphatidylinositol synthesis is prevented by  $\gamma$ -hexachlorocyclohexane. Moreover, we have demonstrated that these cells will regain their ability to proliferate if they are cultured in the absence of  $\gamma$ -hexachlorocyclohexane, even after extended exposure to  $\gamma$ -hexachlorocyclohexane.

#### Materials and Methods

**Cell Culture Conditions.** Fu-1, a non-fusing variant of  $L_8$  myoblasts was obtained from Dr. Kaufman (University of Illinois) and  $M_3A$ , a non-fusing variant of  $L_6$  myoblasts was obtained from Dr. Schubert (Salk Institute). Growth characteristics of these cells were described by Kaufman et al. (1980) and Schubert and Lacorbienic (1982), respectively. Cell lines were cloned and maintained by serial passage as described by Kaufman et al. (1980) and Schubert and Lacorbienic (1982). Cells

were grown in Dulbecco's modified Eagle's Medium<sup>6</sup> supplemented with 10% fetal calf serum (DMEM + 10% FBS) at 37 C in a 10%  $\text{CO}_2$  high-humidity atmosphere.

**Cell Growth.** Approximately  $5 \times 10^4$  cells were plated in 35-mm<sup>2</sup> tissue culture dishes<sup>7</sup> in 2 ml of DMEM + 10% FBS and allowed to attach. After attachment, the media was removed and new media corresponding to the appropriate treatment added. Control media consisted of DMEM + 10% FBS and .035% dimethylsulfoxide (DMSO). To study the effect of blocking phosphatidylinositol synthesis on the cell growth, both cell lines were grown in DMEM + 10% FBS containing  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane<sup>8</sup>.  $\gamma$ -hexachlorocyclohexane was dissolved in DMSO and appropriate volumes were added to the media to give the desired final concentrations of  $\gamma$ -hexachlorocyclohexane, i.e.,  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M. This media was then sterilized by filtration (.2  $\mu\text{M}$ , Millipore) and added to the dishes (both control and treatments contained .035% DMSO). Twenty-four hours after original cell plating, media was removed and new media corresponding to appropriate treatment was added to the dishes. At 0, 24, 48, 72 and 96 h after the original plating, cells were trypsinized (.05% trypsin in  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ -free Earle's Salt Solution) and counted in a ZBI Coulter Counter<sup>9</sup>.

**Labelling and Extraction of Water-Soluble Metabolites.** At the same time cells were plated to measure growth rate, parallel sets of 35-mm<sup>2</sup> plates were prepared by the same procedure for labelling and extraction of water-soluble metabolites according to the procedure of Berridge et al., 1982. Four hours after plating, media was removed and fresh media, corresponding to the appropriate treatment (control,  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane), containing 1  $\mu\text{Ci}/\text{plate}$  of myo-[2- $^3\text{H}$ ]inositol (specific activity 14 Ci/mmol, Amersham) was added. Twenty-four hours after plating, the media was removed and fresh media corresponding to appropriate treatment and containing myo-[2- $^3\text{H}$ ]inositol (1  $\mu\text{Ci}/\text{plate}$ ) was added to the dishes. After 72 h of pre-labelling with myo-[2- $^3\text{H}$ ]inositol, the labelled media was removed, cells were rinsed twice and incubated for 1 h with 2 ml of phosphate-buffered saline (PBS), pH 7.4 (Berridge et al., 1984). After an additional rinse with 2 ml of PBS, the reaction was stopped by addition of 1 ml of ice-cold 15% trichloroacetic

<sup>6</sup> Gibco, Grand Island, NY.

<sup>7</sup> Costar, Cambridge, MA.

<sup>8</sup> Sigma Chemical Co., St. Louis, MO.

<sup>9</sup> Coulter Elect. Inc., Hialeah, FL.

acid (TCA). The dishes were then left on ice for 30 min. Cells were then harvested with rubber policemen and centrifuged at  $2,000 \times g$  for 5 min. This procedure was repeated two additional times with the addition of 1 ml of  $H_2O$  each time for maximum extraction of the water-soluble metabolites. The supernatants from the TCA extractions and  $H_2O$  washes were combined and TCA was removed by extraction with diethylether (four extractions). The final extract was neutralized and the  $^3H$  content was determined.

### Results and Discussion

Results presented in figures 1 and 2 document the effect of  $\gamma$ -hexachlorocyclohexane on the proliferative capacities of the non-fusing myoblast variants, fu-1 and  $M_3A$ . Growth curves for control cells are similar to those of Kaufman et al. (1980) for fu-1 and Schubert and Lacorbienic (1982) for  $M_3A$  cells lines. These results (figures 1 and 2) clearly indicate that the presence of  $\gamma$ -hexachlorocyclohexane in the media negatively affected the ability of these cells to divide. For example, for  $M_3A$  cells, cell population in the media lacking  $\gamma$ -hexachlorocyclohexane increased 7.6 times after 96 h of incubation, but only 2.4 and 2.1 times in the presence of  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane, respectively. The same analysis for fu-1 cells were 7.5, 2.6

and 2.1 times for control,  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane, respectively.

Hoffmann et al. (1980) reported that  $3 \times 10^{-4}$  M concentration of  $\gamma$ -hexachlorocyclohexane was near the toxic level for rat embryo fibroblasts and led to significant cell death in some of their experiments. In order to ensure that the effects of  $\gamma$ -hexachlorocyclohexane observed in our experiments (figures 1 and 2) was not due to cell death, we monitored the appearance of the treated cells via phase microscopy (figures 3 and 4). Although cell proliferation was substantially attenuated, the cells remained viable and did not become refractile and release from the substrate with  $\gamma$ -hexachlorocyclohexane treatment. Moreover, when the  $\gamma$ -hexachlorocyclohexane media was removed after 96 h and was replaced by fresh media (DMEM + 10% FBS alone), we were able to reverse the inhibitory effect of  $\gamma$ -hexachlorocyclohexane (figure 5) and show a dramatic increase in  $M_3A$  proliferation after 48 h in normal media. The fact that these cells regained normal proliferative rates suggest that the  $\gamma$ -hexachlorocyclohexane arrested the cells in  $G_1$ , stage of cell cycle. Phase micrographs of cells after 96 h of incubation in the presence of  $\gamma$ -hexachlorocyclohexane (figure 6a) and 48 h in normal media (figure 6b) show the ability of these cells to regain their normal proliferative properties after the removal of  $\gamma$ -hexachlorocyclohexane. In addition to resuming normal




Figure 1. Growth curve of fu-1 cells in the presence and absence of  $\gamma$ -hexachlorocyclohexane.




Figure 2. Growth curve of  $M_3A$  cells in the presence and absence of  $\gamma$ -hexachlorocyclohexane.




Figure 3. Phase contrast micrograph of fu-1 cells grown in the presence and absence of  $\gamma$ -hexachlorocyclohexane ( $\times 120$ ). a, b and c represent control,  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane, respectively at 0 h. d, e and f represent control,  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane, respectively at 96 h.




Figure 4. Phase contrast micrograph of M<sub>3</sub>A cells grown in the presence and absence of  $\gamma$ -hexachlorocyclohexane ( $\times 120$ ). a, b and c represent control,  $1 \times 10^{-4}$  and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane, respectively at 0 h. d, e and f represent control,  $1 \times 10^{-4}$  M and  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane, respectively at 96 h.




Figure 5. Growth curve of M<sub>3</sub>A cell grown in the presence of  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane, then in its absence. Arrow indicates start of growth in the absence of inhibitor.

proliferative activities, removal of  $\gamma$ -hexachlorocyclohexane resulted in generation of water-soluble inositol metabolites at a rate similar to that in control cells (table 1). The results of this experiment provide evidence that the effect of  $\gamma$ -hexachlorocyclohexane could not be due to a toxic effect.

Although the hexachlorocyclohexane-treat-

ed cells remained viable even after 96 h of treatment, treated cells, both fu-1 and M<sub>3</sub>A cells, displayed substantial morphological differences when compared with the control cells. Fu-1 cells appear to be more elongated, flattened and somewhat larger (figures 3e and f). This latter impression may result from the fact that these cells are more flattened. Treated M<sub>3</sub>A cells also exhibit an altered morphology (figures 4e and f), again demonstrating some elongation and extensive pseudopod formation. M<sub>3</sub>A cells treated for 96 h with  $2 \times 10^{-4}$  M hexachlorocyclohexane (figure 6a) appear almost arborized. However, these cells, when refed with normal media for 48 h, certainly have regained their proliferative ability and morphologically are indistinguishable from control M<sub>3</sub>A cells seen in figure 4d. The results presented in figures 3 and 4 constitute the first evidence documenting morphological changes associated with  $\gamma$ -hexachlorocyclohexane treatment.

Causes of the morphological changes observed in treated cells were not investigated in this study. However, other researchers have noted that diacylglycerol, an end-product of phosphatidylinositol hydrolysis, may effect membrane enzymes and can induce membrane fusion (Das and Rand, 1984). Thus, it is possible that by inhibiting phosphatidylinositol metabolism, thereby reducing the amount of




Figure 6. Phase micrograph ( $\times 120$ ) of M<sub>3</sub>A cells grown in presence of  $2 \times 10^{-4}$  M  $\gamma$ -hexachlorocyclohexane (96 h; a) and then in its absence (48 h; b).

TABLE 1. EFFECT OF  $\gamma$ -HEXACHLOROCYCLOHEXANE ON THE LEVELS OF WATER-SOLUBLE INOSITOL METABOLITES<sup>a</sup>

Cell line	Treatment	Count $\cdot$ min <sup>-1</sup> $\cdot$ dish <sup>-1</sup>
Fu-1	Control	8,321 $\pm$ 155
Fu-1	1 $\times$ 10 <sup>-4</sup> M $\gamma$ -hexachlorocyclohexane	2,432 $\pm$ 110
Fu-1	2 $\times$ 10 <sup>-4</sup> M $\gamma$ -hexachlorocyclohexane	588 $\pm$ 28
M <sub>3</sub> A	Control	9,584 $\pm$ 238
M <sub>3</sub> A	1 $\times$ 10 <sup>-4</sup> M $\gamma$ -hexachlorocyclohexane	1,108 $\pm$ 61
M <sub>3</sub> A	2 $\times$ 10 <sup>-4</sup> M $\gamma$ -hexachlorocyclohexane	553 $\pm$ 22

<sup>a</sup>The results are expressed as radioactivity in the total inositol phosphate, and are means  $\pm$  SE for five determinations in a single experiment. Similar results were found in four other experiments.

diacylglycerol available, we have in some way altered the plasma membranes.

Assay of the effects of  $\gamma$ -hexachlorocyclohexane on phosphatidylinositol metabolism showed that, at 1  $\times$  10<sup>-4</sup> M  $\gamma$ -hexachlorocyclohexane, the <sup>3</sup>H content (i.e., water-soluble metabolites) in fu-1 and M<sub>3</sub>A cells was 29.2 and 7.0%, respectively, of that of control cells (table 1). At 2  $\times$  10<sup>-4</sup> M  $\gamma$ -hexachlorocyclohexane, the <sup>3</sup>H content were 11.6 and 5.8 % of control for fu-1 and M<sub>3</sub>A cells, respectively. The results in table 1 also indicate that M<sub>3</sub>A cells in DMEM + 10% FBS generate water-soluble metabolites more rapidly than fu-1 cells (9,584 vs 8,321 cpm). Comparing these results on phosphatidylinositol metabolism with the results on cell proliferation (figures 1a and b) indicates that  $\gamma$ -hexachlorocyclohexane does not affect these two properties identically. Doubling the concentration of  $\gamma$ -hexachlorocyclohexane in the media from 1  $\times$  10<sup>-4</sup> M to 2  $\times$  10<sup>-4</sup> M has little effect on cell proliferation (figures 1a and b) but reduces water-soluble metabolites by 50% or more (table 1).

Because we saw a considerable reduction in the incorporation of myo-[2-<sup>3</sup>H]inositol into water-soluble metabolites (end-product of phosphatidylinositol hydrolysis), we suggest that  $\gamma$ -hexachlorocyclohexane acts, as has been shown previously (Hokin and Brown, 1969; Fisher and Mueller, 1971; Ristow et al., 1973; Hoffman et al., 1980; Vu and Zelenka, 1983; Zelenka and Vu, 1984) to block phosphatidylinositol synthesis. In addition, since we (M. Koohmaraie and J. E. Schollmeyer, unpublished results) and others (Hoffman et al., 1980) could not override the inhibitory effect of  $\gamma$ -hexachlorocyclohexane by the addition of excess inositol, we conclude that  $\gamma$ -hexachlorocyclohexane could not be functioning as a

myo-inositol analogue as proposed by Hoffman et al. (1980). The ability of these cells to continue to proliferate for the first few hours in the presence of  $\gamma$ -hexachlorocyclohexane suggests that there are sufficient stores of inositol lipids to support that proliferation.

These results, when considered collectively with similar results with other cell types (Zelenka, 1980; Ishima et al., 1982), lend substantial support to the hypothesis that cell proliferation and phosphatidylinositol turnover are linked. The precise manner in which these two events are interrelated is not yet understood. However, the fact that various cell types, when presented with mitogenic stimuli, display elevated phosphatidylinositol hydrolysis (Ristow et al., 1973; Hoffman et al., 1974; Habenicht et al., 1981; Sawyer and Cohen, 1981; Berridge et al., 1984) lends support to the hypothesis that the two events are linked. Because of our interest in control of muscle cell proliferation, we are currently investigating whether the mitogenic response of muscle cells depends, in part, on the promotion of phosphatidylinositol turnover, and whether we can stimulate muscle cell proliferation by enhancing phosphatidylinositol turnover by means other than mitogens.

#### Literature Cited

- Aub, D. L. and J. W. Putney, Jr. 1985. Properties of receptor-controlled inositol trisphosphate formation in parotid acinar cells. *Biochem. J.* 225:263.
- Berridge, M. J. 1984. Inositol trisphosphate and diacylglycerol as second messengers. *Biochem. J.* 220:345.
- Berridge, M. J., C. P. Downes and M. R. Hanley. 1982. Lithium amplifies agonist-dependent phosphatidylinositol responses in brain and salivary glands. *Biochem. J.* 206:587.
- Berridge, M. J., J. P. Heslop, R. F. Irvine and K. D.

- Brown, 1984. Inositol trisphosphate formation and calcium mobilization in Swiss 3T3 cells in response to platelet-derived growth factor. *Biochem. J.* 222:195.
- Biden, T. J., M. Prentki, R. F. Irvine, M. J. Berridge and C. B. Wollheim. 1984. Inositol 1,4,5-trisphosphate mobilizes intracellular  $Ca^{++}$  from permeabilized insulin-secreting cells. *Biochem. J.* 223:467.
- Burgess, G. M., P. P. Godfrey, J. S. McKinney, M. J. Berridge, R. F. Irvine and J. W. Putney, Jr. 1984a. The second messenger linking receptor activation to internal Ca release in liver. *Nature* 309:63.
- Burgess, G. M., R. F. Irvine, M. J. Berridge, J. S. McKinney and J. W. Putney, Jr. 1984b. Actions of inositol phosphate on  $Ca^{++}$  pools in guinea pig hepatocytes. *Biochem. J.* 224:741.
- Das, S. and R. P. Rand. 1984. Diacylglycerol causes major structural transitions in phospholipid bilayer membranes. *Biochem. Biophys. Res. Comm.* 124:491.
- Dawson, A. P. and R. F. Irvine. 1984. Inositol (1,4,5) trisphosphate-promoted  $Ca^{++}$  release from microsomal fraction of rat liver. *Biochem. Biophys. Res. Comm.* 120:858.
- Fisher, D. B. and G. C. Mueller. 1971. Gamma-hexachlorocyclohexane inhibits the initiation of lymphocyte growth by phytohemagglutinin. *Biochem. Pharm.* 20:2515.
- Folch, J. 1949. Brain diphosphoinositide a new phosphatide having inositol metadiphosphate as a constituent. *J. Biol. Chem.* 117:505.
- Habenicht, A. J., J. A. Glomset, W. C. King, C. Nist, C. D. Mitchell and R. Ross. 1981. Early changes in phosphatidylinositol and arachidonic acid metabolism in Quiescent Swiss 3T3 cells stimulated to divide by platelet-derived growth factor. *J. Biol. Chem.* 256:12329.
- Hoffmann, R., P. Erzberger, W. Frank and H-J. Ristow. 1980. Increased phosphatidylinositol synthesis in rat embryo fibroblasts after growth stimulation and its inhibition by  $\gamma$ -hexachlorocyclohexane. *Biochem. Biophys. Acta* 618:282.
- Hoffmann, R., H-J. Ristow, H. Pachowsky and W. Frank. 1974. Phospholipid metabolism in embryonic rat fibroblasts following stimulation by a combination of serum protein S1 and S2. *Eur. J. Biochem.* 49:317.
- Hokin, M. R. and D. F. Brown. 1969. Inhibition by  $\gamma$ -hexachlorocyclohexane of acetylcholine-stimulated phosphatidylinositol synthesis in cerebral cortex slices and of phosphatidic acid-inositol transferase in cerebral cortex particulate fractions. *J. Neurochem.* 16:475.
- Irvine, R. F., K. D. Brown and M. J. Berridge. 1984. Specificity of inositol trisphosphate-induced calcium release from permeabilized Swiss-mouse 3T3 cells. *Biochem. J.* 221:269.
- Ishima, Y., Y. Nakagawa and K. Waku. 1982. The high turnover rate of phosphatidylinositol in chicken heart ventricular cells during the earlier stages of development. *Arch. Biochem. Biophys.* 214:180.
- Joseph, S. K., A. P. Thomas, R. J. Williams, R. F. Irvine and J. R. Williamson. 1984a. Myo-inositol 1,4,5-trisphosphate. A second messenger for the release of the hormonal mobilization of intracellular  $Ca^{++}$  in liver. *J. Biol. Chem.* 259:3077.
- Joseph, S. K., R. J. Williams, B. E. Corkey, F. M. Matschinsky and J. R. Williamson. 1984b. The effect of inositol trisphosphate on  $Ca^{++}$  fluxes in insulin-secreting tumor cells. *J. Biol. Chem.* 259:12952.
- Kaufman, S. J., C. M. Parks, J. Bohn and L. E. Faiman. 1980. Transformation is an alternative to normal skeletal muscle development. *Exp. Cell Res.* 125:333.
- Kishimoto, A., Y. Takai, T. Mori, U. Kikkawa and Y. Nishizuka. 1980. Activation of calcium and phospholipid-dependent protein kinase by diacylglycerol, its possible relation to phosphatidylinositol turnover. *J. Biol. Chem.* 255:2273.
- Kolesnick, R. N. and M. C. Gershengorn. 1984.  $Ca^{++}$  ionophores affect phosphoinositide metabolism differently than thyrotropin-releasing hormone in  $GH_3$  pituitary cells. *J. Biol. Chem.* 259:9514.
- Michell, R. H. 1979. Inositol lipids in membrane function. *Trends Biochem. Sci.* 4:128.
- Prentki, M., T. J. Biden, C. D. Janjie, R. F. Irvine, M. J. Berridge and C. B. Wollheim. 1984. Rapid mobilization of  $Ca^{++}$  from rat insulinoma microsomes by inositol-1,4,5-trisphosphate. *Nature* 309:562.
- Rebeci, M. J., R. N. Kolesnick and M. C. Gershengorn. 1983. Thyrotropin-releasing hormone stimulates rapid loss of phosphatidylinositol and its conversion to 1,2 diacylglycerol and phosphatidic acid in rat mammatropic pituitary cells. Association with calcium mobilization and prolactin secretion. *J. Biol. Chem.* 258:227.
- Ristow, H-J., W. Frank and M. Frohlich. 1973. Stimulation of embryonic rat cells by calf serum: metabolism of choline and inositol lipids. *Z. Naturforsch.* 28c:188.
- Ristow, H-J., R. Hoffmann, H. Pachowsky and W. Frank. 1975. Influence of potassium and calcium on the ptdins and phosphatidylcholine metabolism in rat fibroblasts after growth stimulation by calf serum. *Eur. J. Biochem.* 56:413.
- Sawyer, S. T. and S. Cohen. 1981. Enhancement of calcium uptake and phosphatidylinositol turnover by epidermal growth factor in A-431 cells. *Biochem.* 20:6280.
- Schubert, D. and M. Lacorbienic. 1982. Properties of extracellular adhesion-mediating particles in myoblast clone and its adhesion-deficient variant. *J. Cell Biol.* 94:108.
- Streb, H., R. F. Irvine, M. J. Berridge and I. Schulz. 1983. Release of  $Ca^{++}$  from a nonmitochondrial intracellular store in pancreatic acinar cells by inositol 1,4,5-trisphosphate. *Nature* 306:67.
- Vu, N. D., G. Chepko and P. Zelenka. 1983. Decreased turnover of phosphatidylinositol accompanies in vitro differentiation of embryonic chicken lens epithelial cells into lens fibers. *Biochem. Biophys. Acta* 750:105.
- Zelenka, P. S. 1980. Changes in phosphatidylinositol metabolism during differentiation of lens epithelial cells into lens fiber cells in the embryonic chick. *J. Biol. Chem.* 255:1296.
- Zelenka, P. S. and N. D. Vu. 1984. Correlation between phosphatidylinositol degradation and cell division in embryonic chicken lens epithelia. *Dev. Biol.* 105:325.