

Physiology of the Meishan boar

D. D. Lunstra¹, J. J. Ford¹, J. Klindt² and T. H. Wise¹

USDA, ARS, ¹Reproduction Research Unit and ²Nutrition Research Unit,
R. L. Hruska US Meat Animal Research Center, Clay Center, Nebraska 68933-0166, USA

Onset of puberty (sperm production) occurs at a much younger age (56-84 days) in Meishans than in conventional boars (120-180 days). Throughout postnatal development, Meishans exhibit markedly higher (two- to ten-fold) concentrations of serum FSH, LH and androgens compared with conventional boars, and these high hormone concentrations are maintained at maturity. Increased gonadotrophin concentrations occur only in Meishan males, since concentrations in female Meishans and conventional females do not differ. In the Meishan boar, FSH increases steadily during establishment of sperm production and LH increases markedly after tubule diameter stabilizes, while FSH and LH concentrations are low and change little during this period of rapid testicular growth in conventional boars. The proportion of the testis occupied by Leydig cells is two-fold greater and average Leydig cell size is two- to four-fold larger in Meishans than in conventional boars during the onset of spermatogenesis and into maturity. Testis size in Meishans is only half that of conventional boars at maturity, and the number of Sertoli cells is markedly reduced in Meishans. However, Meishan testes exhibit reduced degeneration of germ cells and spermatids during spermatogenesis, and spermatogenic efficiency per Sertoli cell in Meishans is double that in conventional boars. At maturity, Meishan pituitaries have greater mass, larger gonadotrophs, higher content of FSH, and greater expression of genes for FSH and LH subunits than found in conventional boars. The high concentrations of serum gonadotrophins in the mature Meishan exhibit a typical postcastration rise, respond to GnRH stimulation and are suppressed by gonadal steroids, indicating that the hypothalamo-pituitary-gonadal axis functions normally but with much higher set points in Meishan boars. These unusual endocrine and physiological attributes in boars of the highly prolific Meishan breed offer a unique model for further investigation of factors influencing early onset of puberty and efficiency of male reproductive function.

Introduction

There are over 75 breeds of pig in China today; some of these have been selected for over 7000 years (Li and Enfield, 1989). Hierarchical classification of Chinese breeds based on geographic distribution, history, body size, physical appearance and reproductive characteristics identified one group, known as Taihu pigs, which were characterized by high rates of reproduction and average size (Li and Enfield, 1989). Taihu pigs originate from the Lake Taihu region west of Shanghai in southern China. Meishans are typical of this group; they exhibit early sexual maturity and high prolificacy (Cheng, 1983; Wang, 1990). The importation of Meishan pigs into France (1979), the United Kingdom (1987), United States (1989) and other countries allowed research that documented greater prolificacy in Meishans than in conventional pigs of these countries, averaging three to five more pigs per litter (Haley *et al.*, 1995). Female Meishans reach puberty at very young ages (85-118 days of age), at least 3-4 months earlier than pubertal age (200-235 days of age) in conventional pigs (Wang, 1990; Christenson, 1993; Hunter *et al.*, 1993; White *et al.*, 1993). In this review, conventional breeds refers to typical pig breeds of US and European origin. Meishan boars, compared with

conventional breeds of pig, offer a unique model for investigation of factors influencing male reproductive function that relate to high prolificacy, but only limited data on pubertal development and mature reproductive function have been reported for these boars. In addition, no study has characterized relationships between testicular structure and endocrine function in Meishan boars. The purpose of this review is to summarize results on pubertal development, endocrinology, and mature reproductive function in Meishan boars and to compare these with observations in conventional breeds of boar to provide insights concerning the primary mechanisms controlling reproductive function in male pigs.

Testicular Development and Onset of Sperm Production in Meishan Boars

During pubertal development in Meishan boars, body weight (BWT), paired testis weight (PTW) and epididymal weights increase continuously from birth to 16 weeks of age (Fig. 1; Harayama *et al.*, 1991; Lunstra *et al.*, 1993). The testis exhibits a typical rapid increase in mass associated with pubertal development beyond 6 weeks of age (Fig. 1). Elongated spermatids first appear within seminiferous tubules at 6 weeks of age, and homogenization-resistant spermatid nuclei consistently occur in the testes by 8 weeks of age (BWT = 10.3 kg, PTW = 30.9 g), which is slightly earlier than the appearance of elongated spermatids at 60–75 days of age reported for Meishan boars in Japan (Harayama *et al.*, 1991). Percentage of tubules exhibiting elongated spermatids increases steadily and > 80% have elongated spermatids by 12 weeks of age (BWT = 19.2 kg, PTW = 63.3 g). Lumina are first observed in seminiferous tubules at 4 weeks of age; the majority of tubules exhibit lumina by 6 weeks; and spermatozoa are present in the lumina of seminiferous tubules at 8 weeks. Thus, onset of spermatogenesis (spermatozoa formed and released into the lumen of the seminiferous tubules) in the Meishan occurs at 8 weeks of age (Fig. 1). Spermatozoa first appear in the cauda epididymis at 8 weeks of age; first motile spermatozoa (about 20% motile) are present by 10 weeks of age, and appreciable sperm motility (> 70% motile) is attained by 14 weeks and beyond, which is in agreement with results of other workers (Cheng, 1983; Wang, 1990; Harayama and Kato, 1992).

Onset of sperm production (puberty) in Meishan boars occurs much earlier (56–84 days) and at lower testis weight (30–60 g PTW) compared with the 120–180 days of age and 160–240 g PTW at puberty reported for various breeds of conventional boar (FlorCruz and Lapwood, 1978; Allrich *et al.*, 1982, 1983; Colenbrander *et al.*, 1982; Ford, 1983; Young *et al.*, 1986; Lee *et al.*, 1987; Kosco *et al.*, 1987; Trudeau *et al.*, 1992). Data from Meishans extend the range of differences among breeds within a species for body weight and testis weight at given ages during pubertal development (Young *et al.*, 1986). In addition, puberty occurs earlier in males than in females, even within the Meishan, since puberty in Meishan boars occurs 30–60 days earlier than in Meishan females.

Changes in Testicular Structure During Pubertal Development in Meishan Boars

During postnatal testicular development in Meishan boars, the patterns of change in diameter of seminiferous tubules and in volume percentages of the testis occupied by tubules, interstitial tissue and Leydig cells (Fig. 1) are very similar to patterns exhibited by conventional boars but are more compressed in time. The diameter of seminiferous tubules in the Meishan is low at birth (about 70 μm), increases rapidly and linearly through first sperm production (about 200 μm ; 8 weeks), and reaches a plateau by 12 weeks of age and thereafter at 250–290 μm (Fig. 1 and Table 1; Harayama *et al.*, 1991). Tubule diameter in conventional boars is also small at birth (about 60 μm), increases peripubertally and reaches a plateau at approximately 24 weeks of age and beyond (FlorCruz and Lapwood, 1978; van Straaten and Wensing, 1977; Allrich *et al.*, 1983). Maximal tubule diameter tends to be larger in Meishan (250–290 μm) than in conventional boars (190–257 μm ; Table 1; FlorCruz and Lapwood, 1978; van Straaten and Wensing, 1977; Allrich *et al.*, 1983; Lee *et al.*, 1987). Appearance of lumina in seminiferous tubules indicates establishment of the blood–testis barrier; thus this barrier is established at a markedly younger age in Meishan (about 40 days of age) than in conventional boars

factors influencing male
on pubertal development
In addition, no study has
function in Meishan boars.
ment, endocrinology, and
mechanisms controlling

Meishan Boars

paired testis weight (PTW)
age (Fig. 1; Harayama *et al.*,
associated with pubertal
appear within seminiferous
consistently occur in the
earlier than the appearance
in Japan (Harayama *et al.*,
steadily and > 80% have
lumina are first observed
ules exhibit lumina by
es at 8 weeks. Thus, onset
the seminiferous tubules)
the cauda epididymis at
ent by 10 weeks of age,
and beyond, which is in
ma and Kato, 1992).

earlier (56–84 days) and at
age and 160–240 g PTW at
apwood, 1978; Allrich *et al.*
et al., 1987; Kosco *et al.*,
ices among breeds within
development (Young *et al.*,
within the Meishan, since
es.

Meishan Boars

of change in diameter of
tubules, interstitial tissue
ional boars but are more
is low at birth (about 70
at 200 μm ; 8 weeks), and
and Table 1; Harayama *et al.*
(about 60 μm), increases
and beyond (FlorCruz and
nal tubule diameter tends
tm; Table 1; FlorCruz and
et al., 1987). Appearance of
carrier; thus this barrier is
an in conventional boars




Fig. 1. Patterns of change in (a) body weight (\square) and paired testes weight (\circ); (b) seminiferous tubule diameter (\square) and volume (%) seminiferous tubules (\circ); (c) volume (%) interstitial tissue (\square) and volume (%) Leydig cells (\circ) in Meishan boars from birth through puberty. An average of seven Meishan boars per age point were castrated and evaluated. (\bullet) A comparison of changes in Leydig cell size during postnatal development in Meishan (\square) and conventional boars (\circ). Allrich *et al.*, 1983; \circ : van Straaten and Wensing, 1978) is shown.

Table 1. Differences between mature Meishan and conventional boars in size of testes, sperm production and various aspects of testicular structure^a

	Meishan (n=5)	Composite ^b (n=5)
Age (days)	225 ± 3	346 ± 2
Body weight (kg)	78 ± 2	133 ± 6
Paired testes weight (g)	269 ± 19	541 ± 35
Volume (%) seminiferous tubules	48.6 ± 1.7	68.7 ± 1.6
Volume (%) interstitial tissue	51.4 ± 1.9	31.3 ± 0.8
Volume (%) Leydig cells	18.5 ± 0.6	7.1 ± 0.2
Volume (%) vascular	10.0 ± 0.9	6.5 ± 0.8
Seminiferous tubules, diameter (µm)	287 ± 15	257 ± 8
Daily sperm production/g ⁻¹ testis ^c	24.47 ± 1.26	24.77 ± 1.43
Total daily sperm production ^d	6.04 ± 0.52	12.46 ± 1.43
Total no. Sertoli cells ^d	3.87 ± 0.59	14.45 ± 1.33
DSP per Sertoli cell	1.56 ± 0.32	0.86 ± 0.20
Total no. Leydig cells ^d	15.27 ± 0.66	34.79 ± 1.76
Leydig cell size (µm ³)	2855 ± 124	960 ± 49

^aCalculated from data of Okwun *et al.* (1996a,b). All data shown are presented as means ± SEM. Volume (%): volume percentage of testicular parenchyma occupied by respective structure or cell type.

^bComposite boars: crossbred boars (1/4 Chester White, 1/4 Landrace, 1/4 Large White, 1/4 Yorkshire). Each mean ± SEM for Meishan differs ($P < 0.10$ to $P < 0.01$) from that for Composite boars, except ($P > 0.20$) for average tubule diameter and daily sperm production g⁻¹ testis.

^cDaily sperm production (DSP) × 10⁶ per gram of testicular parenchyma.

^dTotal daily sperm production (TDSP) × 10⁹ per paired testes. Total number of Sertoli cells × 10⁶ per paired testes. Total number of Leydig cells × 10⁶ per paired testes.

(90–120 days; FlorCruz and Lapwood, 1978; Allrich *et al.*, 1983; Colenbrander *et al.*, 1982). Rapid proliferation of Sertoli cells occurs primarily during the first month after birth in conventional boars (Tran *et al.*, 1981; Kosco *et al.*, 1987), and establishment of the blood–testis barrier indicates cessation of proliferation of Sertoli cells thereby defining the total population of Sertoli cells and maximum mature testis size (Russell *et al.*, 1990; Chubb, 1992). The markedly earlier onset of puberty in the Meishan requires curtailment of Sertoli cell proliferation sooner in the early postnatal period. Indeed, at sexual maturity, Meishan boars have substantially smaller paired testis weight and fewer Sertoli cells than do conventional boars (Table 1).

In Meishan (Fig. 1) and conventional boars, the percentage of the testis occupied by total interstitial tissue and by Leydig cells exhibits a major early postnatal peak at 1–4 weeks of age followed by a rapid decline through onset of puberty and maintenance at reduced values after puberty (van Straaten and Wensing, 1977, 1978; FlorCruz and Lapwood, 1978; Colenbrander *et al.*, 1982; Allrich *et al.*, 1983). While similar in pattern of change, the absolute values for volume percentage of the testis occupied by interstitial tissue (35–50%) and by Leydig cells (15–25%) in Meishan boars beyond 10 weeks of age and into maturity remains substantially greater than the values reported for conventional boars beyond 20 weeks of age (15–30% interstitial, 7–10% Leydig, respectively). Thus, during the latter part of pubertal development and into maturity (Table 1), Meishan boars consistently exhibit a 1.5- to 2.5-fold higher proportion of the testes occupied by interstitium and Leydig cells than do conventional boars.

Average volume per Leydig cell, in both Meishan (Fig. 1 and Table 1) and conventional boars, exhibits a major early postnatal peak at 1–4 weeks of age, followed by a reduction in size, a transient increase during onset of spermatogenesis, and a slow decline in size that is maintained into maturity (van Straaten and Wensing, 1977, 1978; FlorCruz and Lapwood, 1978; Colenbrander *et al.*, 1982; Allrich *et al.*, 1983). Despite the similarity in pattern, absolute values for Leydig cell size in the

Meishan
larger at
Straaten
with the
and am
unpubli
steroid-s
boars de
increas
also exhib
and both
12 and 14
in the Me
prepuber
relatively

In Meisha
concentrat
2). One str
LH, testost
developme
(Table 2, a
and Raesc
1984; Kosco

A major
androgens
Although t
serum test
preceding
also occur
1982; Kosco
2; note sm
concentrat
Meishan (8
Meishans,
period of in
maintained
but the incr
after the ea
puberty and
coinciding
(Colenbrand
1996), and s
changes (A
Despite the
serum LH d
Meishan and

The high
postpubertal
somewhat be

boars in size of
structure^a

Composite^b
(n=5)

346 ± 2
133 ± 6
541 ± 35
68.7 ± 1.6
31.3 ± 0.8
7.1 ± 0.2
6.5 ± 0.8
257 ± 8
4.77 ± 1.43
2.46 ± 1.43
4.45 ± 1.33
0.86 ± 0.20
1.79 ± 1.76
960 ± 49

± SEM. Volume (%):

1/4 Yorkshire). Each
experiment (P > 0.20) for

cells × 10⁶ per paired

nder *et al.*, 1982). Rapid
in conventional boars
carrier indicates cessation
toli cells and maximum
onset of puberty in the
early postnatal period.
testis weight and fewer

testis occupied by total
ak at 1–4 weeks of age
at reduced values after
78; Colenbrander *et al.*,
lute values for volume
eydig cells (15–25%) in
ntially greater than the
terstitial, 7–10% Leydig,
into maturity (Table 1),
the testes occupied by

and conventional boars,
ction in size, a transient
maintained into maturity
lenbrander *et al.*, 1982;
Leydig cell size in the

Meishan are markedly larger than in other breeds (Fig. 1 and Table 1), and average two- to four-fold larger at every age examined than reported values for Leydig cell size in conventional boars (van Straaten and Wensing, 1978; Peyrat *et al.*, 1981; Allrich *et al.*, 1983; Lunstra *et al.*, 1986). Concomitant with the increase in cell size, increases in average volume of the nucleus, number of mitochondria and amount of endoplasmic reticulum occur within Leydig cells of Meishan testis (Lunstra, unpublished data), and increases in these intracellular components generally reflect increased steroid-synthesizing capacity (Zirkin *et al.*, 1980; Lunstra *et al.*, 1986). Leydig cell size in conventional boars declines and remains small (smaller than at birth) between 6 and 16 weeks of age before increasing transiently coincident with increasing spermatogenesis (18–26 weeks of age). Meishans also exhibit a less noticeable increase coinciding with increasing spermatogenesis (12 weeks of age) and both the percentage interstitial tissue and Leydig cells exhibit small but consistent increases at 12 and 14 weeks of age (Fig. 1). Thus, the early postnatal and pubertal peaks occur so close together in the Meishan, and overall Leydig cell size is so much larger, that a pronounced regression in prepubertal Leydig cell volume does not occur, and the rebound in Leydig cell size at puberty is relatively small (Fig. 1).

Endocrinology During Pubertal Development in Meishan Boars

In Meishan boars, serum FSH, LH, testosterone and dehydroepiandrosterone sulfate (DHEAS) concentrations all change markedly and exhibit distinct patterns during pubertal development (Fig. 2). One striking characteristic in Meishans is that serum concentrations of all four hormones (FSH, LH, testosterone and DHEAS) are each substantially higher (two- to ten-fold) throughout postnatal development, both before and after puberty, than in conventional boars, as assayed in our laboratory (Table 2, and unpublished data) or reported by others for various breeds of conventional boar (Tan and Raeside, 1980; Allrich *et al.*, 1982, 1983; Colenbrander *et al.*, 1982; Ford, 1983; Schinckel *et al.*, 1984; Kosco *et al.*, 1987; Trudeau *et al.*, 1992).

A major peak in serum concentrations of FSH (1–2 weeks of age), LH (3–4 weeks of age) and androgens (3–4 weeks of age) occurs early in the postnatal period in Meishan boars (Fig. 2). Although the early postnatal peak in testosterone appears to be quite subtle (Fig. 2), the increase in serum testosterone concentrations at 3–4 weeks of age is significantly higher than that during the preceding (2 week) and following (5 week) time frames. Early postnatal peaks in serum hormones also occur at comparable ages or even slightly earlier in conventional boars (Colenbrander *et al.*, 1982; Kosco *et al.*, 1987; Trudeau *et al.*, 1992; Ford *et al.*, 1997), although absolute values are lower (Fig. 2; note smaller scale of y-axis for conventional boars). After the early postnatal peak, serum concentrations of gonadotrophins and androgens decline at least until the onset of puberty in both Meishan (8 weeks) and conventional boars (16 weeks). However, with onset of spermatogenesis in Meishans, serum FSH, testosterone and DHEAS all increase steadily and markedly during the period of increasing testis mass (Figs 1 and 2), and very high concentrations of these hormones are maintained in adulthood (Table 2). Serum concentrations of LH in Meishans also increase markedly, but the increase is delayed until maximum tubule diameter is achieved (Figs 1 and 2). In contrast, after the early postnatal peak in conventional boars, serum FSH remains relatively low through puberty and on through sexual maturity, although a small increase in FSH immediately preceding or coinciding with the onset of sperm production has been reported in conventional boars (Colenbrander *et al.*, 1982; Schinckel *et al.*, 1984; Kosco *et al.*, 1987; Trudeau *et al.*, 1992; Mariscal *et al.*, 1996), and serum concentrations of LH remain low into adulthood with no significant peripubertal changes (Allrich *et al.*, 1982; Colenbrander *et al.*, 1982; Schinckel *et al.*, 1984; Kosco *et al.*, 1987). Despite the large differences among breed in serum concentrations of hormones, it is surprising that serum LH declines before and changes relatively little during the onset of spermatogenesis in both Meishan and conventional boars (Fig. 2).

The highest peripubertal concentrations of FSH, LH, testosterone and DHEAS all occur postpubertally at 15–16 weeks of age in Meishan boars. Although serum LH concentrations decline somewhat between 16 weeks of age and maturity in Meishan boars (compare Fig. 2 and Table 2), the

Table 2. Differences between mature Meishan and conventional boars in body weight, testes size and circulating plasma hormone concentrations

Characteristic	Meishan (Borg <i>et al.</i> , 1993)	Duroc ^a (Borg <i>et al.</i> , 1993)	Meishan (Borg <i>et al.</i> , 1993)	Duroc ^a (Borg <i>et al.</i> , 1993)	Meishan Composite ^b (Wise <i>et al.</i> , 1996)	Meishan (Ford <i>et al.</i> , 1997)	Composite ^b (Ford <i>et al.</i> , 1997)
Number of boars	9	8	9	8	8	22	22
Age (days)	533 ± 5	453 ± 11	706 ± 5	627 ± 11	682 ± 55	509 ± 44	440 ± 41
Body weight (kg) ^c	140 ± 5	184 ± 4	164 ± 7	197 ± 6	147 ± 4	115 ± 5	147 ± 6
Paired testes weight (g) ^c	417 ± 37 ^d	701 ± 64 ^d	408 ± 31 ^d	969 ± 31 ^d	394 ± 53	325 ± 24	547 ± 25
Plasma FSH (ng ml ⁻¹) ^c	634 ± 79	65 ± 11	656 ± 73	91 ± 11	554 ± 50	704 ± 48	136 ± 48
Plasma LH (ng ml ⁻¹) ^c	nd	nd	nd	nd	1.3 ± 0.1	nd	nd
Plasma testosterone (ng ml ⁻¹) ^c	15.1 ± 5.0	6.9 ± 1.7	20.8 ± 6.9	7.5 ± 1.7	14.5 ± 0.6	16.8 ± 2.6	3.8 ± 1.3
Plasma inhibin (fmol ml ⁻¹) ^c	97 ± 9	147 ± 18	117 ± 17	269 ± 16	131 ± 13	nd	nd

^aPurebred Duroc boars.

^bComposite boars: crossbred boars (1/4 Chester White, 1/4 Landrace, 1/4 Large White, 1/4 Yorkshire).

^cWithin each study, mean for Meishan boars differs significantly ($P < 0.05$ to $P < 0.01$) from that for conventional boars, except ($P > 0.10$) for body weight (kg) and plasma concentrations of inhibin reported by Wise *et al.* (1996).

^dPaired testis volume (cm³) calculated from caliper measurements (equivalent to weight, assuming 1.0 cm³ = 1.0 g).

nd: not determined. All data shown are presented as means ± SEM. Data from Borg *et al.* (1993), Wise *et al.* (1996); and Ford *et al.* (1997).




Fig. 2. Plasma concentrations of (a) DHEAS, (b) testosterone and (c) LH in Meishan (filled bars) and Duroc (open bars) boars.

higher sex steroid concentrations at sexual maturity (age), but similar sex steroid concentrations at sexual maturity.




Fig. 2. Patterns of change in (a,b) FSH, (c,d) LH, (e,f) testosterone and (g,h) dehydroepiandrosterone sulfate (DHEAS) concentrations in (a,c,e,g) Meishan and (b,d,f,h) conventional boars from birth through puberty. (b) Data from (□) Kosco *et al.*, 1987; (◇) Colenbrander *et al.*, 1982; and (●) Schinckel *et al.*, 1984; (d) data from (□) Kosco *et al.*, 1987; (◇) Colenbrander *et al.*, 1982 and (○) Allrich *et al.*, 1982; (f) data combined from Tan and Raeside (1980), Allrich *et al.* (1982), Ford (1983) and Kosco *et al.* (1987); (h) data from Tan and Raeside (1980).

higher serum concentrations of FSH, LH, and testosterone present at 16 weeks of age are maintained at sexual maturity (Table 2). One possible explanation is that the sensitivity of the hypothalamus to steroid feedback in the Meishan boar may be greatly reduced during this time frame (14–16 weeks of age), but the reduction must be temporary, since feedback sensitivity seems to be fully functional at sexual maturity (see discussion on sexual maturity below). Regardless of the mechanism, Meishan

boars continue to exhibit greater concentrations of serum FSH, LH, and androgens than do conventional boars at sexual maturity (Table 2).

In Meishan and conventional boars, the major early postnatal peaks in serum concentrations of FSH, LH and androgens coincide well with the timing of early postnatal peaks observed in volume percentages of the testis occupied by interstitial tissue and Leydig cells, and with the marked early peak in Leydig cell size (compare Fig. 1 and Fig. 2). Although Meishans achieve puberty at a much younger age than do conventional boars, the coincidence between structural and hormonal peaks indicate that there is a strong interdependence and cause-and-effect relationship amongst these characteristics, even in these two very different types of pig. Presumably, the high early postnatal concentrations of gonadotrophins stimulate the marked increase in Leydig cell size, and the increased steroidogenic machinery in the enlarged Leydig cells synthesize and secrete enough steroids to produce the early postnatal peak in serum androgens.

In conventional boars, the general decreases in serum LH and FSH through puberty and into sexual maturity, with perhaps a small transient increase in FSH preceding or coinciding with the onset of sperm production, are not due to decreases in pituitary gonadotrophin content. Indeed, hypothalamic content of GnRH and pituitary content of FSH and LH increases progressively from birth through puberty in conventional male pigs (McNamara *et al.*, 1985; Lee *et al.*, 1987). It is more likely that the rapid increase in testis size, increasing total Leydig cell mass, and increasing numbers of receptors for both LH and FSH (presumably occurring in Meishans, as has been demonstrated in conventional boars) collectively contribute to rising serum concentrations of androgens during this time frame, i.e., an increase in testicular responsiveness. The increasing serum androgens presumably allow steroid feedback to maintain low serum concentrations of LH during this peripubertal period. Assuming that this occurs, the feedback suppression on gonadotrophins appears to be expressed differently in the Meishan, since LH is suppressed (at least until 13 weeks of age) while FSH increases steadily and markedly at 8 weeks of age and beyond (Fig. 2).

General Reproductive Characteristics in Mature Meishan Boars

In general, mature Meishan boars exhibit lower body weight and much smaller testicular size than do conventional breeds of boar at comparable ages (Table 2). The number of spermatozoa per ejaculate from Meishan boars is approximately half that of conventional boars (Borg *et al.*, 1993; Gerfen *et al.*, 1994; Huang and Johnson, 1996), but percentage progressively motile spermatozoa, percentage spermatozoa with normal morphology, and other semen quality characteristics are similar, despite major breed differences in mature testis size. In mammals, the number of Sertoli cells determines mature testicular size (Russell *et al.*, 1990; Chubb, 1992), and findings in boars concur with this. Testes of mature Meishans that are approximately half the size of those of conventional boars (Table 1) have substantially fewer Sertoli cells (Okwun *et al.*, 1996a, 1996b). Of significance, daily sperm production per gram of testis (DSP g⁻¹) does not differ between Meishan and conventional boars (Table 1); therefore, major differences in total DSP primarily reflect breed differences in paired testis weight. However, the percentage of the testis occupied by seminiferous tubules is 1.4-fold lower in Meishan than in conventional boars (Table 1). Meishan boars produce approximately twice as many spermatozoa per Sertoli cell (Table 1) and almost twice as many elongated spermatids per type A spermatogonium compared with conventional boars to accommodate this change in testis composition (Okwun *et al.*, 1996a, 1996b). The yield of spermatids per type A spermatogonium is a reliable measure of efficiency of spermatogenesis (Berndtson and Igboeli, 1989). Thus, mature Meishan boars produce spermatozoa much more efficiently (per gram of seminiferous tubule, per Sertoli cell, and per spermatogonium) than do conventional boars and appear to achieve this via reduced degeneration of germ cells and spermatids during the spermatogenic process. Although the mechanism that allows Meishan boars to achieve this greater efficiency is unknown, it probably relates to constant exposure of the Sertoli cells to greater concentrations of FSH and androgens.

Sensitivity to GnRH Stimulation in Mature Meishan and Conventional Boars

When mature Meishan and conventional boars (Composite = 1/4 Chester White, 1/4 Landrace, 1/4 Large White, 1/4 Yorkshire) of approximately the same body weight (BWT) are challenged with a high (1500 ng kg⁻¹ BWT) or a low (150 ng kg⁻¹ BWT) dose of GnRH, both breeds respond with significant increases in serum FSH, LH and testosterone concentrations (Wise *et al.*, 1996), and greater increases occur in response to the high dose of GnRH in both types of boar. Despite major pretreatment differences in circulating hormone concentrations, the FSH response to GnRH is comparable (i.e., approximately 125% at low dose and 160% at high dose above pretreatment values) in Meishan and Composite boars. In Meishan boars, the substantially increased serum concentrations of FSH and LH take longer to return to pretreatment values. Although paired testis size is significantly lower in Meishan boars, serum inhibin concentrations differ minimally at pretreatment and changed little during GnRH challenge (Wise *et al.*, 1996). Accepting the limitations of the radioimmunoassay for the α -subunit of inhibin, serum inhibin does not appear to be a primary determinant of high circulating concentrations of FSH in Meishan boars. Thus, under normal physiological conditions, pituitaries of Meishan boars are not functioning under maximal GnRH stimulation and are not secreting maximal amounts of gonadotrophins. High gonadotrophin concentrations in Meishan boars apparently are under control of a functional and responsive GnRH system.

Interestingly, differences in gonadotrophin secretion during the oestrous cycle are not apparent between females of Meishan and European breeds (Hunter *et al.*, 1996). In contrast to the situation in males, ovariectomized European gilts release greater concentrations of FSH and LH in response to GnRH stimulation than do Meishan gilts and are more responsive in their reduction of FSH concentrations after treatment with pig follicular fluid (inhibin effect; Tilton *et al.*, 1994). A mechanism to explain these sex differences in Meishans is not obvious, but in mice that possess a transgene for human FSH β , expression is greater in males, but not in females, relative to normal mice (Kumar and Low, 1993).

Negative Feedback Regulation of FSH in Mature Meishan and Conventional Boars

When mature Meishan and Composite boars that differed 4- to 9-fold in circulating FSH concentrations were castrated, FSH exhibited a postcastration rise, and the rate of this increase (0–48 h) in both breeds did not differ (Wise *et al.*, 1996). In addition, differences in postcastration FSH concentrations were maintained between breeds through at least 16 days. The pattern of decline in serum testosterone did not differ between breeds after castration, although Meishan boars had greater serum testosterone before castration. These results demonstrate that gonadal hormones exert negative feedback on gonadotrophin concentrations in the mature intact Meishan boar, apparently in a manner similar to that observed in intact conventional boars.

Injection of castrate Meishan and Composite boars with the same dosages of gonadal steroids (testosterone + oestradiol, i.m. on kg⁻¹ BWT basis) was equally effective, in both breeds, at reducing the increased concentrations of serum FSH present in castrates (Wise *et al.*, 1996). The rate of decline in FSH after injection did not differ among the breeds, and the relative degree of suppression by gonadal steroids appeared to be similar. The relative difference in serum FSH before steroid injection continued after steroid treatment. Furthermore, serum FSH began to escape exogenous steroid suppression earlier in Meishan than in Composite castrates. Thus, feedback sensitivity to testicular steroids in these two divergent breeds of boar seems to be fully functional and coordinate within their relative concentrations of circulating hormones. Collectively, these data indicate that the high concentrations of gonadotrophins and androgens in Meishans occur under the control of a hypothalamo-pituitary-gonadal axis that functions responsively but with much higher set points in the Meishan boar.

Meishan Pituitary Mass and Biological Activity of FSH

Pituitary size is greater in mature Meishan than in mature Composite boars, and both content and concentration of FSH are substantially higher in anterior pituitaries of Meishan boars (Zanella *et al.*, 1996). The greater expression of the genes for α -subunit and for FSH β and LH β subunits accounts, in part, the observed differences in gonadotrophin concentrations within pituitaries of Meishan boars. In addition, nucleated pituitary cells that are positive immunocytochemically for FSH or LH (gonadotrophs) have larger cell volumes in Meishan than in Composite boars (Macdonald *et al.*, 1995).

Furthermore, in an *in vitro* bioassay for FSH activity that utilized a cell line possessing recombinant FSH receptors coupled with a cAMP-responsive system, the activity of pituitary extracts from Meishan and Composite boars is highly correlated with FSH content determined by radioimmunoassay ($r = 0.92$; Macdonald *et al.*, 1995). In association with this difference in pituitary FSH concentration, gene expression for activin, a stimulator of FSH secretion, is higher in pituitaries of Meishan than in Composite boars (Li *et al.*, 1997). In total, the difference in FSH concentrations between Meishan and Composite boars may be due more to differences in activin regulated synthesis rather than to differences in GnRH regulation of secretion or differences in sensitivity to inhibitors of secretion.

Negative Relationship Between FSH and Mature Testis Size

When the variation in mature testis size and serum FSH concentration was examined, a negative correlation became apparent in mature Meishan boars as well as in mature Composite and Meishan \times Composite crossbred boars ($r = -0.74$, Fig. 3; Lunstra *et al.*, 1996). Owing to the positive correlation between mature testis weight and total daily sperm production (TDSP), there is a negative correlation between FSH and TDSP in these boars. Thus, a three times greater serum FSH concentration was associated with 32% smaller mature testis weight and 33% lower TDSP. In Meishan \times Composite boars destined to exhibit major differences in serum FSH concentration at maturity, we detected divergence in serum FSH concentrations at 2–8 weeks of age, and this divergence remained during pubertal development (4–6 months of age) and at 1 year of age (Ford *et al.*, 1997). In addition, we have found that serum FSH concentration has a moderate heritability ($h^2 = 0.42$) in Meishan \times Composite boars (Rohrer *et al.*, 1997), comparable in magnitude to the heritability of testis size in conventional boars (Young *et al.*, 1986).

The reason for this negative relationship between mature testis size and serum FSH concentration in boars is not clear. In contrast to our findings, FSH secretion (Schinckel *et al.*, 1984; Mariscal *et al.*, 1996) was not influenced by generations of selection for increased testis size in conventional boars (Huang and Johnson, 1996). This lack of an effect on FSH secretion may reflect selection that was not based on testis size at maturity; instead, boars were selected for larger testis size at 150 days of age which would select primarily for earlier pubertal development and secondarily for mature testis size. In contrast to our findings, most other evidence supports a concept that high serum FSH, especially during the early postnatal period, would contribute to greater proliferation of Sertoli cells and result in a larger mature testis size. Treatment of rodents with FSH during early postnatal life causes testicular hypertrophy and increases in numbers of Sertoli cells that persist into adulthood (Meachem *et al.*, 1996). In addition, treatment of conventional boars with FSH from 8 to 40 days of age caused no testicular hypertrophy by 100 days of age, but appeared to delay puberty, to increase the length of seminiferous tubules and the total number of Sertoli cells, and, theoretically, would have increased mature testis size (Swanlund *et al.*, 1995). The latter study contradicts the association between testis size and high serum FSH in Meishan boars (i.e., earlier puberty and reduced mature testis size) and in conventional breeds of boar (i.e., reduced mature testis size). Moreover, other researchers have begun to question the necessity of FSH for regulation of spermatogenesis (Zirkin *et al.*, 1994). Thus, further studies are needed to clarify relationships between extremes in FSH concentrations and extremes in age at puberty and mature testis size.

FSH

ars, and both content and
ishan boars (Zanella *et al.*,
LH β subunits accounts, in
in pituitaries of Meishan
chemically for FSH or LH
boars (Macdonald *et al.*,

and a cell line possessing
the activity of pituitary
H content determined by
his difference in pituitary
on, is higher in pituitaries
nce in FSH concentrations
nces in activin regulated
differences in sensitivity to

Testis Size

was examined, a negative
mature Composite and
6). Owing to the positive
action (TDSP), there is a
times greater serum FSH
and 33% lower TDSP. In
am FSH concentration at
weeks of age, and this
and at 1 year of age (Ford *et al.*
s a moderate heritability
ble in magnitude to the

s size and serum FSH
ion (Schinckel *et al.*, 1984;
r increased testis size in
FSH secretion may reflect
selected for larger testis
pubertal development and
her evidence supports a
iod, would contribute to
reatment of rodents with
ses in numbers of Sertoli
ent of conventional boars
days of age, but appeared
al number of Sertoli cells,
l., 1995). The latter study
eishan boars (i.e., earlier
oar (i.e., reduced mature
y of FSH for regulation of
to clarify relationships
and mature testis size.




Fig. 3. Relationship between testis size and serum FSH concentration in mature (◆) Meishan, (□) Composite and (○) Meishan × Composite boars. All boars were > 1 year of age. Composite boars: 1/4 Chester White, 1/4 Landrace, 1/4 Large White, 1/4 Yorkshire.

Of particular relevance may be that serum TSH concentration is greater and expression of the β -subunit gene for thyroid-stimulating hormone (TSH- β) in pituitaries is higher in mature Meishan than in conventional boars (Li *et al.*, 1996). Sertoli cells have receptors for TSH, and TSH has potency equivalent to FSH for stimulation of androgen binding protein secretion by Sertoli cells (Hutson and Stocco, 1981). Increased serum TSH concentrations (Ford *et al.*, unpublished data) in combination with high FSH concentrations are associated with reduced testis size exhibited by mature boars with high FSH. Early prepubertal treatment of male rodents with propylthiouracil, a compound that blocks T3 and T4 production (hypothyroidism), causes TSH and FSH to remain at low concentrations into adulthood and results in increased numbers of Sertoli cells, increased mature testis size, and increased sperm production (Hess *et al.*, 1993; Simorangkir *et al.*, 1995). The implication from these propylthiouracil studies is that the opposite effect (high concentrations of TSH and FSH) prepubertally would cause the opposite of Sertoli cells and smaller adult testis size. Contrary to prediction, mature testis size of conventional boars was not increased by propylthiouracil treatment from 1 to 9 weeks of age (Tarn *et al.*, 1996), but this treatment failed to decrease serum FSH. At present the prepubertal manipulation of FSH and TSH in conventional boars has not duplicated effects on testicular development comparable to those observed in Meishan boars.

Conclusion

Major endocrine differences and unique differences in the structural and functional efficiency of the testis are exhibited in Meishan boars during pubertal development and at maturity in comparison with boars of conventional breeds. Many fundamental questions remain concerning the factors that control these aspects of reproduction in boars. Why is high FSH secretion throughout development associated with increased spermatogenic efficiency but an unexpected reduction in testis size, while high LH secretion is associated with predictable increases in Leydig cell size and testicular steroidogenesis? These unusual endocrine and physiological attributes in boars of the highly prolific Meishan breed offer a unique model for further investigation of factors influencing male reproductive function. In particular, can the greater efficiency of spermatogenesis in Meishan boars be introgressed into boars of conventional breeds? Hopefully, future results using this model will provide new insights and contributions to the expansion of our understanding of the physiological basis for the reproductive superiority of hyperprolific pigs.

The authors thank A. Kruger, P. Sherman and S. Hassler for technical assistance, personnel of USMARC Swine Operations for care of animals, and D. Griess for secretarial assistance. Mention of trade names is necessary to report factually on available data; however, the USDA neither guarantees nor warrants the standard of the product, and the use of the same by USDA implies no approval of the product to the exclusion of others that may also be suitable.

References

- Allrich RD, Christenson RK, Ford JJ and Zimmerman DR (1982) Pubertal development of the boar: testosterone, estradiol-17 β , cortisol and LH concentrations before and after castration at various ages *Journal of Animal Science* **55** 1139-1146
- Allrich RD, Christenson RK, Ford JJ and Zimmerman DR (1983) Pubertal development of the boar: age-related changes in testicular morphology and *in vitro* production of testosterone and estradiol-17 β *Biology of Reproduction* **28** 902-909
- Berndtson WE and Igboeli G (1989) Numbers of Sertoli cells, quantitative rates of sperm production, and the efficiency of spermatogenesis in relation to the daily sperm output and seminal quality of young beef bulls *American Journal of Veterinary Research* **50** 1193-1197
- Borg KE, Lunstra DD and Christenson RK (1993) Semen characteristics, testicular size, and reproductive hormone concentrations in mature Duroc, Meishan, Fengjing, and Minzhu boars *Biology of Reproduction* **49** 515-521
- Cheng PL (1983) A highly prolific pig breed of China - the Taihu breed (Parts I and II) *Pig News and Information* **4** 407-425
- Christenson RK (1993) Ovulation rate and embryonic survival in Chinese Meishan and white crossbred pigs *Journal of Animal Science* **71** 3060-3066
- Chubb C (1992) Genes regulating testis size *Biology of Reproduction* **47** 29-36
- Colenbrander B, Frankenhuus M and Wensing CJG (1982) Male sexual development In *Control of Pig Reproduction* pp 3-24; Eds D Cole and G Foxcroft. Butterworth Scientific, London
- FlorCruz SV and Lapwood KR (1978) A longitudinal study of pubertal development in boars *International Journal of Andrology* **1** 317-330
- Ford JJ (1983) Serum estrogen concentrations during postnatal development in male pigs *Proceedings of the Society for Experimental Biology and Medicine* **174** 160-164
- Ford JJ, Wise TH and Lunstra DD (1997) Negative relationship between blood concentrations of follicle stimulating hormone (FSH) and testicular size in mature boars *Journal of Animal Science* **75** 790-795
- Gerfen RW, White BR, Cotta MA and Wheeler MB (1994) Comparison of semen characteristics of Fengjing, Meishan and Yorkshire boars *Theriogenology* **41** 461-469
- Haley CS, Lee CJ and Ritchie M (1995) Comparative reproductive performance in Meishan and Large White pigs and their crosses *Animal Sciences* **60** 259-267
- Harayama H and Kato S (1992) Changes in motility and morphology of spermatozoa during their transit through the epididymis in Meishan boars at various ages *Animal Science Technology* **63** 462-467
- Harayama H, Nanjo I, Kanda S and Kato S (1991) Testicular development in Chinese Meishan boars *Theriogenology* **36** 637-643
- Hess RA, Cooke PS, Bunick D and Kirby JD (1993) Adult testicular enlargement induced by neonatal hypothyroidism is accompanied by increased Sertoli and germ cell numbers *Endocrinology* **132** 2607-2613
- Huang Y-T and Johnson RK (1996) Effect of selection for size of testes in boars on semen and testis traits *Journal of Animal Science* **74** 750-760
- Hunter MG, Biggs C, Foxcroft GR, McNeilly AS and Tilten JE (1993) Comparisons of endocrinology and behavioural events during the periovulatory period in Meishan and Large-White hybrid gilts *Journal of Reproduction and Fertility* **97** 475-480
- Hunter MG, Picton HM, Biggs C, Mann GE, McNeilly AS and Foxcroft GR (1996) Periovulatory endocrinology in high ovulating Meishan sows *Journal of Endocrinology* **150** 141-147
- Hutson JC and Stocco DM (1981) Regulation of Sertoli cell function by thyrotropin *Biology of Reproduction* **25** 303-306
- Kosco MS, Bolt DJ, Wheaton JE, Loseth KJ and Crabo BG (1987) Endocrine responses in relation to compensatory testicular growth after neonatal hemicastration in boars *Biology of Reproduction* **36** 1177-1185
- Kumar TR and Low MJ (1993) Gonadal steroid hormone regulation of human and mouse follicle stimulating hormone β -subunit gene expression *in vivo*. *Molecular Endocrinology* **7** 898-906
- Lee K-H, Diekman MA, Moss GE and Allrich RD (1987) Pituitary gonadotropins, hypothalamic gonadotropin-releasing hormone, and testicular traits of boars exposed to natural or supplemental lighting during pubertal development *Biology of Reproduction* **36** 1164-1169
- Li MD and Enfield FD (1989) A characterization of Chinese breeds of swine using cluster analysis *Journal of Animal Breeding Genetics* **106** 379-388
- Li MD, Matteri RL, Macdonald GJ, Wise TH and Ford JJ (1996) Overexpression of β -subunit of thyroid-stimulating hormone in Meishan swine identified by differential display *Journal of Animal Science* **74** 2104-2111
- Li MD, Macdonald GJ and Ford JJ (1997) Breed differences in expression of inhibin/activin subunits in porcine anterior pituitary glands *Endocrinology* **138** 712-718
- Lunstra DD, Ford JJ, Christenson RK and Allrich RD (1986) Changes in Leydig cell ultrastructure and function during pubertal development in the boar *Biology of Reproduction* **34** 145-158
- Lunstra DD, Borg KE and Klindt J (1993) Changes in porcine testicular structure during development in Meishan boars *Biology of Reproduction* **48** (Supplement 1) 126 (Abstract)
- Lunstra DD, Ford JJ and Wise TH (1996) Selection for extremes in serum FSH concentrations results in reduced testis size and fertility in Meishan and White composite boars. In *Advances in Swine in Biomedical Research* Vol. 2 pp. 523-531 Eds ME Tumbleson and LB Schook. Plenum Press, New York

personnel of USMARC Swine trade names is necessary to arrange the standard of the exclusion of others that may

D and Kirby JD (1993) Adult induced by neonatal hypothyroidism increased Sertoli and germ cell numbers in the testis *Journal of Animal Science* 77 2607-2613

D and Kirby JD (1996) Effect of selection for size of testis on reproductive traits *Journal of Animal Science* 82 1155-1161

GR, McNeilly AS and Tilton JE (1994) Gonadotropin-releasing hormone and luteinizing hormone-releasing hormone during the oestrous cycle in Meishan and European pig *Journal of Reproduction and Fertility* 66 115-121

GR, Mann GE, McNeilly AS and Tilton JE (1995) Testicular endocrinology in high and low prolific Meishan boars *Journal of Endocrinology* 150 105-111

GR (1981) Regulation of Sertoli cell function in the testis *Journal of Reproduction and Fertility* 25 303-306
E, Loseth KJ and Crabo BG (1987) Effect of castration on compensatory testicular growth in boars *Journal of Animal Science* 65 1155-1161

GR (1993) Gonadal steroid hormone and mouse follicle stimulating hormone expression *in vivo*. *Molecular Reproduction and Development* 10 1-10

GR and Allrich RD (1987) Pituitary gonadotropin-releasing hormone in the testis of boars exposed to natural selection during pubertal development *Journal of Animal Science* 65 1164-1169

GR (1991) A characterization of Chinese pig clusters analysis *Journal of Animal Science* 73 388-394

GR, Wise TH and Ford JJ (1996) Identification of a subunit of thyroid-stimulating hormone in swine identified by differential screening *Journal of Animal Science* 82 2104-2111

GR and Ford JJ (1997) Breed differences in the expression of thyroid-stimulating hormone subunits in porcine anterior pituitary *Journal of Animal Science* 84 712-718

GR, Johnson RK and Allrich RD (1986) Testicular ultrastructure and function during development in the boar *Journal of Reproduction and Fertility* 58 115-121

GR and Ford JJ (1993) Changes in porcine testicular development in Meishan boars *Journal of Animal Science* 77 (Supplement 1) 126 (Abstract)

GR (1996) Selection for extremes in prolificacy results in reduced testis size and altered testicular ultrastructure in white composite boars. In *Advances in Pig Production Research* Vol. 2 pp. 523-531 Eds by G. M. Ross and J. C. M. Cook. Plenum Press, New York

Macdonald GJ, Wise T, Sluss PM, Lunstra DD and Ford JJ (1995) Differences in pituitary gonadotropins from male pigs representing divergent breeds *Journal of Reproduction and Fertility* 57 (Supplement 1) 77 (Abstract)

Mariscal DV, Wolfe PL, Bergfeld EG, Cupp AS, Kojima FN, Fike KE, Sanchez T, Wehrman ME, Johnson RK, Kittok RJ, Ford JJ and Kinner JE (1996) Comparison of circulating concentrations of reproductive hormones in boars of lines selected for size of testes or number of ovulations and embryonal survival to concentrations in respective control lines *Journal of Animal Science* 82 1155-1161

McNamara M, Vandalem JL, Hennen G, Coienbrander B, MacDonald AA, Van De Wiel D and Meijer JC (1985) The ontogeny of hypothalamic pituitary function in the pig. I Pituitary LH and FSH in the fetus and neonate *Molecular and Cellular Endocrinology* 39 119-124

Meachem SJ, McLachlan RI, de Kretser DM, Robertson DM and Wreford NG (1996) Neonatal exposure of rats to recombinant follicle stimulating hormone increases adult Sertoli and spermatogenic cell numbers *Journal of Reproduction and Fertility* 58 36-44

Okwun OE, Igboeli G, Ford JJ, Lunstra DD and Johnson L (1996a) Sertoli cell number and function, spermatogonial number and yield, and daily sperm production in three breeds of boars *Journal of Reproduction and Fertility* 107 137-149

Okwun OE, Igboeli G, Lunstra DD, Ford JJ and Johnson L (1996b) Testicular composition, number of A spermatogonia, germ cell ratios, and number of spermatids in three different breeds of boars *Journal of Andrology* 17 301-309

Peyrat J-P, Meusy-Dessolle N and Garnier J (1981) Changes in Leydig cells and luteinizing hormone receptors in porcine testis during postnatal development *Endocrinology* 108 625-631

Rohrer GA, Ford JJ, Wise TH and Young LD (1997) Estimates of genetic correlations between serum FSH concentrations in males and measures of female reproduction in a crossbred Meishan population of pigs *Journal of Animal Science* 84 (Supplement 1) 31 (Abstract)

Russell LD, Ping Ren H, Sinha Hikim I, Schultze W and Sinha Hikim AR (1990) A comparative study in twelve mammalian species of volume density, volumes, and numerical densities of selected testis components, emphasizing those related to the Sertoli cell *American Journal of Anatomy* 188 21-30

Schinckel AP, Johnson RK and Kittok RJ (1984) Testicular development and endocrine characteristics of boars selected for either high or low testis size *Journal of Animal Science* 58 675-685

Simorangkir DR, de Kretser DM and Wreford NG (1995) Increased numbers of Sertoli and germ cells in adult rat testes induced by synergistic action of transient neonatal hypothyroidism and neonatal hemicastration *Journal of Reproduction and Fertility* 104 207-213

Swanlund DJ, N'Diaye MR, Loseth KJ, Pryor JL and Crabo BG (1995) Diverse testicular responses to exogenous growth hormone and follicle-stimulating hormone in prepubertal boars *Journal of Reproduction and Fertility* 53 749-757

Tan HS and Raeside JJ (1980) Developmental patterns of plasma

dehydroepiandrosterone sulfate and testosterone in male pigs *Animal Reproduction Science* 3 73-81

Tarn CY, Rosenkrans CF, Jr and Kirby JD (1996) Effects of 6-N-propyl-2-thiouracil on boars. II Growth and reproductive traits *Journal of Animal Science* 82 (Supplement 1) 155 (Abstract)

Tilton JE, Biggs C, Hunter MG and Foxcroft GR (1994) Gonadotropin secretion in ovariectomized Chinese Meishan and hybrid large white gilts: responses to challenges with estradiol benzoate, gonadotropin-releasing hormone, or porcine follicular fluid *Journal of Reproduction and Fertility* 51 963-970

Tran D, Meusy-Dessolle N and Josso N (1981) Waning of anti-Mullerian activity: an early sign of Sertoli cell maturation in the developing pig *Journal of Reproduction and Fertility* 24 923-931

Trudeau VL, Meijer JC, Erkens JHF, Van De Wiel DFM and Wensing CJG (1992) Pubertal development in the male pig: effects of treatment with a long-acting gonadotropin-releasing hormone agonist on plasma luteinizing hormone, follicle stimulating hormone and testosterone *Canadian Journal of Veterinary Research* 56 102-109

van Straaten HWM and Wensing CJG (1977) Histomorphometric aspects of testicular morphogenesis in the pig *Journal of Reproduction and Fertility* 17 467-472

van Straaten HWM and Wensing CJG (1978) Leydig cell development in the testis of the pig *Journal of Reproduction and Fertility* 18 86-93

Wang RX (1990) Reproductive characteristics of the Fengjing pigs. In *Proceedings of Chinese Pig Symposium* Toulouse pp. 6-16 Eds M Molenat and C Legault, INRA (Institut National de la Recherche Agronomique), Jouy-en-Josas Cedex, France

White BR, McLaren DG and Dzuik PJ (1993) Age at puberty, ovulation rate, uterine length, prenatal survival and litter size in Chinese Meishan and Yorkshire females *Theriogenology* 40 85-97

Wise T, Lunstra DD and Ford JJ (1996) Differential pituitary and gonadal function of Chinese Meishan and European white composite boars: effects of GnRH stimulation, castration, and steroidal feedback *Journal of Reproduction and Fertility* 54 146-153

Young LD, Leymaster KA and Lunstra DD (1986) Genetic variation in testicular development and its relationship to female reproductive traits in swine *Journal of Animal Science* 63 17-26

Zanella EL, Ford JJ, Wise T and Hamernik DL (1996) Pituitary relationship of gonadotropins and messenger ribonucleic acid for gonadotropin subunits in white composite and Meishan boars *Journal of Reproduction and Fertility* 54 154-159

Zirkin BR, Johnson LL, Kromann N and Cochran RC (1980) Testosterone secretion by rat, rabbit, guinea pig, dog, and hamster testes perfused *in vitro*: correlation with Leydig cell ultrastructure *Endocrinology* 107 1867-1874

Zirkin BR, Awoniyi C, Griswold MD, Russell LD and Sharpe R (1994) Is FSH required for adult spermatogenesis? *Journal of Andrology* 15 273-276