

Functional Approach to Simulating Short-Rotation Woody Crops in Process-Based Models

Tian Guo¹ · Bernard A. Engel¹ · Gang Shao² · Jeffrey G. Arnold³ ·
Raghavan Srinivasan⁴ · James R. Kiniry³

Published online: 9 April 2015
© Springer Science+Business Media New York 2015

Abstract Short-rotation woody crops (SRWCs) such as *Populus* have great potential as biofuel feedstocks. Biomass yields and yield stability at potential sites are important considerations when SRWCs are widely planted. The process-based, daily time-step simulation model Agricultural Land Management Alternative with Numerical Assessment Criteria (ALMANAC) offers promise as a useful tool to evaluate tree growth over large ranges of conditions. The objective of this study was to develop algorithms and growth parameters of hybrid poplar ‘Tristis #1’ (*Populus balsamifera* L. × *Populus tristis* Fisch) and eastern cottonwood (*Populus deltoides* Bartr.) in ALMANAC and to improve simulation of leaf area index (LAI) and plant biomass as well as biomass partitioning. ALMANAC with the improved algorithms for LAI and weight of falling leaves was applied to hybrid poplar plots in Wisconsin and cottonwood plots in Mississippi, and the modeled biomass yield and LAI were compared with

measured data to modify and evaluate the location-specific ALMANAC models. Improved algorithms for LAI and biomass simulation and suggested values and potential parameter ranges for hybrid poplar and cottonwood were reasonable (Nash-Sutcliffe model efficiency (NSE) 0.81~0.99 and R^2 0.76~0.99). ALMANAC with modified algorithms and parameters for *Populus* growth realistically simulated LAI, aboveground woody biomass, and root biomass of *Populus*. Thus, this model can be used for biofeedstock production modeling for *Populus*. The improved algorithms of LAI and biomass simulation for tree growth should also be useful for other process-based models, such as Soil and Water Assessment Tool (SWAT), Environmental Policy Integrated Climate (EPIC), and Agricultural Policy/Environmental eXtender (APEX).

Keywords Bioenergy · Short-rotation woody crops · Hybrid poplar · Cottonwood · Process-based models · Biofuel production modeling

USDA is an equal opportunity provider and employer.

Electronic supplementary material The online version of this article (doi:10.1007/s12155-015-9615-0) contains supplementary material, which is available to authorized users.

✉ Tian Guo
guo190@purdue.edu

¹ Department of Agricultural and Biological Engineering, Purdue University, 225 South University Street, West Lafayette, IN 47907-2093, USA

² Department of Forest and Natural Resources, Purdue University, 715 West State Street, West Lafayette, IN 47907-2061, USA

³ Grassland Soil and Water Research Laboratory, USDA-ARS, 808 East Blackland Rd, Temple, TX 76502, USA

⁴ Spatial Sciences Laboratory, Texas A&M University, College Station, TX 76502, USA

Introduction

Increasing energy demand and high sustained oil prices have encouraged the use of alternative forms of energy. The majority of biofuel production in the USA comes from sugar-rich maize (*Zea mays* L.) and soybean (*Glycine max* (L.) Merr) oil. However, with the combination of a global increasing demand for renewable energy and food, the problems of food-fuel competition for land, higher food prices [1], and lower food production [2] will be created. Thus, beneficial biofuels should provide sustainable biofeedstocks that neither compete with food crops nor cause clearing of native forests. Non-food bioenergy crops—crop residues [3–7], cellulosic perennial crops (e.g., miscanthus (*Miscanthus* × *giganteus*), switchgrass

(*Panicum virgatum* L.), mixed grasses) [8–16], and woody biomass crops (e.g., *Populus*)—offer great potential [17].

Short-rotation intensive culture of trees is considered a promising way to increase wood biomass productivity [18]. Dry matter production of wood plus bark in short-rotation hardwood plantations are up to 20,000 kg/ha/year, three to five times more than that for some natural stands [19]. Interest has increased in growing short-rotation plantations for energy production, since the oil embargo in 1973 [20]. The *Populus* genus is highly productive under short-rotation intensive culture system and is a good raw material for reconstituted forest products, due to its genetic diversity, rapid growth, vegetative propagation ease, and coppice regeneration [21].

Biomass productivity may increase with narrower tree spacing under short-rotation intensive culture system. Strong and Hansen [22] concluded that biomass differences related to spacing were minor in hybrid poplar plantations with 18 clone/spacing combinations for up to 16-year growth periods in northern Wisconsin. Productivity of hybrid poplar was mainly influenced by clone, irrigation, and disease. Similarly, Cannell and Smith [23] showed that close spacing was not essential for high biomass yield of hybrid poplar. However, tree spacing can affect time to canopy closure and the time needed to achieve maximum mean annual biomass increment (MABI). Hybrid poplar trees with wide tree spacing have longer rotations and more flexible harvest scheduling as well as lower costs [22, 23].

Prediction of *Populus* growth is critical for managers and policy makers to establish and manage short-rotation woody crops (SRWCs) and to obtain high yields. Some researchers studied simulation of hybrid poplar growth using tree growth models. For instance, Ek [24] used a model for regression estimation of branch weights of *Populus* which was found to be more precise than the models based on branch diameter. An individual-tree-based stand simulation model, FOREST, was used to simulate the periodic growth of hybrid poplar and showed that plot design, establishment techniques, cultural and environmental factors, measurement procedures, and model limitation can explain differences between the projected and observed harvest [25]. Meldahl [26] modified the FOREST model to simulate biomass yields of hybrid poplar and reduce the differences between projected and observed values. Moreover, Landsberg and Wright [27] simulated annual biomass production of two hybrid *Populus* clones in two locations using an energy conversion which assumes that plant biomass is proportional to the radiant energy absorbed by the canopy. Use of a radiant energy equation, also used in the Agricultural Land Management Alternative with Numerical Assessment Criteria (ALMANAC) models as described below, resulted in better simulation performance of *Populus* biomass yields than other simulations based on tree branch weight or stand [27].

The ALMANAC model [28] is a process-based, daily time-step simulation model that has been parameterized and validated for a wide range of crop (corn and soybean), grass (switchgrass, miscanthus) and northern tree species [29] (lodgepole pine (*Pinus contorta* Douglas ex Loudon), white spruce (*Picea glauca* var. *glauca*), black spruce (*Picea mariana*), and trembling aspen (*Populus tremuloides* Michx.)). The model uses readily available USDA-NRCS soils data and readily available daily temperature and rainfall data. ALMANAC plant growth simulation processes include light interception, dry matter production, and biomass partitioned into plants [28, 30]. Biomass is calculated based on light interception and species-specific radiant use efficiency (RUE), which is the amount of dry biomass produced per unit of intercepted light [31, 32]. Three attributes useful for quantifying potential plant growth are as follows: RUE, LAI, and the light extinction coefficient (k) used to calculate the fraction of light intercepted by leaves [33].

Generally, RUE values for woody species are between 1.3 and 1.9 g/MJ intercepted photosynthetically active radiation (PAR) and for crops are between 2.2 and 3.5 g/MJ intercepted PAR [34]. Kiniry measured RUE values for eastern red cedar (*Juniperus virginiana*) (1.60 g/MJ intercepted PAR) and honey mesquite (*Prosopis glandulosa*) (1.61 g/MJ intercepted PAR) to allow better prediction of their growth in ALMANAC [33]. Mean RUE values were 1.5 for poplar in Wisconsin and Pennsylvania, USA [27], and RUE values were between 2.4 and 3.4 for intensively cultured poplar in Scotland [35]. The standard RUE values (g/MJ) should be multiplied by 10, to obtain the values (kg/ha)/(MJ/m²) used in the ALMANAC and Soil and Water Assessment Tool (SWAT) [36].

Nineteen parameters for annual and long-term forest growth were incorporated and modified in the model to simulate successional forest regrowth after disturbance of forest ecosystems. Ranges of parameters were derived from scientific literature or yield tables. The ranges of RUE and k values for mixed forest used in ALMANAC were determined as 15–20 and 0.5–0.55, respectively [37, 29]. However, research on biomass yields of trees simulated by ALMANAC is limited, since parameters and equations modified in the model are for mixed forest stands consisting of various woody species rather than a specific woody species [29].

Moreover, accurate LAI, biomass yield, and biomass partitioning simulation for *Populus* in ALMANAC has not been adequately developed, and it is important to quantify fast-growing tree growth accurately. In ALMANAC and SWAT, leaf area development, a sigmoid curve, is a function of the growing season for mature plants, during which mature plants can reach maximum LAI with the increase of heat units [36]. As LAI for juvenile trees cannot increase to maximum LAI, the leaf area algorithm used in the model was not suitable for juvenile tree growth simulation. Thus, ALMANAC can only simulate

Fig. 1 Location of hybrid poplar site at the USDA Forest Service Harshaw Experimental Farm near Rhinelander, WI, and cottonwood site at the Delta Research and Extension Center at Stoneville, MS



plant growth after plants reach maturity [36]. However, SRWCs were usually harvested once they reach maturity or even before maturity [21]. Thus, it is also important to improve the model to reasonably simulate tree growth from tree planting to maturity.

This work is a first effort to improve *Populus* growth algorithms and parameters in ALMANAC with published region-specific *Populus* growth data. The objectives of this study were to (1) develop algorithms and growth parameters of hybrid poplar ‘Tristis #1’ (*Populus balsamifera* L.

Table 1 Data for hybrid poplar and cottonwood growth simulation by ALMANAC

Plant	Data type	Source	Format	Date
Hybrid poplar	SSURGO	USDA Web Soil Survey	Polygon shapefile	
	Precipitation and temperature	NCDC		1970–1980
	Annual aboveground woody biomass yield (metric ton (t)/ha)	Scientific literature ^a		1970–1980
	Annual LAI	Scientific literature ^a		1970–1980
Cottonwood	SSURGO	USDA Web Soil Survey	Polygon shapefile	
	Precipitation and temperature	NCDC		1995–1997
	Annual aboveground biomass yield (t/ha)	Unpublished report ^b		1995–1997
	Annual root biomass (t/ha)	Unpublished report ^b		1995–1997

SSURGO, Soil Survey Geographic Database, USDA US Department of Agriculture, NCDC National Climate Data Center

^a Hansen [21]

^b Pettry et al. (1997), unpublished annual progress report

Table 2 Management operations for hybrid poplar site at the USDA Forest Service Harshaw Experimental Farm near Rhinelander, WI

Plant	Date	Management operation	Rate (kg/ha)
Hybrid poplar	30 May	Tillage, roto-tiller (mixing depth 5 mm, mixing efficiency 0.80)	
	1 June	Planting	
	1 June	Pesticide application (as linuron)	2.2 ^{a,b}
	1 June	Nitrogen application (as anhydrous ammonia)	200 ^{a,b}
	1 June	Phosphorus application (as elemental phosphorus)	50 ^{a,b}
	31 Dec	The end of the operation scheduling for a year	

^a Ek and Dawson [40]^b Srinivasan and Cibir (2014), personal communication

× *Populus tristis* Fisch) and eastern cottonwood (*Populus deltoides* Bartr.) in ALMANAC and to improve simulation of leaf area and plant biomass as well as biomass partitioning, (2) use the modified model to simulate LAI and aboveground woody biomass of hybrid poplar in Wisconsin and aboveground woody biomass and root biomass of cottonwood in Mississippi, and (3) compare simulated LAI and biomass results from the modified model with observed values for verification of improved algorithms and growth parameters of *Populus*.

Materials and Methods

Hybrid Poplar Site in Northern Wisconsin and Cottonwood Site in Western Mississippi

This study was conducted using data in the literature from two study sites (Fig. 1). The poplar site was a short-rotation intensive culture plantation at the USDA Forest Service Harshaw Experimental Farm near Rhinelander, WI, USA (45.6° N, 89.5° W) [38, 39]. Hybrid poplar cuttings were planted in early June, 1970, on a prepared site [40]. The site was sowed to rye, plowed, and rototilled before planting [22]. The soil of the plantation is the Padus series, a silt loam, overlaying sand, and gravel at depths of 30 to 60 cm with slope reaching at

most 1 %. The pH is from 6.7 to 7.0 [40]. The average growing season of hybrid poplar in this region is 120 days.

The cottonwood site was at the Delta Research and Extension Center at Stoneville, Mississippi, in the Tennessee Valley region [41], which was on agricultural land with a Bostket silt loam soil, a fine loamy, mixed, thermic Mollic Hapludalfs. The slope gradient is 0.2 %. Soil quality changes were determined based on soil physical characteristics measured at the site in 1995 (prior to tree establishment) and in 1997 (at the end of growing season) [42]. Cottonwood cuttings 20–30 cm long were planted with spacing of 1.2×3.6 m (population 23 trees/100 m²) on 3 February, 1995 [43], and harvested during 1–20 November, 1997 (Pettry et al. 1997, unpublished annual progress report).

ALMANAC Model Setup and Management Schedules

ALMANAC 2011 (Version 1.0.3 Beta 2) with Interface (Version 1.0.3) was used in this project. A new crop named “Poplar Tian Low” and “Cottonwood” were added to represent hybrid poplar and cottonwood, respectively. Lat 45.6° and long 89.5°, and lat 33.34° and long 90.85° were used for the hybrid poplar and cottonwood sites, respectively. The fraction of total tree biomass partitioned to roots was assumed to be 0.5 for hybrid poplar [21] and 0.2 for cottonwood (Pettry et al. 1997, unpublished annual progress report). Table 1 describes the primary data required for ALMANAC model setup.

Table 3 Management operations for cottonwood site at the Delta Research and Extension Center at Stoneville, MS

Plant	Date	Management operation	Rate (kg/ha)
Cottonwood	3 Feb	Tillage, roto-tiller (mixing depth 5 mm, mixing efficiency 0.80)	
	3 Feb	Planting	
	3 Feb	Pesticide application (as linuron)	2.2 ^{a,b}
	1 June	Nitrogen application (as anhydrous ammonia)	200 ^{a,b}
	1 June	Phosphorus application (as elemental phosphorus)	30 ^{a,b}
	31 Dec	The end of the operation scheduling for a year	

^a Thornton et al. [43] and Joslin and Schoenholtz [41]^b Srinivasan and Cibir (2014), personal communication

Table 4 Hybrid poplar and cottonwood growth data for model calibration and validation

<i>Populus</i>	Population (trees/100 m ²)	Density level	Outputs (annual aboveground woody biomass (AAWB), LAI, annual aboveground biomass (AAB), and root biomass (RB))	Data usage
Hybrid poplar	278	High	LAI	Model calibration
	278	High	AAWB (t/ha)	
	69	Medium	AAWB (t/ha)	
	17	Low	LAI	
	17	Low	AAWB (t/ha)	
	1111	High	AAWB (t/ha)	Model validation
	83	High	LAI	
	83	High	AAWB (t/ha)	
	25	Medium	LAI	
	25	Medium	AAWB (t/ha)	
	8	Low	AAWB (t/ha)	
Cottonwood	23	Medium	AAB (t/ha)	
	23	Medium	RB (t/ha)	

ALMANAC management includes planting and end of schedule dates, yearly tillage, pesticide, and nutrient application rates. Tables 2 and 3 represent management operations for hybrid poplar growth in 1970 and cottonwood growth in 1995. Fertilizer and auto irrigation were also added to these two location-specific models to ensure that *Populus* growth was not under water stress or nutrient stress. Nutrient application dates and rates for hybrid poplar growth from years 1971 to 1980 and cottonwood growth during years 1996 and 1997 were the same as nitrogen and phosphorus application in Tables 2 and 3, respectively. Hybrid poplar planting was on 22 May, 1970, and harvest was on 1 May, 1980. Cottonwood planting was on 3 February, 1995, and harvest was on 30 Nov, 1997.

Algorithm and Parameter Changes in the Model

Deciduous tree LAI increases both within each growing season prior to late-season senescence and among years as the maximum seasonal LAI increases. The seasonal leaf area development curve in the model can be used in years prior to maturity year after adjusting each year's potential LAI. Published yearly

LAI values for *Populus* trees with various planting densities ranged from 8 to 1111 trees/100 m² (Table 10 of Appendix). The increase in maximum seasonal LAI across years for *Populus* with various densities was similar to the equation of loss of leaf late in the season [44]. This served as the starting point to derive a new leaf development algorithm to simulate maximum seasonal LAI each year with various densities.

$$y_{yr} = y_{yr-1} * 10^{\log_{10}\left(\frac{y_r}{x_1}\right) * x_2} \quad (1)$$

where yr is the current growth year, y_{yr} is the LAI value for current year, y_{yr-1} is the LAI value for previous year, x_1 is the number of years until maximum LAI is attained (CLAIYR), and x_2 is a new tree leaf factor (TreeD) in the LAI algorithm, representing how LAI increases to the maximum potential LAI (DMLA) with varying densities.

CLAIYR values for *Populus* trees with various densities were obtained from a previous study [21]. A specific density of *Populus* trees has an associated TreeD value representing its LAI development. Based on published LAI values for different years and CLAIYR values, TreeD in Eq. (1) was

Table 5 Hybrid poplar tree growth parameters for various spacings for used in LAI simulation in the modified ALMANAC

Population (trees/100 m ²)	Spacing (m×m)	DMLA (maximum LAI) in ALMANAC)	Observed DMLA	TreeD (LAI factor)	CLAIYR (year to attain maximum LAI)
1111	0.3×0.3	9.5	8.6	0.5	6
278	0.6×0.6	9.5	8.6	0.75	6
83	1.1×1.1	9.5	8.6	1.5	6
69	1.2×1.2	9.5	8.6	2.5	6
25	2×2	9.5	8.6	3	6
17	2.4×2.4	9.5	8.6	2	7
8	3.6×3.6	9.5	8.6	4.5	9




Fig. 2 ALMANAC TreeD parameters for hybrid poplar trees with various spacings

calibrated manually for various populations to match observed values.

The management parameter “POPULATION” is the number of trees per 100 m². Previously, ALMANAC did not include a specific parameter for population effects on maximum seasonal LAI over years. In this new version, TreeD values in the crop database are used for different populations for *Populus* trees to calculate these seasonal maximums.

Total tree biomass consists of root biomass, senescent dropped leaf weight, and aboveground biomass (leaves, stems, and branches). To accurately simulate *Populus* tree biomass partitioning, the algorithm used for dropping leaves was improved (see details in the [Appendix](#)).

Values and Ranges of Parameters Determined Before Model Calibration

Two-week moving average daily temperatures at the USDA Forest Service Harshaw Experimental Farm in Wisconsin and the

Stoneville site in Mississippi were obtained using Matlab 2013 based on NOAA daily temperature data to determine base temperature (TG). The period of emergence was assumed from 1 to 20 April for hybrid poplar and 20 March to 10 April for cottonwood [45–47], which were day of year 90 to 110 and 78 to 98, respectively.

Values of potential heat unit (PHU) for hybrid poplar growth in Wisconsin and cottonwood growth in Mississippi were calculated based on accumulation of heat units during the growing season [48]. The growing season of hybrid poplar on the Harshaw experiment farm and cottonwood at the Stoneville site was assumed from 1 April to 11 October and from 20 March to 31 October, respectively [45–47] (see details in [Appendix](#)).


Values of hybrid poplar and cottonwood growth parameters maximum rooting depth (RDMX); rate of decline in RUE per unit increase in vapor pressure deficit (WAVP); plant nitrogen (N) at emergence (BN₁), 50 % maturity (BN₂), and maturity (BN₃); phosphorus fraction at emergence (BP₁), 50 % maturity (BP₂), and maturity (BP₃) [33, 29]; and harvest index (HI) for optimal growing conditions [36, 46] were derived from previous studies (see details in [Appendix](#)).

Values of plant maximum stomatal conductance (GSI) and maximum canopy height (HMX) for *Populus* growth simulation in the model were assumed before model calibration based on personal communication (Kiniry 2014) (see details in [Appendix](#)).

ALMANAC Model Calibration and Parameterization

Previous hybrid poplar growth studies at the USDA Forest Service Harshaw Experimental Farm in Wisconsin [49, 21, 27, 50, 51] suggested values for RUE (called WA in the model), *k* (called EXTINC in the model), DMLA, two points on optimal leaf development curve parameters (DLAP1 and DLAP2), fraction of growing season when leaf area starts declining (DLAI), plant N fraction in harvested biomass

Fig. 3 Simulated LAI curve for hybrid poplar trees with various populations (number in parentheses is population (trees/100 m²) of hybrid poplar trees)






Fig. 4 Two-week moving average daily temperatures at Harshaw Experiment Farm in Wisconsin and at Stoneville Site in Mississippi (gray bands are periods of emergence). **a** Hybrid poplar site in Wisconsin. **b** Cottonwood site in Mississippi

(CNY), and plant P fraction in harvested biomass (CNP) (see details in [Appendix](#)), providing reasonable ranges of these tree growth parameters for model calibration. Ranges of PHU values were calculated before model calibration. The model was calibrated by changing these *Populus* growth parameters manually to obtain a good fit with published hybrid poplar LAI and aboveground biomass values. Values of WA, EXTINC, DMLA, DLAP1 and DLAP2, DLAI, CNY, and PHU were determined after model calibration.

The LAI and aboveground woody biomass data of hybrid poplar with various spacings and aboveground biomass and root biomass of cottonwood with medium density used for model calibration and validation are summarized in Table 4.

Validation of the Modified ALMANAC Model

The methods used for verifying the model performance [52] include percent bias/percent error (P_{BIAS} [%]), Nash-Sutcliffe model efficiency (NSE) coefficient, and coefficient of determination (R^2). Value of P_{BIAS} [53] is a measure of the average tendency of the simulated data to be larger or smaller than the measured data. The value of 0.0 is the optimal value of P_{BIAS} . Negative values represent overestimation bias, and positive values represent underestimation bias. The NSE [54] describes how well measured versus simulated data plots match the 1:1 line. The NSE value ranges from $-\infty$ to 1, and the optimal value is 1. We assumed that a NSE value of greater than 0.5 meant that model performance is satisfactory [55]. Values of $0.36 \leq NSE \leq 0.72$ and $NSE \geq 0.75$ also have been considered satisfactory and good simulated results, respectively [56, 57]. The R^2 value indicates the strength of the linear relationship between the measured and simulated data. We assumed that an R^2 value of greater than 0.5 indicated reasonable model performance [55].

Results and Discussion

Algorithm and Parameter Changes in the Model

Leaf area cover, as defined by leaf area index (LAI), is a driving variable determining amount of light intercepted and, thus, biomass via the RUE approach. Simulated LAI also drives potential transpiration, an important component of the total evapotranspiration of the system. Deciduous tree LAI increases both within each growing season prior to late-season senescence and among years. Values for LAI also vary with planting density of trees. Within each growing season, LAI decreases late in the season with leaf senescence. Tree spacing was converted to population (Table 5). TreeD, CLAI YR, observed DMLA, and DMLA for various spacings used in LAI simulation in the modified ALMANAC are shown in Table 5. For high-density (population of 1111, 278 or 83 trees/100 m²) and medium-density (population of 69 or 25 trees/

Table 6 Potential heat units for *Populus* during each growing season of different years

Plant	Year	PHU
Hybrid poplar	1974	1670
	1975	1999
	1976	2047
	1977	2149
	1978	1956
	1979	1893
	1980	1986
Cottonwood	1995	2899
	1996	2818
	1997	2421

Table 7 Suggested values and potential parameter ranges for hybrid poplar and cottonwood compared to current parameters for *Populus* in ALMA NAC crop database

Parameter Acronym in ALMANAC	Parameter definition	Hybrid poplar ‘Tristis #1’ <i>Populus balsamifera</i> L. × <i>Populus tristis</i> Fisch (HYPT)	
		Suggested value	Range
TG ^a [PHU] ^{a,c}	Base temperature (°C) Heat units to maturity	4 [1750]	0–6 [2150–1500]
TB ^b	Optimal temperature (°C)	25	25–30
WA ^{c,d}	Radiation use Efficiency in ambient CO ₂ (kg/ha)/(MJ/m ²)	20	20–35
EXTINC ^{c,d}	Light extinction coefficient	0.30	0.20–0.60
DMLA ^{c,e,f}	Maximum LAI	9.50	5.00–9.50
DLAI ^{c,e,f}	Point in growing season when LAI declines	0.99	0.99
TREED ^{c,e}	Tree leaf area decline factor	0.500–4.500	0.500–4.500
BP ₁ ^{g,h}	Plant P fraction at emergence (whole plant)	Existing value	Existing value
GSI ^b	Maximum stomatal conductance	0.0070	0.0040–0.0070
HMX ^b	Maximum canopy height	Existing value	7.00–15.00
BN ₁ ^{g,h}	Plant N fraction at emergence (whole plant)	Existing value	Existing value
BN ₃ ^{g,h}	Plant N fraction at maturity (whole plant)	Existing value	Existing value
BN ₂ ^{g,h}	Plant N fraction at 50 % maturity (whole plant)	Existing value	Existing value
RDMX ^{g,h}	Maximum rooting depth	Existing value	Existing value
CNY ^{c,i,j}	Plant N fraction in harvested biomass	0.0005	0.0005–0.0015
CPY ^{c,k}	Plant P fraction in harvested biomass	0.0002	0.0002–0.0003
BP ₂ ^{g,h}	Plant P fraction at 50 % maturity (whole plant)	Existing value	Existing value
BP ₃ ^{g,h}	Plant P fraction at maturity (whole plant)	Existing value	Existing value
WAVP ^{g,h}	Rate of decline in RUE per unit increase in vapor pressure deficit	Existing value	Existing value
CHTYR ^{c,f}	Number of years required for tree species to reach full development (years)	6–9	6–9
HI ^{l,m}	Harvest index for optimal growing conditions	0.65	0.45–0.70
Optimal leaf development curve parameters			
DLAP1 ^{c,e,f}	Fraction of growing season coinciding with first point	0.05	0.05–0.07
	Fraction of DMLA corresponding to first point	0.05	0.05–0.30
DLAP2 ^{c,e,f}	Fraction of growing season coinciding with second point	0.40	0.40–0.45
	Fraction of DMLA corresponding to second point	0.95	0.95–0.98
Parameter Acronym in ALMANAC	Eastern cottonwood <i>Populus deltoides</i> Bartr. (POEC)		<i>Populus</i> (POPL)
	Suggested value	Range	Database value
TG ^a [PHU] ^{a,c}	8 [2818]	7–15 [2900–2200]	10–
TB ^b	25	25–30	30
WA ^{c,d}	41	30–58	30
EXTINC ^{c,d}	0.60	0.20–0.60	0.45
DMLA ^{c,e,f}	9.50	5.00–9.50	5.00
DLAI ^{c,e,f}	0.99	0.99	0.99
TREED ^{c,e}	0.500–4.500	0.500–4.500	
BP ₁ ^{g,h}	Existing value	Existing value	0.0007
GSI ^b	0.0070	0.0040–0.0070	0.0040
HMX ^b	10.00	10.00–15.00	7.50
BN ₁ ^{g,h}	Existing value	Existing value	0.0060
BN ₃ ^{g,h}	Existing value	Existing value	0.0015
BN ₂ ^{g,h}	Existing value	Existing value	0.0020
RDMX ^{g,h}	Existing value	Existing value	3.50
CNY ^{c,i,j}	0.0005	0.0005–0.0015	0.0015
CPY ^{c,k}	0.0002	0.0002–0.0003	0.0003
BP ₂ ^{g,h}	Existing value	Existing value	0.0004
BP ₃ ^{g,h}	Existing value	Existing value	0.0003
WAVP ^{g,h}	Existing value	Existing value	8.00

Table 7 (continued)

Parameter Acronym in ALMANAC	Eastern cottonwood <i>Populus deltoides</i> Bartr. (POEC)		<i>Populus</i> (POPL)
	Suggested value	Range	Database value
CHTYR ^{c,f}	6–9	6–9	10
HI ^{l,m}	0.60	0.40–0.65	0.76
Optimal leaf development curve parameters			
DLAP1 ^{c,e,f}	0.05	0.05–0.07	0.05
	0.05	0.05–0.30	0.05
DLAP2 ^{c,e,f}	0.40	0.40–0.45	0.40
	0.95	0.95–0.98	0.95

^a Maximum and minimum daily temperature from NOAA^b Assumed^c Modified parameter from hybrid poplar growth simulation^d Landsberg and Wright [27]^e Hansen [21]^f Zavitkovski [51]^g Kiniry [32]^h MacDonald et al. [37]ⁱ Black et al. [49]^j McLaughlin et al. [50]^k Kiniry (2014), personal communication^l Michael et al. [47]^m Arnold et al. [36]

100 m²) hybrid poplar trees, a shorter time (6 years) is needed to attain DMLA. For low-density (population of 17 or 8 trees/100 m²) hybrid poplar trees, a longer time (7 or 9 years) is needed to attain DMLA.

Based on TreeD and tree spacing values (Table 5) for high- and medium-density hybrid poplar trees (Fig. 2), TreeD is linearly related to tree spacing (Eq. (2)). Equation (2) was assumed suitable for short-rotation *Populus* trees which can attain DMLA in 6 years. For *Populus* trees attaining DMLA in 6 years, the higher tree spacing (smaller tree population) is associated with higher TreeD values. For *Populus* trees which attain DMLA in 7 to 9 years, the TreeD values can be found in Table 5.

$$y = 1.579 * x + 0.007, R^2 = 0.898 \quad (2)$$

where y is the TreeD parameter and x is the tree planting spacing (m).

Maximum seasonal LAI values of hybrid poplar with different populations calculated by Eq. (1) are shown in Fig. 3. For high-density hybrid poplar trees, LAI can increase significantly at the beginning (years 1 to 3) and attain maximum LAI in a shorter time (6 years). For low-density hybrid poplar trees, LAI increases slowly at the beginning (years 1 to 3) and attains maximum LAI at a later time (7 or 9 years). Thus, tree

spacing can affect time to canopy closure, and wide tree spacing allows longer rotations of hybrid poplar if canopy closure by harvest year is desirable. This is consistent with results of Strong and Hansen [22].

The algorithm used for dropping leaves was improved to more accurately simulate weight of dropping leaves.

$$y = x_1 * \left(0.4 - 0.3 * \frac{yr - 1}{x_2 - 1} \right) \quad (3)$$

where y is the weight of dropping leaves, yr is the current growth year, x_1 is the aboveground biomass, and x_2 is the number of years to maximum height and maximum LAI of trees (CHTYR).

Values and Ranges of Parameters Determined Before Model Calibration

Two-week moving average daily temperature plots of hybrid poplar and cottonwood growth are shown in Fig. 4. The gray bands were the period of emergence for hybrid poplar (Fig. 4a) and cottonwood (Fig. 4b) growth, respectively. Temperatures in the gray bands were the ranges of TG for *Populus* growth. TG was chosen as 4 °C for hybrid poplar (within the expected range 0–6 °C (Srinivasan R 2014, personal




Fig. 5 Yearly observed and calibrated ALMANAC (modified) simulated LAI during calibration of hybrid poplar with populations of 278 (a) and 17 (b) trees/100 m²

communication)) and 8 °C for cottonwood. The generic optimal temperature for warm season plants, 25 °C, was chosen for optimal temperature (TB) of hybrid poplar and cottonwood growth in this study [36].

The PHU values were calculated during each growing season from 1974 to 1980 for hybrid poplar and from 1995 to

1997 for cottonwood (Table 6). The range of possible values for PHU is 1670–2150 for hybrid poplar in Wisconsin and 2421–2899 for cottonwood in Mississippi.

Assumed values and ranges of RDMX, WAVP, BN₁, BN₂, BN₃, BP₁, BP₂, BP₃, HI, GSI, and HMX for hybrid poplar and cottonwood growth are summarized in Table 7.




Fig. 6 Yearly observed and calibrated ALMANAC (modified) simulated aboveground woody biomass during calibration of hybrid poplar with populations of 278 (a), 69 (b), and 17 (c) trees/100 m²




Fig. 7 Yearly observed and calibrated ALMANAC (modified) simulated LAI during validation of hybrid poplar with populations of 83 (a) and 25 (b) trees/100 m²

ALMANAC Model Calibration for Hybrid Poplar Growth in Wisconsin

Calibrated annual LAI values by ALMANAC were compared with published values for hybrid poplar with

populations of 278 (high density) and 17 (low density) trees/100 m² (Fig. 5). Calibrated annual aboveground woody biomass values were compared with published values for hybrid poplar with populations of 278, 69 (medium density), and 17 trees/100 m² (Fig. 6).




Fig. 8 Yearly observed and calibrated ALMANAC (modified) simulated aboveground woody biomass during validation of hybrid poplar with populations of 1111 (a), 83 (b), 25 (c), and 8 (d) trees/100 m²




Fig. 9 Yearly observed and calibrated ALMANAC (modified) simulated aboveground biomass (a) and root biomass (b) during validation of cottonwood with a population of 23 trees/100 m²

Simulated annual LAI of hybrid poplar with populations of 278 and 17 trees/100 m² had a reasonable good match with observed values, except that the simulated LAI value at year 9 (population of 17 trees/100 m²) was slightly higher than the observed value (Fig. 5).

Projected annual aboveground woody biomass of hybrid poplar with populations of 278, 69, and 17 trees/100 m² fit the observed values reasonably well, except that simulated annual aboveground woody biomass values at years 2 and 3 (population of 278 trees/100 m²) were higher than the observed values (Fig. 6). Simulated annual aboveground woody biomass values at years 2, 3, and 4 (population of 17 trees/100 m²) were higher than the observed values.

Suggested Values and Potential Parameter Range for Hybrid Poplar and Cottonwood in ALMANAC Model

ALMANAC realistically simulated annual LAI and aboveground woody biomass yield of hybrid poplar with various spacings. Suggested values and potential parameter ranges for hybrid poplar and cottonwood were determined (Table 7).

Existing values or ranges of growth parameters used in ALMANAC are globally approximated, since it is time-consuming and difficult to obtain growth data in detail for each species. The existing values or ranges can be adjusted in specific regions before being used for plant growth simulation.

Modified ALMANAC Model Validation for Hybrid Poplar and Cottonwood Growth

Comparisons of annual LAI values modeled by the modified ALMANAC with published values for hybrid poplar with populations of 83 (high density) and 25 (medium density) trees/100 m² are shown in Fig. 7. Comparisons of annual aboveground woody biomass modeled values with published values for hybrid poplar with populations of 1111 (high density), 83, 25, and 8 (low density) trees/100 m² are shown in Fig. 8. Comparisons of modeled annual aboveground biomass and root biomass with published values for cottonwood with a population of 23 trees/100 m² (medium density) are shown in Fig. 9. The modified model was validated based on the percent

Table 8 Evaluation of model outputs with various populations for the modified ALMANAC

Plant	Population (trees/100 m ²)	Density level	Outputs Aboveground woody biomass (AWB), LAI, aboveground biomass (AB), root biomass (RB)	P _{BIAS} (%)	NSE	R ²
Hybrid poplar	1111	High	AWB (t/ha)	2	0.81	0.98
			LAI	4	0.96	0.76
	25	Medium	AWB (t/ha)	−9	0.95	0.79
			LAI	−11	0.98	0.98
			AWB (t/ha)	−22	0.96	0.96
			AWB (t/ha)	1	0.99	0.99
Cottonwood	23	Medium	AB (t/ha)	−0.3	0.99	0.99
			RB (t/ha)	2	0.99	0.99

bias (P_{BIAS} , %), Nash-Sutcliffe model efficiency (NSE), and coefficient of determination (R^2) methods. Evaluation results of modeled outputs are shown in Table 8. Projected MABI values by the modified ALMANAC were compared with measured yields and projected values from the original ALMANAC and FOREST and modified FOREST models for hybrid poplar growth in Rhineland, WI (Table 9).

Projected annual LAI of hybrid poplar with populations of 83 and 25 trees/100 m² had a good match with observed values (Fig. 7). Moreover, NSE (R^2) values for modeled LAI of hybrid poplar with populations of 83 and 25 trees/100 m² were 0.96 (0.76) and 0.98 (0.98), respectively (Table 8). Overall performance of the modeled LAI of hybrid poplar (83 and 25 trees/100 m²) was satisfactory (since $NSE \geq 0.75$ and $R^2 \geq 0.5$). The optimal value of P_{BIAS} is 0, and 4 % (83 trees/100 m²) was close to 0, which also represented accurate model simulation. However, $P_{BIAS} = -11$ % (25 trees/100 m²) meant that simulated annual LAI results were slightly overestimated, which also can be found from Fig. 7b. Simulated annual LAI values for years 3 and 4 were higher than observed values.

Overall performance of the modeled aboveground woody biomass yields of hybrid poplar (1111, 83, 25, and 8 trees/100 m²) was satisfactory (since $NSE \geq 0.75$ and $R^2 \geq 0.5$). Projected annual aboveground woody biomass of hybrid poplar with populations of 1111, 83, 25, and 8 trees/100 m² fit observed values well (Fig. 8). Moreover, NSE (R^2) values for simulated aboveground woody biomass of hybrid poplar with populations of 1111, 83, 25, and 8 trees/100 m² were 0.81 (0.98), 0.95 (0.79), 0.96 (0.96), and 0.99 (0.99), respectively (Table 8). P_{BIAS} values of aboveground woody biomass of hybrid poplar with populations of 1111 and 8 trees/100 m² were 2 and 1 %, which also represented accurate model simulation. However, P_{BIAS} values of hybrid poplar with populations of 83 and 25 trees/100 m² were -9 and -22 % respectively, indicating that modeled annual aboveground woody biomass results were slightly overestimated, which also can be found from Fig. 8b (83 trees/100 m²) and Fig. 8c (25 trees/100 m²). Modeled annual aboveground woody biomass for years 2 and 3 was higher than observed values.

Projected annual aboveground biomass and root biomass of cottonwood with a population of 23 trees/100 m² fit the observed values well (Fig. 9). Moreover, NSE (R^2) values for modeled aboveground biomass and root biomass of cottonwood were 0.99 (0.99) and 0.99 (0.99), respectively (Table 8). Overall performance of the modeled aboveground and root biomass yields of cottonwood was satisfactory (since $NSE \geq 0.75$ and $R^2 \geq 0.5$). P_{BIAS} values of modeled aboveground and root biomass were -0.3 and 2 %, respectively, which also represented accurate model simulation.

Performance of MABI simulation by the modified ALMANAC was superior to the original ALMANAC and FOREST and the modified FOREST models. Measured MABI of the 5-year-old hybrid poplar planting with a population of 69 trees/

Table 9 Comparison of projected and measured MABI of 5-, 9- and 10-year-old short-rotation intensively cultured hybrid poplar grown with various spacings in Wisconsin

Variables	Age (year)	Spacing (m×m)	Population (trees/100 m ²)	Measured harvest	Modeled yields (t/ha/year)		
					Modified ALMANAC	ALMANAC	FOREST
MABI (t/ha/year)	5	1.2×1.2	69	7.6 ^a	7.0 ^b (-8 %)	10.0 ^b (32 %)	10.8 ^{c,d} (42 %)
MABI (t/ha/year)	10	2.4×2.4	17	10.4 ^c	9.2 ^b (-12 %)	1.9 ^b (-82 %)	20.4 ^{c,d} (96 %)
MABI (t/ha/year)	9	3.6×3.6	8	6.2 ^e	7.3 ^b (18 %)	2.2 ^b (-65 %)	17.5 ^{c,d} (182 %)

Number in parentheses represents rate of increase/decrease of simulated results to related measured results

^a Isebrands et al. [59]

^b Present study

^c Ek and Dawson [40]

^d Ek and Dawson [58]

^e Hansen [21]

^f Meldahl [26]

100 m² was 7.6 t/ha/year (Table 9). The modified ALMA NAC, original ALMANAC, and FOREST [40, 58] projections were 8 % (7.0 t/ha/year) lower, 32 % (10.0 t/ha/year) higher, and 42 % (10.8 t/ha/year) higher than the measured value, respectively.

Additionally, measured MABI of the 10-year-old hybrid poplar planting with a population of 17 trees/100 m² was 10.4 t/ha/year (Table 9). The modified ALMANAC, original ALMANAC, and FOREST [40, 58] and the modified FOREST [26] projections were 12 % (9.2 t/ha/year) lower, 82 % (1.9 t/ha/year) lower, 96 % (20.4 t/ha/year) higher, and 81 % (18.8 t/ha/year) higher than the measured value, respectively.

Measured MABI of the 9-year-old hybrid poplar planting with a population of 8 trees/100 m² was 6.2 t/ha/year (Table 9). The modified ALMANAC, original ALMANAC, and FOREST [40, 58] projections were 18 % (7.3 t/ha/year) higher, 65 % (2.2 t/ha/year) lower, and 182 % (17.5 t/ha/year) higher than the measured value, respectively.

Conclusions

SRWCs such as hybrid poplar and cottonwood are important biofuel feedstocks. To simulate biomass yields of hybrid poplar and cottonwood appropriately, the functional components and parameters of hybrid poplar and cottonwood were determined, and related algorithms improved in ALMANAC for leaf area, plant biomass, and biomass partitioning. The improved tree growth simulation in ALMANAC was applied to hybrid poplar plots in Wisconsin and cottonwood plots in Mississippi. The simulated LAI, total biomass, and biomass partitioning between above-ground and roots were compared with published data to modify and evaluate the location-specific ALMANAC model parameters.

Simulated aboveground woody biomass and LAI results from the modified ALMANAC for the hybrid poplar site with various spacings in Wisconsin were satisfactory (P_{BIAS} -22~4, NSE 0.81~0.99, and R^2 0.76~0.99). Additionally, modeled aboveground biomass and root biomass for the cottonwood site in Mississippi were good (P_{BIAS} -0.3~2, NSE 0.99~0.99, and R^2 0.99~0.99). Generally, simulations by the modified ALMANAC model of LAI and biomass yield of *Populus* were good (P_{BIAS} -22~4, NSE 0.81~0.99, and R^2 0.76~0.99) and improved relative to simulations by the original ALMANAC, FOREST, and modified FOREST models. Thus, the new algorithm for estimating LAI development for *Populus* (Eq. (1)), the new equation for calculating falling leaves weight (Eq. (3)), and suggested values of newly added parameter tree leaf factor (Table 5 and Eq. (2)) for various populations (high, medium, and low density) were reasonable. The suggested values and potential parameter range for hybrid poplar and cottonwood (Table 7) were reasonable, which provide guidance for simulation of poplar growth in the

midwestern USA and cottonwood growth in the southern USA. The modified ALMANAC model is able to simulate biofeedstock production of juvenile and mature *Populus* trees with various populations. The improved algorithms of LAI and biomass simulation for tree growth could also be used in other process-based models, such as Soil and Water Assessment Tool (SWAT), Environmental Policy Integrated Climate (EPIC), and Agricultural Policy/Environmental eXtender (APEX).

The LAI and biomass yield data of *Populus* trees used in this work were from previous studies during 1970–1980 or 1995–1997. The data were limited (for some tree populations, only 4-year data were observed). Moreover, tree planting techniques and applied pesticide were different from those in recent hybrid poplar trials. Short-rotation woody crop growth models and parameters could potentially be improved using additional *Populus* tree growth data. Moreover, suggested ranges and values for *Populus* growth parameters could be adjusted in specific regions before being used for tree growth simulation.

Acknowledgements We thank Amber Williams and Daren Harmel with USDA-ARS for model setup and comments on this manuscript. We thank Lynn Wright with WrightLink Consulting Inc and Oak Ridge National Laboratory for providing data and suggestions to this manuscript.

References

- Johansson DJA, Azar C (2007) A scenario based analysis of land competition between food and bioenergy production in the US. *Clim Chang* 82(3–4):267–291. doi:10.1007/s10584-006-9208-1
- Wolf J, Bindraban P, Luijten J, Vleeshouwers L (2003) Exploratory study on the land area required for global food supply and the potential global production of bioenergy. *Agric Syst* 76(3):841–861
- Cibin R, Chaubey I, Engel B (2012) Simulated watershed scale impacts of corn stover removal for biofuel on hydrology and water quality. *Hydrol Process* 26(11):1629–1641
- Raj C (2013) Optimal land use planning on selection and placement of energy crops for sustainable biofuel production. Dissertation, Purdue University
- Thomas M, Engel B, Chaubey I (2009) Water quality impacts of corn production to meet biofuel demands. *J Environ Eng* 135(11):1123–1135
- Thomas M, Engel B, Chaubey I (2011) Multiple corn stover removal rates for cellulosic biofuels and long-term water quality impacts. *J Soil Water Conserv* 66(6):431–444
- Thomas MA, Ahiablame LM, Engel BA, Chaubey I, Mosier N (2014) Modeling water quality impacts of cellulosic biofuel production from corn silage. *Bioenergy Res* 7(2):636–653
- Behman KD, Keitt TH, Kiniry JR (2014) Modeling differential growth in switchgrass cultivars across the Central and Southern Great Plains. *Bioenergy Res*. doi:10.1007/s12155-014-9450-8
- Boles CM (2013) Swat model simulation of bioenergy crop impacts in a tile-drained watershed. Dissertation, Purdue University
- Casler MD (2010) Changes in mean and genetic variance during two cycles of within-family selection in switchgrass. *Bioenergy Res* 3(1):47–54

11. Cortese LM, Honig J, Miller C, Bonos SA (2010) Genetic diversity of twelve switchgrass populations using molecular and morphological markers. *Bioenergy Res* 3(3):262–271
12. Kiniry JR, Anderson L, Johnson M-V, Behrman K, Brakie M, Burner D, Cordsiemon R, Fay P, Fritschi F, Houx J III (2013) Perennial biomass grasses and the Mason–Dixon line: comparative productivity across latitudes in the Southern Great Plains. *Bioenergy Res* 6(1):276–291
13. Schmer MR, Mitchell R, Vogel K, Schacht W, Marx DB (2010) Spatial and temporal effects on switchgrass stands and yield in the Great Plains. *Bioenergy Res* 3(2):159–171
14. Thomas MA (2011) Environmental implications of feedstock production practices for bioenergy. Dissertation, Purdue University
15. Thomas MA, Ahiablame LM, Engel BA, Chaubey I (2014) Modeling water quality impacts of growing corn, switchgrass, and *Miscanthus* on marginal soils. *J Water Resour Prot* 06(14):1352–1368. doi:10.4236/jwarp.2014.614125
16. Trybula EM, Cibin R, Burks JL, Chaubey I, Brouder SM, Volenec JJ (2014) Perennial rhizomatous grasses as bioenergy feedstock in SWAT: parameter development and model improvement. *GCB Bioenergy*. doi:10.1111/gcbb.12210
17. Tilman D, Socolow R, Foley JA, Hill J, Larson E, Lynd L, Pacala S, Reilly J, Searchinger T, Somerville C (2009) Beneficial biofuels—the food, energy, and environment trilemma. *Science* 325(5938):270
18. U. S. Department of Agriculture FS (1980) Energy & wood from intensively cultured plantations: research and development program. Gen. Tech. Rep. St. Paul, MN. North Central Forest Experiment Station, (NC-58), 28. http://www.nrs.fs.fed.us/pubs/gtr/gtr_nc058.pdf. Accessed 1 July 2014
19. Hansen E, Baker J (1979) Biomass and nutrient removal in short rotation intensively cultured plantations. In Proceedings: Impact of intensive harvesting on forest nutrient cycling, ate 1979. State University of New York, College of Environmental Science and Forestry, pp 130–151
20. Hansen EA (1991) Poplar woody biomass yields: a look to the future. *Biomass Bioenergy* 1(1):1–7
21. Hansen EA (1983) Intensive plantation culture: 12 years research. General Technical Report, North Central Forest Experiment Station, USDA Forest Service (NC-91)
22. Strong T, Hansen E (1993) Hybrid poplar spacing/productivity relations in short rotation intensive culture plantations. *Biomass Bioenergy* 4(4):255–261
23. Cannell M, Smith R (1980) Yields of minirotation closely spaced hardwoods in temperate regions: review and appraisal. *For Sci* 26(3):415–428
24. Ek AR (1979) Notes: a model for estimating branch weight and branch leaf weight in biomass studies. *For Sci* 25(2):303–306
25. Isebrands J, Ek A, Meldahl R (1982) Comparison of growth model and harvest yields of short rotation intensively cultured *Populus*: a case study. *Can J For Res* 12(1):58–63
26. Meldahl RS (1979) Yield projection methodology and analysis of hybrid poplars based on multispatial plots. Dissertation, University of Wisconsin, Madison
27. Landsberg J, Wright L (1989) Comparisons among *Populus* clones and intensive culture conditions, using an energy-conservation model. *For Ecol Manag* 27(2):129–147
28. Kiniry J, MacDonald J, Kemanian AR, Watson B, Putz G, Prepas EE (2008) Plant growth simulation for landscape-scale hydrological modelling. *Hydrol Sci J* 53(5):1030–1042
29. MacDonald JD, Kiniry J, Putz G, Prepas E (2008) A multi-species, process based vegetation simulation module to simulate successional forest regrowth after forest disturbance in daily time step hydrological transport models This article is one of a selection of papers published in this Supplement from the Forest Watershed and Riparian Disturbance (FORWARD) Project. *J Environ Eng Sci* 7(S1):127–143
30. Kiniry JR, Johnson M-VV, Bruckerhoff SB, Kaiser JU, Cordsiemon R, Harmel RD (2012) Clash of the titans: comparing productivity via radiation use efficiency for two grass giants of the biofuel field. *Bioenergy Res* 5(1):41–48
31. Kiniry J, Burson B, Evers G, Williams J, Sanchez H, Wade C, Featherston J, Greenwade J (2007) Coastal bermudagrass, bahiagrass, and native range simulation at diverse sites in Texas. *Agron J* 99(2):450–461
32. Kiniry J, Tischler C, Van Esbroeck G (1999) Radiation use efficiency and leaf CO₂ exchange for diverse C₄ grasses. *Biomass Bioenergy* 17(2):95–112
33. Kiniry J (1998) Biomass accumulation and radiation use efficiency of honey mesquite and eastern red cedar. *Biomass Bioenergy* 15(6):467–473
34. Kiniry J, Jones C, O'toole J, Blanchet R, Cabelguenne M, Spanel D (1989) Radiation-use efficiency in biomass accumulation prior to grain-filling for five grain-crop species. *Field Crop Res* 20(1):51–64
35. Cannell M, Sheppard L, Milne R (1988) Light use efficiency and woody biomass production of poplar and willow. *Forestry* 61(2):125–136
36. Arnold J, Kiniry J, Srinivasan R, Williams J, Haney E, Neitsch S (2011) Soil and water assessment tool input/output file documentation version 2009. US Department of Agriculture–Agricultural Research Service, Grassland, Soil and Water Research Laboratory, Temple, TX and Blackland Research and Extension Center, Texas AgriLife Research, Temple, TX Texas Water Resources Institute Technical Report (365)
37. Johnson M-VV, MacDonald JD, Kiniry JR, Arnold J, Gassman P (2009) ALMANAC: a potential tool for simulating agroforestry yields and improving SWAT simulations of agroforestry watersheds. *Int Agric Eng J* 18(1):51
38. Hansen EA, McNeel H, Netzer DA, Phipps HM, Roberts PS, Strong TF, Tolsted D, Zavitskovski J (1979) Short-rotation intensive culture practices for northern Wisconsin. Proceedings, 16th Annual Meeting, North American Poplar Council, Joint Meeting of the United States and the Canadian Chapters, ate., pp 47–63
39. Nelson ND, Michael D (1982) Photosynthesis, leaf conductance, and specific leaf weight in long and short shoots of *Populus* 'Tristis# 1' grown under intensive culture. *For Sci* 28(4):737–744
40. Ek AR, Dawson DH (1976) Actual and projected growth and yields of *Populus* 'Tristis# 1' under intensive culture. *Can J For Res* 6(2):132–144
41. Joslin J, Schoenholtz S (1997) Measuring the environmental effects of converting cropland to short-rotation woody crops: a research approach. *Biomass Bioenergy* 13(4):301–311
42. Tolbert VR, Thornton FC, Joslin JD, Bock BR, Bandaranayake WE, Tyler DD, Pettry D, Green TH, Makik R, Houston AE (1998) Soil and water quality aspects of herbaceous and woody energy crop production: lessons from research-scale comparisons with agricultural crops. Proceedings, BioEnergy, ate., pp 1272–1281
43. Thornton FC, Dev Joslin J, Bock BR, Houston A, Green T, Schoenholtz S, Pettry D, Tyler DD (1998) Environmental effects of growing woody crops on agricultural land: first year effects on erosion, and water quality. *Biomass Bioenergy* 15(1):57–69
44. Kiniry JR, Williams J, Gassman PW, Debaeke P (1992) A general, process-oriented model for two competing plant species. *Trans ASAE (USA)* 35(3):801–810
45. Isebrands J, Nelson N (1983) Distribution of [14C]-labeled photosynthates within intensively cultured *Populus* clones during the establishment year. *Physiol Plant* 59(1):9–18
46. Michael D, Dickmann D, Isebrands J, Nelson N (1990) Photosynthesis patterns during the establishment year within two

- Populus clones with contrasting morphology and phenology. *Tree Physiol* 6(1):11–27
47. Michael D, Isebrands J, Dickmann D, Nelson N (1988) Growth and development during the establishment year of two *Populus* clones with contrasting morphology and phenology. *Tree Physiol* 4(2):139–152
48. Neitsch S, Arnold J, Kiniry J, Williams J (2011) Soil and water assessment tool theoretical documentation version 2009. Institute Texas Water Resources, College Station
49. Black BL, Fuchigami LH, Coleman GD (2002) Partitioning of nitrate assimilation among leaves, stems and roots of poplar. *Tree Physiol* 22(10):717–724
50. McLaughlin RA, Hansen EA, Pope PE (1987) Biomass and nitrogen dynamics in an irrigated hybrid poplar plantation. *For Ecol Manag* 18(3):169–188
51. Zavitkovski J (1981) Characterization of light climate under canopies of intensively-cultured hybrid poplar plantations. *Agric Meteorol* 25:245–255
52. Kumar S, Merwade V (2009) Impact of watershed subdivision and soil data resolution on SWAT model calibration and parameter uncertainty. *JAWRA J Am Water Resour Assoc* 45(5):1179–1196
53. Gupta HV, Sorooshian S, Yapo PO (1999) Status of automatic calibration for hydrologic models: comparison with multilevel expert calibration. *J Hydrol Eng* 4(2):135–143
54. Nash J, Sutcliffe J (1970) River flow forecasting through conceptual models part I—a discussion of principles. *J Hydrol* 10(3):282–290
55. Moriasi D, Arnold J, Van Liew M, Bingner R, Harmel R, Veith T (2007) Model evaluation guidelines for systematic quantification of accuracy in watershed simulations. *Trans ASABE* 50(3):885–900
56. Larose M, Heathman G, Norton L, Engel B (2007) Hydrologic and atrazine simulation of the Cedar Creek watershed using the SWAT model. *J Environ Qual* 36(2):521–531
57. Van Liew M, Arnold J, Garbrecht J (2003) Hydrologic simulation on agricultural watersheds: choosing between two models. *Trans ASAE* 46(6):1539–1551
58. Ek AR, Dawson DH (1976) Yields of intensively grown *Populus*: actual and projected. USDA Forest Service General Technical Report NC
59. Isebrands J, Sturos J, Crist J (1979) Integrated utilization of biomass: a case study of short-rotation intensively cultured *Populus* raw material. Tappi [Technical Association of the Pulp and Paper Industry] (USA)