

Fay, P.A., Polley, H.W., Wilsey, B., Tiner, K., Jin, V.L. 2008. Plant composition and the effects of temperature, soil moisture, and aboveground biomass on soil respiration in grassland. European Geosciences Union General Assembly, April 13-18, 2008, Vienna, Austria. A-09146. CDROM.

Future changes in climate are likely to cause shifts in plant species composition. In grasslands, the growth form composition (i.e., grass/forb) and species diversity of vegetation are important controls on ecosystem carbon cycling because species and growth forms vary in their capacities for carbon uptake, productivity, and allocation belowground. The dominant grasses determine many aspects of ecosystem function, but few studies have examined the relationship of growth form composition and diversity to soil respiration (JCO₂; root + microbial respiration), and how this relationship may influence the response of JCO₂ to soil temperature and moisture. We studied effects of plant growth form and species diversity on JCO₂ by measuring the 2007 annual course of JCO₂, soil temperature (T_{soil}, 0-15 cm), soil moisture (SWC, % volumetric, 0-10 cm), and aboveground biomass in plots of varying composition and diversity. The plots contained a total of 8 C₄ grasses, 1 C₃ grass, and 4 C₃ forbs grown in equal-density monocultures or in mixtures of 2, 4, or 8 species planted in either even or geometric rank abundance distributions. There was strong seasonality in JCO₂ associated with soil temperature (R² = 0.41, p < 0.0001), with less seasonal variation in the geometric than even species distribution plots (p = 0.017). The temperature sensitivity of JCO₂ (the slope of JCO₂ vs. T_{soil}) differed strongly with the composition of the plots (p < 0.0001), with weaker temperature sensitivity in forb monocultures compared to grass monocultures or species mixtures. JCO₂ also correlated with SWC although less so than with T_{soil} (R² = 0.04, p < 0.0001), and the SWC sensitivity of JCO₂ was lower in forb monocultures than in either grass monocultures or mixtures. These results suggest that the growth form composition of vegetation mediates the effects of T_{soil} and SWC on JCO₂. Aboveground biomass did not correlate with JCO₂. However, the ratio of JCO₂ to biomass was 3 to 4-times greater in 2-species mixtures compared to 4 or 8-species mixtures (p = 0.007). This ratio is an index of the net carbon balance in the plots, therefore this result suggests that lower species number is associated with reduced net carbon uptake. Overall, these results indicate that plant growth form composition and diversity may play an important role in soil respiration responses to abiotic drivers and ultimately the net carbon balance of grasslands.