

Complementary epistasis involving *Sr12* explains adult plant resistance to stem rust in Thatcher wheat (*Triticum aestivum* L.)

Matthew N. Rouse · Luther E. Talbert ·
Davinder Singh · Jamie D. Sherman

Received: 31 January 2014 / Accepted: 24 April 2014 / Published online: 17 May 2014
© Springer-Verlag Berlin Heidelberg (outside the USA) 2014

Abstract

Key message Quantitative trait loci conferring adult plant resistance to Ug99 stem rust in Thatcher wheat display complementary gene action suggesting multiple quantitative trait loci are needed for effective resistance.

Abstract Adult plant resistance (APR) in wheat (*Triticum aestivum* L.) to stem rust, caused by *Puccinia graminis* f. sp. *tritici* (*Pgt*), is desirable because this resistance can be *Pgt* race non-specific. Resistance derived from cultivar Thatcher can confer high levels of APR to the virulent *Pgt* race TTKSK (Ug99) when combined with stem rust resistance gene *Sr57* (*Lr34*). To identify the loci conferring APR in Thatcher, we evaluated 160 RILs derived from Thatcher crossed to susceptible cultivar McNeal for field stem rust

reaction in Kenya for two seasons and in St. Paul for one season. All RILs and parents were susceptible as seedlings to race TTKSK. However, adult plant stem rust severities in Kenya varied from 5 to 80 %. Composite interval mapping identified four quantitative trait loci (QTL). Three QTL were inherited from Thatcher and one, *Sr57*, was inherited from McNeal. The markers closest to the QTL peaks were used in an ANOVA to determine the additive and epistatic effects. A QTL on 3BS was detected in all three environments and explained 27–35 % of the variation. The peak of this QTL was at the same location as the *Sr12* seedling resistance gene effective to race SCCSC. Epistatic interactions were significant between *Sr12* and QTL on chromosome arms 1AL and 2BS. Though *Sr12* cosegregated with the largest effect QTL, lines with *Sr12* were not always resistant. The data suggest that *Sr12* or a linked gene, though not effective to race TTKSK alone, confers APR when combined with other resistance loci.

Communicated by A. Zhang.

M. N. Rouse (✉)
Cereal Disease Laboratory, United States Department
of Agriculture-Agricultural Research Service (USDA-ARS),
1551 Lindig Street, St. Paul, MN 55108, USA
e-mail: matthew.rouse@ars.usda.gov

M. N. Rouse
Department of Plant Pathology, University of Minnesota,
485 Borlaug Hall, St. Paul, MN 55108, USA

L. E. Talbert · J. D. Sherman
Department of Plant Sciences and Plant Pathology, Montana
State University, P.O. Box 173150, Bozeman, MT 59717, USA

D. Singh
International Maize and Wheat Improvement Center (CIMMYT),
Mexico, DF, Mexico

Present Address:

D. Singh
Plant Breeding Institute, University of Sydney, Private Bag 4011,
Narellan, NSW 2567, Australia

Introduction

The stem rust fungus *Puccinia graminis* Pers.:Pers. f. sp. *tritici* Eriks. & E. Henn., *Pgt*, has caused severe yield losses in bread wheat (*Triticum aestivum* L.), durum wheat [*T. turgidum* subsp. *durum* (Desf.) Husnot], barley (*Hordeum vulgare* L.), and triticale (*X Triticosecale* Wittmack) (Roelfs et al. 1992). Wheat stem rust was historically problematic throughout wheat-producing areas of the world. In the United States, yield losses exceeded 25 % of state production during epidemic years in Minnesota and the Dakotas (Roelfs 1978). Cultivar Thatcher was bred specifically for stem rust resistance and released in 1935 (Hayes et al. 1936). Though races emerged in the 1950s with virulence to the seedling genes in Thatcher, the adult plant resistance

remained moderately effective. Removal of the widespread alternate host of the stem rust pathogen (*Berberis vulgaris* L.) combined with breeding resistant cultivars with multiple stem rust resistance genes, including resistance derived from Thatcher, has prevented stem rust epidemics since 1956 (Kolmer et al. 1991).

The detection of race TTKSK of the stem rust pathogen in Uganda in 1998 (commonly referred to as Ug99) and the subsequent spread of the disease throughout East and Southern Africa and the Middle East has caused concern over the current susceptibility of wheat to Ug99 (Nazari et al. 2009; Pretorius et al. 2000, 2010; Singh et al. 2011a; Wanyera et al. 2006). Since the original characterization of Ug99 as race TTKSK, variants of the pathogen have been detected with virulence to additional resistance genes increasing the danger of the Ug99 race group and reminding wheat breeders of the short longevity of single major resistance genes (Jin et al. 2007, 2008, 2009; Pretorius et al. 2012).

To mitigate the threat of Ug99, wheat accessions were screened to identify sources of resistance (Jin and Singh 2006; Jin et al. 2007; Njau et al. 2010; Rouse et al. 2011). At least 18 Ug99-effective resistance genes were identified and mapped from diploid or tetraploid wheat relatives (Saal and Wricke 1999; Mago et al. 2002, 2005; Faris et al. 2008; Sambasivam et al. 2008; Tsilo et al. 2008; Wu et al. 2009; Liu et al. 2010; Olson et al. 2010; Zhang et al. 2010; Liu et al. 2011a, 2011b; Niu et al. 2011; Qi et al. 2011; Simons et al. 2011; Ghazvini et al. 2012a; Klindworth et al. 2012), and at least four Ug99-effective resistance genes have been identified and mapped from bread wheat (Hiebert et al. 2010, 2011; Rouse et al. 2012; Ghazvini et al. 2012b). The availability of multiple Ug99-effective stem rust resistance genes and linked molecular markers provides the opportunity to combine multiple genes into cultivars to reduce the risk of the development of pathogen virulence to these genes. In addition, wheat cultivars have been characterized as resistant to Ug99 at the adult plant stage, but susceptible at the seedling stage (Njau et al. 2010). Such cultivars have been characterized to possess not single major-effect resistance genes, but multiple minor-effect resistance genes effective only at the adult plant stage (Singh et al. 2011b). The genes in these varieties include *Sr2*, *Sr57* (*Lr34*), and other uncharacterized genes, many of which are thought to originate from cultivar Thatcher (Singh et al. 2011b). Molecular markers for wheat stem rust adult plant resistance (APR) have been developed only for *Sr2* (Mago et al. 2011) and *Sr57/Lr34* (Lagudah et al. 2006).

Since APR cannot be assessed in efficient and highly replicable wheat seedling assays and the effects of APR genes are minor, mapping stem rust APR must be assisted by genome-wide molecular markers. Three association analyses detected 12, 15, and three marker loci significantly

associated with resistance to Ug99 in spring wheat, winter wheat, and durum wheat, respectively (Yu et al. 2011, 2012; Singh et al. 2013b). QTL were identified linked to Ug99 APR on seven chromosomes from a biparental spring wheat population (Singh et al. 2013c). A total of nine stem rust resistance loci were identified in a durum wheat biparental population (Haile et al. 2012). Though the majority of the variation in Ug99 stem rust response was explained by the additive effects of APR loci, several of the loci detected in these five studies exhibited significant and complex epistatic interactions. Tests for epistatic interactions were not conducted in biparental populations of spring bread wheat where four and five APR quantitative trait loci (QTL) were identified (Njau et al. 2013; Singh et al. 2013c) or in an association analysis of durum wheat germplasm where 36 loci associated with Ug99 stem rust resistance were identified (Letta et al. 2013).

Since stem rust resistance derived from cultivar Thatcher has been used in diverse wheat varieties since its release (Hayes et al. 1936), previous researchers have made progress in dissecting the components of stem rust resistance in Thatcher. At least four seedling resistance genes (*Sr5*, *Sr9g*, *Sr12*, and *Sr16*) are present in Thatcher (Luig 1983) and these genes explain some but not all of the resistance exhibited at the adult plant stage, suggesting the involvement of additional adult plant resistance genes (Green and Dyck 1975; Knott 2000, 2001; Gavin Vanegas et al. 2008). APR gene *Lr34* was previously demonstrated to enhance Thatcher APR (Gavin Vanegas et al. 2008; Kolmer et al. 2011). In a biparental population segregating for both Thatcher resistance and *Lr34*, a QTL on chromosome arm 2BL exhibited weak effectiveness to Ug99, but was enhanced by *Lr34* (Kolmer et al. 2011). The QTL on chromosome arm 2BL mapped to a location consistent with *Sr9* indicating that the APR could involve *Sr9g* (Tsilo et al. 2007). These studies have determined that the adult plant resistance in Thatcher is complex, difficult to genetically map, and related, but not completely explained by seedling resistance genes in Thatcher.

The availability of a biparental population derived from Thatcher and susceptible cultivar McNeal (Lanning et al. 1994) with genotypic data for 615 polymorphic molecular markers (Sherman et al. 2014), allowed for a more complete characterization of the APR in Thatcher. Characterizing Thatcher APR is relevant to current wheat breeding as varieties with APR presumably derived from Thatcher such as Kingbird and Tom are used by current breeding programs (Singh et al. 2011b; Anderson et al. 2012). Identification of markers linked to Thatcher APR will facilitate the selection of APR in available breeding germplasm without introgression of resistance from unadapted germplasm, which would likely possess linkage drag. Our objectives were to (1) identify stem rust adult plant resistance loci in

the Thatcher/McNeal population, (2) determine the additive and epistatic effects of APR loci involved, and (3) determine if the APR loci correspond to seedling resistance genes in Thatcher.

Materials and methods

Plant materials

Cultivars McNeal (USDA National Small Grains Collection accession Number GSTR 10201) and Thatcher (GSTR 10202) and a population of 160 recombinant inbred lines (RILs) derived from single F_5 plants (GSTR 10203 through GSTR 10362) were used in previous studies to identify QTL for agronomic traits (Naruoka et al. 2011; Sherman et al. 2014). We used this population and parents to characterize stem rust resistance derived from Thatcher. Cultivar Prelude (CI 4323) and genetic stock PdSr12tc (*Sr12* backcrossed into Prelude from Thatcher) were obtained from the United States Department of Agriculture Agricultural Research Service (USDA-ARS) Cereal Disease Laboratory and used as *Sr12* negative and positive controls, respectively. *Sr16* isogenic wheat lines ISr16-Ra and CnsTc2B/W3498 were obtained from the USDA-ARS Cereal Disease Laboratory and used as positive controls for *Sr16*.

Stem rust phenotyping

Stem rust seedling assays were performed at the USDA-ARS Cereal Disease Laboratory in St. Paul, Minnesota. Thatcher, McNeal, Prelude, PdSr12tc, and all RILs were evaluated with *P. graminis* f. sp. *tritici* races SCCSC (isolate 09ID73-2) and TTKSK (isolate 04KEN156/05; Ug99). Race names were derived according to internationally recognized race nomenclature (Roelfs and Martens 1988; Jin et al. 2008). Evaluation with race TTKSK was performed in a biocontainment safety level 3 greenhouse. Five seeds of each line were planted in trays filled with vermiculite and inoculated after full emergence of the primary leaf ~8 days after planting. Urediniospores of *P. graminis* f. sp. *tritici* were inoculated as described previously (Rouse et al. 2011). After dew chamber incubation, the plants were then placed in greenhouses maintained at 22 ± 2 °C during the day and 18 ± 2 °C during the night with supplemental lighting in the morning and evening to provide for a photoperiod of 16 h. A total of 14 days after inoculation, stem rust was assessed on the primary leaves by recording infection types (ITs) on the '0' to '4' scale as described by Stakman et al. (1962). The sums of lines in discrete infection type classes were tested against expected segregation ratios by Chi-square (χ^2) goodness-of-fit tests.

The population was tested for response to stem rust in three field seasons. During 2008 and 2009, one replication of the population and parents were evaluated in Njoro, Kenya at the Kenya Agricultural Research Institute, Plant Breeding Station. In 2011, one replication of the population and parents were evaluated in St. Paul, Minnesota, USA. Planting, plot maintenance, and *P. graminis* f. sp. *tritici* inoculation in both Kenya and St. Paul were performed as described in Rouse et al. (2011). In Kenya, bulked urediniospores collected from Kenyan wheat varieties were used to inoculate susceptible spreader plants planted in the experimental plots. Subsequent race analysis of collections made from these plots identified race TTKST (Ug99 variant with virulence to *Sr24*) as the only race present (Y. Jin and M.N. Rouse, unpublished). In St. Paul, the susceptible spreader rows were inoculated with equal amounts of North American *P. graminis* f. sp. *tritici* races TPMKC, RKQQC, RCRSC, QTHJC, QFCSC, and MCCFC (Rouse et al. 2011). Disease evaluations were performed when the experimental entries reached the soft dough growth stage. Disease response was characterized by recording severity (0–100 %) according to the modified Cobb scale (Peterson et al. 1948) and infection response according to Roelfs et al. (1992). Infection response classifications discriminated among size and shape of uredia in addition to the amount of chlorosis in plant tissue adjacent to uredia. Infection responses observed in this study included moderately susceptible (MS) and susceptible (S).

According to the *Pgt* race nomenclature that is based on the response of *Pgt* isolates to wheat stem rust resistance genes (Roelfs and Martens 1988; Jin et al. 2008), the races of *Pgt* used in this study including SCCSC, TTKSK, TTKST, TPMKC, RKQQC, RCRSC, QTHJC, QFCSC, and MCCFC are virulent to resistance genes *Sr5* and *Sr9g*. Gene *Sr16* is not present in the race nomenclature panel of resistance genes. To evaluate the effectiveness of *Sr16* to these races we inoculated *Sr16* isogenic lines ISr16-Ra and CnsTc2B/W3498 at the seedling stage with the *Pgt* races used in this study.

We evaluated Thatcher, McNeal, Prelude, PdSr12tc for response to stem rust in three treatments in 2013: evaluation with race TTKST in Njoro, evaluation with race TPMKC in a single race nursery in Rosemount, Minnesota, and evaluation with race RCRSC in a separate single race nursery in Rosemount. At least two replicates of each line were included in the three treatments.

Genetic map

A total of 615 markers were obtained from a previous study utilizing the same RIL population (Sherman et al. 2014). These markers included 334 DArT markers, 274 SSR markers, seven additional markers linked to storage protein,

hardness, and height traits. Presence and absence of two morphological traits in the RIL population, awnedness and glume color, were also available from Sherman et al. (2014). We genotyped DNAs previously extracted from the RIL population with marker *Xcssfr6* according to the protocol used by Lagudah et al. (2009). This marker was previously developed to determine the presence or absence of gene *Lr34* that has been identified as a putative ABC transporter (Krattinger et al. 2009). The same putative ABC transporter gene was also designated as stem rust resistance gene *Sr57* by the catalogue of gene symbols for wheat managed by the International Wheat Genetics Symposium (McIntosh et al. 2012). The presence or absence of stem rust resistance gene *Sr12* in the RILs was determined by analysis of the seedling IT data derived from screening with *Pgt* race SCCSC (see “Results”). Data on presence or absence of *Sr57* and *Sr12* were included with the data derived from the 615 molecular markers and two morphological traits for construction of a genome-wide genetic map using MapDisto (Lorieux et al. 1995). The Place Locus function (LOD 2.3 minimum and rmax of 0.3) was used to add the *Sr57* and *Sr12* traits to the map created by Sherman et al. (2014).

QTL analysis

Stem rust severities for each RIL were used to identify QTL for each environment. Composite interval mapping (Zeng 1993, 1994) was performed using QTL Cartographer (Wang et al. 2007). LOD values were set by 1,000 permutations at an experiment-wise $p < 0.01$ using a window size of ten map units. RILs were classified as possessing the Thatcher or McNeal allele of each significant QTL if they possessed the same parental allele at the peak marker of the QTL. The additive and two-way epistatic effects contributed by the significant QTL were determined by analysis of variance (R Development Core Team 2010), using the data on allelic state for each significant QTL. These effects were calculated for each environment independently and also for the combined data. The size of the RIL population prevented effective analysis of three-way interactions.

Results

Stem rust phenotypic analyses and segregation of resistance

Thatcher and *Sr12*-positive control PdSr12tc displayed seedling ITs ‘0’ and ‘0;’, respectively, to *Sr12*-avirulent *Pgt* race SCCSC. McNeal and Prelude displayed ITs ‘2+3-’ and ‘3+’, respectively, which were not characteristics of *Sr12*. The RILs segregated for *Sr12*-positive ITs ranging from ‘0’ to ‘0;’ and *Sr12*-negative ITs ranging from ‘22+’

to ‘3’. All *Pgt* races displayed ITs of ‘3+’ to ‘4’ on both *Sr16* isogenic lines except race SCCSC that displayed IT ‘22+’ to the *Sr16* isogenic lines. Since race SCCSC is avirulent to *Sr16* and the low IT for *Sr16* is ‘22+’ in response to race SCCSC, we were able to discriminate among plants with *Sr12* that had low ITs of ‘0’ to ‘0;’ and plants that did not possess *Sr12* with ITs of ‘22+’ or greater. A total of six RILs were heterogeneous for *Pgt* race SCCSC seedling IT. Of the remaining 154 RILs, 69 were classified as *Sr12*-positive and 85 were classified as *Sr12*-negative. The segregation of *Sr12* did not deviate from a 1:1 ratio expected for a single gene ($\chi^2 = 1.66$, $p = 0.20$). McNeal, Prelude, and PdSr12tc all displayed susceptible ITs of ‘3+’ to *Pgt* race TTKSK. All RILs displayed susceptible ITs of ‘33+’ to ‘3+’ to race TTKSK. Infrequently, single plants within the five seedlings for each line in the seedling assay displayed a slightly mesothetic reaction to race TTKSK such as IT ‘33+;’. Similarly, Thatcher was observed to display IT ‘3+’ and infrequently a mesothetic ‘33+;’ infection type to *Pgt* race TTKSK as recorded in Rouse et al. (2012). Since the mesothetic reaction was inconsistent among plants within a given line and difficult to distinguish from a susceptible IT, classification of lines with and without mesothetic seedling resistance was not attempted.

Thatcher displayed intermediate severities and moderately susceptible to susceptible infection responses in the field tests: Kenya 2008, 40-50 MS-S; Kenya 2009, 40 S-MS; and St. Paul 2011, 30-55 MS. McNeal displayed high severities and a susceptible infection response: Kenya 2008, 80 S; Kenya 2009, 50 S; and St. Paul 2011, 70 S. The RILs segregated for both traits: Kenya 2008, 5-80 MS-S; Kenya 2009, 5-80 MS-S; and St. Paul, 15-80 MS-S. The severity values of the RILs among environments were significantly and positively correlated (Pearson’s r ; $p < 0.001$) in all three pairwise comparisons of environments: Kenya 2008 and Kenya 2009 ($r = 0.57$); Kenya 2008 and St. Paul 2011 ($r = 0.67$); and Kenya 2009 and St. Paul 2011 ($r = 0.63$).

In 2013 Thatcher, McNeal, PdSr12tc, and Prelude displayed moderately susceptible to susceptible infection responses and average severities of 15.625, 35, 40, and 38.75, respectively, in response to race TTKST; 53.75, 63.75, 67.5, and 62.5, respectively, in response to race TPMKC; and 59.375, 68.125, 70, and 67.5, respectively, in response to race RCRSC. In all three treatments, Thatcher displayed lower severities compared to the other three wheat lines including PdSr12tc. *Sr12* did not reduce stem rust severity in PdSr12tc compared to Prelude.

QTL analysis

A total of four stem rust resistance QTL were identified, two of which were significant in all three environments

(Fig. 1; Table 1). The QTL with the largest effect in each environment coincided with *Sr12* seedling resistance to *Pgt* race SCCSC. We designated this QTL as *QSr.cdl-3BS*. Thatcher contributed the resistant allele of *QSr.cdl-3BS*. Since *Sr12* was the peak locus at this QTL, we used presence and absence of the *Sr12* resistance gene as indicative of the allelic state of *QSr.cdl-3BS*. A QTL on chromosome arm 1AL was observed in three environments we designated as *QSr.cdl-1AL*. Thatcher contributed the resistant allele of *QSr.cdl-1AL*. The peak marker of *QSr.cdl-1AL* was *XwPt6869*. A QTL was observed in the 2008 and 2009 Kenya experiments that coincided with the *Xcssfr6* marker for *Sr57*. McNeal contributed *Sr57*. A QTL was observed in the 2008 Kenya field experiment on chromosome arm 2BS. Thatcher contributed the resistant allele of this QTL that we designated as *QSr.cdl-2BS*. The peak marker of *QSr.cdl-2BS* was *Xcfd238*. Marker *Xcfd238* was significantly

associated with stem rust resistance in both the 2008 Kenya and 2011 St. Paul field experiments (Table 1).

Several of the two-way epistatic effects among the four QTL were significant (Table 1). The interaction between *Sr12* and *QSr.cdl-1AL* was significant in all three environments and explained a greater proportion of the phenotypic variation than the additive effect of *QSr.cdl-2BS* in each environment. The interaction between *Sr12* and *QSr.cdl-2BS* was also significant in all three environments, but explained a relatively small amount of the variation. The two-way epistatic interactions between (1) *Sr12* and *Sr57* and (2) *QSr.cdl-1AL* and *QSr.cdl-2BS* were significant only in the 2009 Kenya environment. In each of the significant two-way epistatic interactions, the effect of possessing resistant alleles from both loci resulted in greater resistance than was expected based upon the reduction of disease conferred by single resistant alleles (Fig. 2). This pattern was




Fig. 1 Linkage maps of genomic regions significantly associated with stem rust severity in three environments and corresponding plots of the level of significance. The three environments are represented by a dotted line (Kenya 2008), a dotted-dashed line (Kenya 2009),

and a dashed line (St. Paul 2011) on the plots to the right of the linkage maps. Linkage maps are from Sherman et al. (2014) with the addition of *Sr12* and *Sr57*

validated by dissecting the mean severities of RILs classified by their allelic state at each of the four QTL (Fig. 3). RILs with a single resistance allele from any of the four QTL did not display decreased stem rust severity. Even the QTL with the greatest additive effect, *QSr.cdl-3BS* (*Sr12*) that explained at least 26.67 % of the variation in severity did not decrease stem rust severity when present alone. With the exception of the category where resistant alleles

Table 1 Percentage of the variation in stem rust disease severity from three field experiments explained by the effects of four QTL and their two-way interactions from ANOVA

Effect	Kenya 2008	Kenya 2009	St. Paul 2011	Combined
<i>Sr12</i> ^a	26.67***	28.92***	34.62***	38.86***
1AL ^b	4.23***	10.54***	5.49***	8.73***
<i>Sr57</i>	9.49***	6.96***	1.25	6.79***
2BS ^c	5.00***	1.38	3.07**	4.12**
<i>Sr12</i> -1AL	7.92***	4.21**	3.69**	7.52***
<i>Sr12</i> - <i>Sr57</i>	0.92	6.09***	0.09	1.54*
<i>Sr12</i> -2BS	1.53*	5.25**	2.32*	4.69***
1AL- <i>Sr57</i>	0.34	0.16	0.66	0.16
1AL-2BS	0.09	2.23*	0.42	1.37
<i>Sr57</i> -2BS	0.5	1.2	0.02	1.22

^a *Sr12* was used to indicate *QSr.cdl-3BS*

^b Peak marker *XwPt6869* was used to indicate the QTL observed on chromosome arm 1AL named *QSr.cdl-1AL*

^c Peak marker *Xcfd238* was used to indicate the QTL observed on chromosome arm 2BS named *QSr.cdl-2BS*

* $p < 0.05$

** $p < 0.01$

*** $p < 0.001$

are present at the four QTL, only four categories of allelic states were observed with significantly decreased severities (Fig. 3). Three of these four categories included all possible combinations possessing resistant alleles from both *Sr12* and *QSr.cdl-1AL*. The fourth category was the combination of resistant alleles from *Sr12*, *Sr57*, and *QSr.cdl-2BS*. The data indicate that, at a minimum, the combination of resistant alleles from (1) *Sr12* and *QSr.cdl-1AL* or (2) *Sr12*, *Sr57*, and *QSr.cdl-2BS* are necessary for any detectable reduction in stem rust severity in the Thatcher/McNeal population.

We tested for two-way epistatic interactions between the four peak markers and all other markers using ANOVA. Eight interactions were detected with p values < 0.0001 , however, these interactions were between markers linked to *Sr12* (*QSr.cdl-3BS*) and markers linked to *XwPt6869* (*QSr.cdl-1AL*). We did not identify epistatic interactions between the four QTL and any other marker not linked to the four QTL that explained > 2 % of the variation in stem rust severity from the combined environments.

Discussion

There are four results from this study that have implications for plant disease resistance breeding and biology. The first result is the observation that complementary epistasis is responsible for adult plant resistance to stem rust in wheat. Epistasis, the interaction of alleles at different loci, is often conferred by duplicate gene action (Snape and Riggs 1975) in plant disease resistance studies. For example, Lowe et al. (2011) identified four stripe rust resistance QTL in a biparental population, and presence of the QTL with the largest




Fig. 2 Allelic interaction plots calculated as combined environment stem rust severity means of RILs of the Thatcher/McNeal population by allelic combination at pairs of loci: **a** *Sr12* and *QSr.cdl-1AL* (represented as ‘1AL’), **b** *Sr12* and *QSr.cdl-2BS* (represented as ‘2BS’), and **c** *Sr12*

and *Sr57*. Error bars were calculated as plus or minus twice the standard error. Alleles from Thatcher are designated as ‘A’ (red font), whereas McNeal alleles are designated as ‘B’ (blue font). p values describe the significance of the two-way interactions calculated by ANOVA




Fig. 3 Combined environment stem rust severity means of RILs of the Thatcher/McNeal population by allelic combination at four loci: *Sr12*, *QSr.cdl-1AL* (represented as ‘1AL’), *QSr.cdl-2BS* (represented as ‘2BS’), and *Sr57*. Error bars were calculated as plus or minus twice the standard error. Significant differences among allelic combinations were calculated by Tukey’s Honestly Significant Difference test

effect alone, *QYr.ucw-3BS*, was sufficient for adequate reduction in stripe rust severity. The effects of the other QTL were masked in the presence of *QYr.ucw-3BS*. The resulting ANOVA indicated significant epistatic interaction among five of the six possible two-way comparisons among the QTL because of masking effects. Similarly, the significant epistatic interactions reported by Singh et al. (2013a) were duplicate interactions where one major gene, likely *SrCad*, was sufficient to substantially reduce stem rust disease. Though Haile et al. (2012) reported significant epistatic interactions, the type of epistasis was not reported. Since the population was skewed with a large proportion of lines reported as resistant, we expect that the epistasis was duplicate similar to epistasis in Lowe et al. (2011) and Singh et al. (2013a) where populations were similarly skewed. The association mapping studies conducted by Yu et al. (2011, 2012) reported substantial epistasis among significant QTL identified and among additional loci not significantly associated with stem rust resistance alone. Yu et al. (2012) reported multiple types of epistasis, though these were not disclosed. We report complementary epistasis as the form of gene action responsible for the complex resistance to stem rust in Thatcher wheat. Though our ANOVA indicated that more of the variation was explained

by the additive effects of the significant QTL than the epistatic effects (Table 1), comparing means of categories of RILs indicated that presence of any single significant QTL was not sufficient to reduce stem rust severity. This result (1) warns plant breeders of the ability of complementary gene action to overestimate the effects of single QTL, (2) suggests that the effect of single QTL should be assessed without the presence of other resistance loci, and (3) suggests that combining multiple QTL may provide greater than expected disease resistance because of complementary epistasis.

A second result from this study is the observation that a defeated resistance gene, namely *Sr12*, coincides with APR to a virulent pathogen race and that this resistance is only effective in the presence of additional APR genes. Resistance gene *Sr12* alone (PdSr12tc) was not sufficient to reduce stem rust severity. Gene *Sr12* was previously reported to correlate with field adult plant resistance in populations fixed or segregating for *Sr7a* in response to *Pgt* races that were avirulent to *Sr7a* at the seedling stage (Singh and McIntosh 1987). Our data suggest that *Sr12* can complement other resistance loci in a population devoid of seedling resistance to the pathogen race used in the field (McNeal and Thatcher were both susceptible as seedlings to race TTKSK). The observation of QTL coinciding with defeated major resistance genes has been reported in the rice blast and potato late blight pathosystems (Young 1996). It is possible that the observed QTL and coinciding defeated resistance gene in our study and in other systems are conferred by tightly linked independent loci. We did not observe field resistance to be associated with the chromosome arms where Thatcher seedling resistance genes *Sr5*, *Sr9g*, and *Sr16* are located. The identification of qualitative resistance genes that are associated with APR even when they are defeated, such as *Sr12*, suggests that at least some qualitative resistance genes may be useful in breeding programs even after they are defeated. Additional data are needed to validate whether or not the *Sr12* gene confers the associated APR to virulent *Pgt* races.

The third informative result from this study is the description of QTL conferring APR to Ug99 and linked markers. Validation and optimization of the markers linked to stem rust resistance QTL described in this study are needed for effective marker-assisted selection in wheat breeding. The significant association of Thatcher alleles at *QSr.cdl-3BS*, *QSr.cdl-1AL*, and *QSr.cdl-2BS* with stem rust resistance in both Kenya and St. Paul suggests that these QTL are effective to multiple races of *Pgt*. We did not identify a QTL on chromosome arm 2BL to be associated with APR as in Kolmer et al. (2011). Since Thatcher appears twice in the pedigree of McNeal, it is possible that the population is fixed at this locus. Alternatively, the use of different parents in Kolmer et al. (2011) may explain the

discrepancy in identification of this QTL. Below, we have written a discussion of each of the four QTL identified in this study.

1. *QScd1-3BS*: Our description of *Pgt* race SCCSC resistance (*Sr12*) as coinciding with *QScd1-3BS* allows for screening of germplasm with *Sr12*-avirulent races to identify *Sr12*-associated APR to virulent races such as TTKSK (Ug99). In addition, we described markers linked to *Sr12*: *XwPt6047* and *XwPt0544* (Fig. 1) that could be used to develop PCR-based markers for this locus for marker-assisted selection. Markers potentially near the *Sr12* locus were identified as significantly correlated with APR to Ug99 in CIMMYT (International Maize and Wheat Improvement Center) spring wheat germplasm by Yu et al. (2011). In addition, Ug99 resistance QTL overlapping the *Sr12* locus were identified in biparental populations derived from CIMMYT germplasm (Yu et al. 2014). As many CIMMYT spring wheats have Thatcher in their pedigree, it is possible that *Sr12* is present in CIMMYT germplasm. Yu et al. (2011) also reported epistatic interactions between markers possibly near the *Sr12* locus and markers on chromosome 2B, providing additional evidence that Thatcher resistance is present in CIMMYT germplasm.
2. *QScd1-IAL*: Markers *XwPt6869* and *XwPt0827* could be used to select for *QScd1-IAL* that must be complemented by *Sr12* to confer resistance to Ug99. Interestingly, marker *XwPt6869* was identified to significantly associate with epistatic interactions for leaf rust resistance in an association analysis in durum wheat (Singh et al. 2013b). Since Thatcher possesses Iumillo durum as a parent and *QScd1-IAL* was observed to have epistatic interactions, *QScd1-IAL* and the resistance associated with *XwPt6869* observed by Singh et al. (2013b) may be conferred by the same gene.
3. *Sr57*: We confirmed that *Sr57* confers field resistance to Ug99 as reported by previous studies (Kolmer et al. 2011; Yu et al. 2011, 2012). Similar to several of the previous studies, we found that *Sr57* alone was not sufficient to decrease stem rust disease severity. We did not observe an effect of *Sr57* at the St. Paul 2011 test.
4. *QScd1-2BS*: The QTL on chromosome 2BS displayed relatively weak effects and interacted significantly with *Sr12* in all environments. Similarly, a region of 2BS was identified in an association mapping study of CIMMYT winter wheat that also exhibited epistatic interactions (Yu et al. 2012). Marker *Xcfd238* could be used to select for *QScd1-2BS*.

The fourth result is the identification of germplasm with desirable APR to wheat stem rust and Ug99 in particular. Though Thatcher possesses intermediate levels of field

resistance to stem rust, transgressive segregants combining resistance from Thatcher and *Sr57* from McNeal were identified. We recommend the use of GSTR 10285 and GSTR 10299 (available from the USDA-ARS National Small Grains Collection) as stem rust resistant donor lines for breeding programs. Both lines have the reduced height allele at *Rht-D1* (Sherman et al. 2014) in addition to very high levels of stem rust resistance. We identified several lines with *Sr12* but without *Sr57*, *QScd1-IAL*, and *QScd1-2BS*. Since these lines did not display reduced levels of stem rust compared to lines without any resistance alleles (Fig. 3), we identified these lines based on seedling resistance to race SCCSC and marker haplotypes at the QTL. These lines include GSTR 10244, GSTR 10256, GSTR 10320, and GSTR 10328 and could be used in further studies to address the effect of *Sr12* without complementation of additional resistance alleles. Additional studies are needed to validate the presence and effectiveness of the Thatcher-derived resistance alleles in conventional germplasm. Correspondence between our study and the association mapping studies described previously suggest that Thatcher resistance may indeed be present in conventional CIMMYT germplasm.

Overall, our data suggest that APR in Thatcher wheat is conferred by at least three loci including the *Sr12* locus. The difficulties in characterizing Thatcher APR in the past were likely due to the complementary epistasis of QTL. Our data suggest that APR to stem rust in wheat may involve complex gene interactions. Knowledge of the complementary nature of these interactions suggests that multiple APR loci are needed to breed for disease resistance. The companion paper summarizing the Ug99 resistance loci identified in 21 studies in a consensus map (Yu et al. 2014) will be a useful resource for wheat breeders to select multiple stem rust resistance loci.

Acknowledgments Funding for this research was provided by the USDA-ARS National Plant Disease Recovery System, the Durable Rust Resistance in Wheat project (funded by the Bill and Melinda Gates Foundation and the United Kingdom Department for International Development), and the USDA-AFRI Triticeae Coordinated Agricultural Project (2011-68002-30029). We acknowledge the Computational Genetics Laboratory of the University of Minnesota Supercomputing Institute for computational support. We are grateful to staff at the Kenya Agricultural Research Institute and at the USDA-ARS Cereal Disease Laboratory for maintenance of field experiments. Mention of trademark, proprietary product, or vendor does not constitute a guarantee or warranty of the product by the USDA, and does not imply its approval to the exclusion of other products and vendors that might also be suitable.

Conflict of interest The authors declare that there are no conflict of interest.

Ethical standards The authors declare that the experiments comply with the current laws of the countries in which the experiments were performed.

References

- Anderson JA, Wiersma JJ, Linkert GL, Kolmer JA, Jin Y, Dill-Macky R, Wiersma JV, Hareland GA, Busch RH (2012) Registration of ‘Tom’ wheat. *J Plant Reg* 6:180–185
- Faris JD, Xu SS, Cai X, Friesen TL, Jin Y (2008) Molecular and cytogenetic characterization of a durum wheat-*Aegilops speltoides* chromosome translocation conferring resistance to stem rust. *Chromosome Res* 16:1097–1105
- Gavin Vanegas CD, Garvin DF, Kolmer JA (2008) Genetics of stem rust resistance in the spring wheat cultivar Thatcher and the enhancement of resistance by *Lr34*. *Euphytica* 159:391–401
- Ghazvini H, Hiebert CW, Zegeye T, Fetch T (2012a) Inheritance of stem rust resistance derived from *Aegilops triuncialis* in wheat line Tr129. *Can J Plant Sci* 92:1037–1041
- Ghazvini H, Hiebert CW, Zegeye T, Liu S, Dilawari M, Tsilo T, Anderson JA, Rouse MN, Jin Y, Fetch T (2012b) Inheritance of resistance to Ug99 stem rust in wheat cultivar Norin 40 and genetic mapping of *Sr42*. *Theor Appl Genet* 125:817–824
- Green GJ, Dyck PL (1975) The reaction of Thatcher wheat to Canadian races of stem rust. *Can Plant Dis Surv* 55:85–86
- Haile JK, Nachit MM, Hammer K, Badebo A, Roder MS (2012) QTL mapping of resistance to race Ug99 of *Puccinia graminis* f. sp. *tritici* in durum wheat (*Triticum durum* Desf.). *Mol Breeding* 30:1479–1493
- Hayes HK, Ausemus ER, Stakman EC, Bailey CH, Wilson HK, Bamberg RH, Markley MC, Crim RF, Levine MN (1936) Thatcher wheat. *Minn Agric Exp Stn Tech Bull* 325
- Hiebert CW, Fetch T Jr, Zegeye T (2010) Genetics and mapping of stem rust resistance to Ug99 in the wheat cultivar Webster. *Theor Appl Genet* 121:65–69
- Hiebert CW, Fetch TG, Zegeye T, Thomas JB, Somers DJ, Humphreys DG, McCallum BD, Cloutier S, Singh D, Knott DR (2011) Genetics and mapping of seedling resistance to Ug99 stem rust in Canadian wheat cultivars ‘Peace’ and ‘AC Cadillac’. *Theor Appl Genet* 122:143–149
- Jin Y, Singh RP (2006) Resistance in US wheat to recent Eastern African isolates of *Puccinia graminis* f. sp. *tritici* with virulence to resistance gene *Sr31*. *Plant Dis* 90:476–480
- Jin Y, Singh RP, Ward RW, Wanyera R, Kinyua M, Njau P, Fetch T Jr, Pretorius ZA, Yahyaoui A (2007) Characterization of seedling infection types and adult plant infection responses of monogenic *Sr* gene lines to race TTKS of *Puccinia graminis* f. sp. *tritici*. *Plant Dis* 91:1096–1099
- Jin Y, Szabo LJ, Pretorius ZA, Singh RP, Ward R, Fetch T Jr (2008) Detection of virulence to resistance gene *Sr24* within race TTKS of *Puccinia graminis* f. sp. *tritici*. *Plant Dis* 92:923–926
- Jin Y, Szabo LJ, Rouse MN, Fetch T Jr, Pretorius ZA, Wanyera R, Njau P (2009) Detection of virulence to resistance gene *Sr36* within the TTKS race lineage of *Puccinia graminis* f. sp. *tritici*. *Plant Dis* 93:367–370
- Klindworth DL, Niu Z, Chao S, Friesen TL, Jin Y, Faris JD, Cai X, Xu SS (2012) Introgression and characterization of a goatgrass gene for a high level of resistance to Ug99 stem rust in tetraploid wheat. *G3* 2:665–673
- Knott DR (2000) The inheritance of stem rust resistance in Thatcher wheat. *Can J Plant Sci* 80:53–63
- Knott DR (2001) The relationship between seedling and field resistance to two races of stem rust in Thatcher wheat. *Can J Plant Sci* 81:415–418
- Kolmer JA, Dyck PL, Roelfs AP (1991) An appraisal of stem and leaf rust resistance in North American hard red spring wheats and the probability of multiple mutations to virulence in populations of cereal rust fungi. *Phytopathology* 81:237–239
- Kolmer JA, Garvin DF, Jin Y (2011) Expression of a Thatcher wheat adult plant stem rust resistance QTL on chromosome arm 2BL is enhanced by *Lr34*. *Crop Sci* 51:526–533
- Krattinger SG, Lagudah ES, Spielmeyer W, Singh RP, Heurta-Espino J, McFadden H, Bossolini E, Selter LL, Keller B (2009) A putative ABC transporter confers durable resistance to multiple fungal pathogens in wheat. *Science* 323:1360–1363
- Lagudah ES, McFadden H, Singh RP, Huerta-Espino J, Bariana HS, Spielmeyer W (2006) Molecular genetic characterization of the *Lr34/Yr18* slow rusting resistance gene region in wheat. *Theor Appl Genet* 114:21–30
- Lagudah ES, Krattinger SG, Herrera-Foessel S, Singh RP, Huerta-Espino J, Spielmeyer W, Brown-Guedira G, Selter LL, Keller B (2009) Gene-specific markers for the wheat gene *Lr34/Yr18/Pm38* which confers resistance to multiple fungal pathogens. *Theor Appl Genet* 119:889–898
- Lanning SP, Talbert LE, McGuire CF, Bowman HF, Carlson GR, Jackson GD, Eckhoff JL, Kushnak GD, Stougaard RN, Stallknecht GF, Wichman DM (1994) Registration of ‘McNeal’ wheat. *Crop Sci* 34:1126–1127
- Letta T, Maccaferri M, Badebo A, Ammar K, Ricci A, Crossa J, Tuberosa R (2013) Searching for novel sources of field resistance to Ug99 and Ethiopian stem rust races in durum wheat via association mapping. *Theor Appl Genet* 126:1237–1256
- Liu S, Yu L, Singh RP, Jin Y, Sorrells ME, Anderson JA (2010) Diagnostic and co-dominant PCR markers for wheat stem rust resistance genes *Sr25* and *Sr26*. *Theor Appl Genet* 120:691–697
- Liu W, Jin Y, Rouse M, Friebe B, Gill B, Pumphrey MO (2011a) Development and characterization of wheat-*Ae. searsii* Robertsonian translocations and a recombinant chromosome conferring resistance to stem rust. *Theor Appl Genet* 122:1537–1545
- Liu W, Rouse M, Friebe B, Jin Y, Gill B, Pumphrey MO (2011b) Discovery and molecular mapping of a new gene conferring resistance to stem rust, *Sr53*, derived from *Aegilops geniculata* and characterization of spontaneous translocation stocks with reduced alien chromatin. *Chromosome Res* 19:669–682
- Lorieux M, Goffinet B, Perrier X, De Gonzales Leon D, Laud C (1995) Maximum likelihood models for mapping genetic markers showing segregation distortion. 1. Backcross populations. *Theor Appl Genet* 90:73–80
- Lowe I, Jankuloski L, Chao S, Chen X, See D, Dubcovsky J (2011) Mapping and validation of QTL which confer partial resistance to broadly virulent post-2000 North American races of stripe rust in hexaploid wheat. *Theor Appl Genet* 123:143–157
- Luig NH (1983) A survey of virulence genes in wheat stem rust, *Puccinia graminis* f. sp. *tritici*. *Advances in plant breeding, Supplement 11 to Journal of Plant Breeding*. Paul Parey, Berlin
- Mago R, Spielmeyer W, Lawrence GJ, Lagudah ES, Ellis JG, Pryor A (2002) Identification and mapping of molecular markers linked to rust resistance genes located on chromosome 1RS of rye using wheat-rye translocation lines. *Theor Appl Genet* 104:1317–1324
- Mago R, Bariana HS, Dundas IS, Spielmeyer W, Lawrence GJ, Pryor AJ, Ellis JG (2005) Development of PCR markers for the selection of wheat stem rust resistance genes *Sr24* and *Sr26* in diverse wheat germplasm. *Theor Appl Genet* 111:496–504
- Mago R, Brown-Guedira G, Dreisigacker S, Breen J, Jin Y, Singh R, Appels R, Lagudah ES, Ellis J, Spielmeyer W (2011) An accurate DNA marker assay for stem rust resistance gene *Sr2* in wheat. *Theor Appl Genet* 122:735–744
- McIntosh RA, Dubcovsky J, Rogers WJ, Morris CF, Appels R, Xia XC (2012) Catalogue of gene symbols for wheat: 2012 supplement. *Annu Wheat Newsltt* 58:259–279
- Naruoka Y, Talbert LE, Lanning SP, Blake NK, Martin JM, Sherman JD (2011) Identification of quantitative trait loci for productive tiller number and its relationship to agronomic traits in spring wheat. *Theor Appl Genet* 123:1043–1053

- Nazari K, Mafi M, Yahyaoui A, Singh RP, Park RF (2009) Detection of wheat stem rust (*Puccinia graminis* f. sp. *tritici*) race TTKSK (Ug99) in Iran. *Plant Dis* 93:317
- Niu Z, Klindworth DL, Friesen TL, Chao S, Jin Y, Cai X, Xu SS (2011) Targeted introgression of a wheat stem rust resistance gene by DNA marker-assisted chromosome engineering. *Genetics* 187:1011–1021
- Njau PN, Jin Y, Huerta-Espino J, Keller B, Singh RP (2010) Identification and evaluation of sources of resistance to stem rust race Ug99 in wheat. *Plant Dis* 94:413–419
- Njau PN, Bhavani S, Huerta-Espino J, Keller B, Singh RP (2013) Identification of QTL associated with durable adult plant resistance to stem rust race Ug99 in wheat cultivar ‘Pavon 76’. *Euphytica* 190:33–44
- Olson EL, Brown-Geudira G, Marshall D, Stack E, Bowden RL, Jin Y, Rouse M, Pumphrey MO (2010) Development of wheat lines having a small introgressed segment carrying stem rust resistance gene *Sr22*. *Crop Sci* 50:1823–1830
- Peterson RF, Campbell AB, Hannah AE (1948) A diagrammatic scale for estimating rust intensity of leaves and stem of cereals. *Can J Res Sect C* 26:496–500
- Pretorius ZA, Singh RP, Wagoire WW, Payne TS (2000) Detection of virulence to wheat stem rust resistance gene *Sr31* in *Puccinia graminis* f. sp. *tritici* in Uganda. *Plant Dis* 84:203
- Pretorius ZA, Bender CM, Visser B, Terefe T (2010) First report of a *Puccinia graminis* f. sp. *tritici* race virulent to the *Sr24* and *Sr31* wheat stem rust resistance genes in South Africa. *Plant Dis* 94:784
- Pretorius ZA, Szabo LJ, Boshoff WHP, Herselman L, Visser B (2012) First report of a new TTKSF race of wheat stem rust (*Puccinia graminis* f. sp. *tritici*) in South Africa and Zimbabwe. *Plant Dis* 96:590
- Qi LL, Pumphrey MO, Friebe B, Zhang P, Qian C, Bowden RL, Rouse MN, Jin Y, Gill BS (2011) A novel Robertsonian translocation event leads to transfer of a stem rust resistance gene (*Sr52*) effective against race Ug99 from *Dasypyrum villosum* into bread wheat. *Theor Appl Genet* 123:159–167
- Roelfs AP (1978) Estimated losses caused by rust in small grain cereals in the United States: 1918–76. USDA Miscellaneous Publication, USA 1363
- Roelfs AP, Martens JW (1988) An international system of nomenclature for *Puccinia graminis* f. sp. *tritici*. *Phytopathology* 78:526–533
- Roelfs AP, Singh RP, Saari EE (1992) Rust diseases of wheat: concepts and methods of disease management. CIMMYT, Mexico
- Rouse MN, Wanyera R, Njau P, Jin Y (2011) Sources of resistance to stem rust race Ug99 in spring wheat germplasm. *Plant Dis* 95:762–766
- Rouse MN, Nava IC, Chao SM, Anderson JA, Jin Y (2012) Identification of markers linked to the race Ug99 effective stem rust resistance gene *Sr28* in wheat (*Triticum aestivum* L.). *Theor Appl Genet* 125:877–885
- Saal B, Wricke G (1999) Development of simple sequence repeat markers in rye (*Secale cereale* L.). *Genome* 42:964–972
- Sambasivam PK, Bansal UK, Hayden MJ, Dvorak J, Lagudah ES, Bariana HS (2008) Identification of markers linked with stem rust resistance genes *Sr33* and *Sr45*. In: Appels R, Eastwood R, Lagudah E, Langridge P, Mackay M, McIntyre L, Sharp P (eds) Proceedings of 11th international wheat genetics symposium. Sydney University Press, Australia, pp 351–353
- Sherman JD, Martin JM, Blake NK, Lanning SP, Talbert LE (2014) Genetic basis of agronomic differences between a modern and a historical spring wheat cultivar. *Crop Sci* 54:1–13
- Simons K, Abate Z, Chao S, Zhang W, Rouse M, Jin Y, Elias E, Dubcovsky J (2011) Genetic mapping of stem rust resistance gene *Sr13* in tetraploid (*Triticum turgidum* ssp. *durum* L.). *Theor Appl Genet* 122:649–658
- Singh RP, McIntosh RA (1987) Genetics of resistance to *Puccinia graminis tritici* in ‘Chris’ and ‘W3746’ wheats. *Theor Appl Genet* 73:846–855
- Singh RP, Hodson DP, Huerta-Espino J, Jin Y, Bhavani S, Njau P, Herrera-Foessel S, Singh PK, Singh S, Govindan V (2011a) The emergence of Ug99 races of the stem rust fungus is a threat to world wheat production. *Annu Rev Phytopathol* 49:465–481
- Singh RP, Huerta-Espino J, Bhavani S, Herrera-Foessel SA, Singh D, Singh PK, Velu G, Mason RE, Jin Y, Njau P, Crossa J (2011b) Race non-specific resistance to rust diseases in CIMMYT spring wheats. *Euphytica* 179:175–186
- Singh A, Knox RE, DePauw RM, Singh AK, Cuthbert RD, Campbell HL, Singh D, Bhavani S, Fetch T, Clarke F (2013a) Identification and mapping in spring wheat of genetic factors controlling stem rust resistance and the study of their epistatic interactions across multiple environments. *Theor Appl Genet* 126:1951–1964
- Singh A, Pandey MP, Singh AK, Knox RE, Ammar K, Clarke JM, Clarke FR, Singh RP, Pozniak CJ, DePauw RM, McCallum BD, Cuthbert RD, Randhawa HS, Fetch TG Jr (2013b) Identification and mapping of leaf, stem and stripe rust resistance quantitative trait loci and their interactions in durum wheat. *Mol Breeding* 31:405–418
- Singh S, Singh RP, Bhavani S, Huerta-Espino J, Lopez-Vera EE (2013c) QTL mapping of slow-rusting, adult plant resistance to race Ug99 of stem rust fungus in PBW343/Muu RIL population. *Theor Appl Genet* 126:1367–1375
- Snappe JW, Riggs TJ (1975) Genetical consequences of single seed descent in the breeding of self-pollinating crops. *Heredity* 35:211–219
- Stakman EC, Steward DM, Loegering WQ (1962) Identification of physiologic races of *Puccinia graminis* var. *tritici*. USDA Agric Res Serv E-617
- R Development Core Team (2010) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>
- Tsilo TJ, Jin Y, Anderson JA (2007) Microsatellite markers linked to stem rust resistance allele *Sr9a* in wheat. *Crop Sci* 47:2013–2020
- Tsilo TJ, Jin Y, Anderson JA (2008) Diagnostic microsatellite markers for the detection of stem rust resistance gene *Sr36* in diverse genetic backgrounds of wheat. *Crop Sci* 48:253–261
- Wang S, Basten CJ, Zeng ZB (2007) Windows QTL Cartographer 2.5. Department of Statistics, North Carolina State University, Raleigh, NC. <http://statgen.ncsu.edu/qtlcart/WQTLCart.htm>
- Wanyera R, Kinyua MG, Jin Y, Singh RP (2006) The spread of stem rust caused by *Puccinia graminis* f. sp. *tritici*, with virulence on *Sr31* in wheat in Eastern Africa. *Plant Dis* 90:113
- Wu S, Pumphrey M, Bai G (2009) Molecular mapping of stem-rust-resistance gene *Sr40* in wheat. *Crop Sci* 49:1682–1686
- Young ND (1996) QTL mapping and quantitative disease resistance in plants. *Annu Rev Phytopathol* 34:479–501
- Yu L, Lorenz A, Rutkoski J, Singh RP, Bhavani S, Huerta-Espino J, Sorrells ME (2011) Association mapping and gene–gene interaction for stem rust resistance in CIMMYT spring wheat germplasm. *Theor Appl Genet* 123:1257–1268
- Yu L, Morgounov A, Wanyera R, Keser M, Singh SK, Sorrells M (2012) Identification of Ug99 stem rust resistance loci in winter wheat germplasm using genome-wide association analysis. *Theor Appl Genet* 125:749–758
- Yu L, Barbier H, Rouse MN, Singh S, Singh RP, Bhavani S, Huerta-Espino J, Sorrells ME (2014) A consensus map for Ug99 stem

- rust resistance loci in wheat. *Theor Appl Genet*. doi:[10.1007/s00122-014-2326-7](https://doi.org/10.1007/s00122-014-2326-7) (companion paper)
- Zhang W, Olson EL, Saintenac C, Rouse M, Abate Z, Jin Y, Akhunov E, Pumphrey MO, Dubcovsky J (2010) Genetic maps of stem rust resistance gene *Sr35* in diploid and hexaploid wheat. *Crop Sci* 50:2464–2474
- Zheng ZB (1993) Theoretical basis for separation of multiple linked gene effects in mapping quantitative trait loci. *Proc Natl Acad Sci USA* 90:10972–10976
- Zheng ZB (1994) Precision mapping of quantitative trait loci. *Genetics* 136:1457–1468