

Nutrient Availability from Douglas Fir Bark in Response to Substrate pH

James E. Altland^{1,3}

USDA-ARS, Application Technology Research Unit, Ohio Agricultural Research and Development Center, 208 Ag. Eng. Building, 1680 Madison Avenue, Wooster, OH 44691

M. Gabriela Buamscha²

USDA Forest Service, 333 SW First Avenue, Portland, OR 97208-3623

Additional index words. *Pseudotsuga menziesii*, lime, calcium carbonate, calcium hydroxide

Abstract. Two studies were conducted to determine the influence of substrate pH on nutrient availability in Douglas fir bark. Douglas fir bark was amended with either calcium carbonate (CaCO₃) or calcium hydroxide [Ca(OH)₂] at 13 rates to generate substrates with low to high pH. A nonamended control was also maintained. The substrates received no other fertilizer amendments. Substrates were filled into nursery containers (2.8 L) and maintained in a nursery setting with 1.2 cm·d⁻¹ overhead irrigation. At 1 and 6 weeks after potting, four containers of each lime treatment were harvested to determine substrate pH and availability of 13 nutrients. Substrate solutions used for analysis were obtained with the saturated media extract method. Water-extractable phosphorus and DTPA-extractable boron, iron, copper, and aluminum decreased with increasing substrate pH. Other nutrients were either nonresponsive to substrate pH or the observed response was deemed more likely caused by calcium competition on cation exchange sites.

Ornamental container crops in the Pacific Northwest are grown primarily in Douglas fir [*Pseudotsuga menziesii* (Mirbel) Franco] bark (DFB). Similar to pine (*Pinus taeda* L.) bark in the southeast United States, DFB comprises the highest portion of most nursery substrates (60% to 80% of the substrate mix, pers. obser.). Chemical and physical properties of DFB, as they pertain to use in container substrates, have recently been documented by Buamscha et al. (2007b). They reported nonamended DFB to have low pH (3.7 to 4.4); little or no available nitrogen (N); relatively high water-extractable phosphorus (P; 10 to 28 mg·L⁻¹) and potassium (K; 78 to 162 mg·L⁻¹); low levels of calcium (Ca), magnesium (Mg), and sulfur (S); and moderate levels of some micronutrients. In a separate study, Buamscha et al. (2007a) documented that DFB alone provides sufficient micronutrients for annual vinca [*Catharanthus roseus* (L.) G. Don 'Peppermint Cooler'] grown at low pH (4.5 to 5.5). Macronutrient and micronutrient availability

may not be sufficient to support plant growth when substrate pH is higher. Altland (2006) reported reduced growth of Japanese maple (*Acer palmatum* var. *atropurpureum* Thunb.), hydrangea [*Hydrangea macrophylla* Thunb. (Ex J.A. Murr.) Ser. 'Endless Summer'], and leucothoe [*Leucothoe axillaris* (Lam.) D. Don] caused by a pH-induced reduction of available N, P, and micronutrients in DFB. Similar observations of reduced plant growth with high substrate pH have been reported for crops grown in pine bark (Wright and Hinsley, 1991).

Little is known empirically about how pH affects nutrient availability in DFB substrates. Lucas and Davis (1961) determined the relationship between pH and nutrient availability in organic soils. They concluded that the ideal pH range (in terms of total nutrient availability) for wood-sedge soils to be between 5.5 and 5.8 and 5.0 for sphagnum peat soils. They further commented that this range is one to 1.5 points lower than what was considered ideal for mineral soils. This report formed the basis for future studies years later as the greenhouse industry switched from mineral soils to those composed primarily of peat or bark. Peterson (1980) documented the effect of substrate pH on macronutrient and micronutrient availability in a well-fertilized commercial greenhouse substrate (peat moss, perlite, vermiculite, granite sand, and composted pine bark; ratios not given). His study agreed with Lucas and Davis (1961) in that the optimum pH range was 5.2 to 5.5, which he characterized as being a whole pH unit or more below what is considered optimum for mineral soils. Peterson (1980) reported decreasing availability of P, iron (Fe), man-

ganese (Mn), boron (B), zinc (Zn), and copper (Cu) with increasing pH. Argo (1998) reviewed the effects of pH on nutrient availability in soilless substrates citing numerous sources and generally agreed with conclusions from Peterson (1980).

In view of the widespread use of DFB in the Pacific Northwest, and the corresponding lack of information available on its chemical properties, two experiments were initiated to document the influence of pH on nutrient availability. The objectives of these experiments were to determine the influence of two lime sources on DFB pH as well as nutrient availability with respect to changes in substrate pH.

Materials and Methods

Expt. 1. On 20 June 2006, pH of 13 batches of DFB (each 0.03 m³) was adjusted using 13 rates of either calcium carbonate (CaCO₃; Imperial Limestone; J.A. Jack and Sons, Seattle, WA) or hydrated lime [calcium hydroxide, Ca(OH)₂]; Kemilime; Ash Grove Cement Co, Portland, OR] (Table 1). A non-amended control (no lime) was also included. Rates for each lime type were selected to provide substrates with a spectrum of low to high pH and were based on previous research by the authors (unpublished data). Calcium carbonate [97% CaCO₃, 2% MgCO₃, 97 calcium carbonate equivalency (CCE)] was processed such that 100% and 80% passed through 40 and 100 mesh, respectively. Hydrated lime [94% Ca(OH)₂, 126 CCE] was processed such that 100% and 99% passed through 40 and 100 mesh, respectively. Bark was ground with a hammer-mill, passed through a 2.2-cm screen, and aged for ≈6 months (Marr Bros. Co., Monmouth, OR). Eight No. 1 containers (2.8 L) were filled with each bark batch and maintained in a retractable roof greenhouse (Cravo Equip. Ltd., Brantford, Ontario, Canada) in Aurora, OR. Containers were overhead-irrigated with groundwater at 1.2 cm·d⁻¹ (Table 2). Four containers of each treatment were randomly selected and harvested 1 and 6 weeks after potting (WAP) by placing the entire contents of the container into a plastic bag for delivery to the laboratory. Bark samples were analyzed for pH, ammonium (NH₄-N), nitrate (NO₃-N), P, K, Ca, Mg, and sulfate (SO₄²⁻) using the saturated media extract (SME) method with deionized water as the extractant (Gavlak et al., 2003; Warncke, 1998). Boron, Fe, Mn, Cu, Zn, and aluminum (Al) were analyzed using a SME with diethylenetriaminepentaacetic acid (DTPA) as the extractant (Warncke, 1998). Following Gavlak et al. (2003), DFB was soaked in the extractant (either water or DTPA) for 24 h. Ammonium and NO₃-N in extracted solutions were analyzed colorimetrically using a Lachat Quick Chem 8000 (Lachat Instruments, Milwaukee, WI). All other elements were analyzed with inductively coupled plasma-emission spectrometry (Thermo Jarrel Ash, Offenbach, Germany).

Received for publication 12 Sept. 2007. Accepted for publication 24 Oct. 2007.

Special thanks to the Oregon Association of Nurseries for funding this research and to Gilda Medina, Vyacheslav Dobryk, and Magdalena Zazirska for skillful technical assistance.

Mention of proprietary products or company is included for the reader's convenience and does not imply any endorsement or preferential treatment by USDA/ARS.

¹Research Horticulturist.

²Western Nursery Specialist.

³To whom reprint requests should be addressed; e-mail James.Altland@ars.usda.gov

Table 1. Water and DTPA-extractable nutrients in Douglas fir bark after incorporation of two lime types with multiple rates.^z

Harvest time (weeks)	Lime type	Rate (kg·m ⁻³)	Water extraction ^y							DTPA extraction					
			pH	P	K	Ca	Mg	SO ₄ ²⁻	Na	B	Fe	Mn	Cu	Zn	Al
1	CaCO ₃	0	4.9	5.5	17.8	2.6	1.2	5.3	6.1	0.30	54.8	10.6	0.40	2.4	22.9
		0.6	5.4	7.1	24.9	5.8	1.7	5.4	6.7	0.28	50.3	13.5	0.37	2.4	18.1
		1.5	6.1	5.8	27.2	8.5	1.9	5.3	7.0	0.23	44.0	14.9	0.33	2.4	12.5
		3.0	6.5	4.5	27.5	11.9	2.2	5.2	7.1	0.16	31.8	15.8	0.28	2.4	5.5
		4.4	6.8	5.2	28.5	15.0	2.5	4.5	6.6	0.12	20.3	15.3	0.23	2.2	1.8
		5.9	6.9	5.0	29.1	18.3	2.8	4.7	6.4	0.10	13.0	13.6	0.20	1.9	1.4
		8.9	7.0	5.4	27.7	18.4	2.8	4.6	6.4	0.11	14.0	14.3	0.22	2.0	1.4
	Ca(OH) ₂	11.9	6.9	4.2	21.6	15.0	2.4	5.6	7.6	0.10	9.9	13.1	0.20	1.9	1.3
		4.4	7.2	3.0	17.4	18.8	2.0	6.2	8.2	0.09	7.3	12.6	0.22	1.9	1.0
		5.9	7.4	4.6	37.0	43.2	3.7	9.1	10.2	0.08	4.3	13.6	0.21	2.0	0.8
		8.9	7.9	2.3	42.8	73.6	4.8	10.0	11.2	0.07	1.7	13.5	0.20	2.1	0.9
		11.9	7.8	2.1	57.4	112.2	6.5	9.8	9.2	0.08	1.0	15.6	0.22	2.6	0.9
		17.8	7.9	1.8	84.7	240.2	11.8	12.7	10.3	0.06	0.8	14.6	0.23	2.6	0.7
		23.7	8.8	1.2	84.2	293.4	9.5	16.0	11.5	0.05	0.3	10.5	0.17	2.2	0.7
6	CaCO ₃	0.0	6.0	1.6	4.3	1.8	0.7	6.5	7.9	0.15	40.8	14.6	0.35	2.5	10.2
		0.6	6.2	2.5	6.1	1.1	0.5	7.2	9.7	0.16	37.7	14.9	0.32	2.3	9.5
		1.5	6.7	2.0	5.7	2.2	0.8	6.0	9.4	0.13	30.4	15.6	0.28	2.3	5.6
		3.0	7.2	2.1	6.6	3.9	1.1	10.2	9.7	0.10	21.0	14.3	0.23	2.0	3.0
		4.4	7.4	2.4	8.0	5.3	1.2	8.9	10.9	0.07	15.9	14.7	0.22	2.0	1.7
		5.9	7.5	2.0	6.2	4.4	1.1	6.7	11.9	0.06	11.3	15.6	0.21	2.1	1.0
		8.9	7.7	1.6	4.9	3.6	0.9	6.7	12.5	0.05	7.1	14.9	0.20	2.0	0.7
	Ca(OH) ₂	11.9	7.7	2.0	7.4	8.9	1.6	7.0	12.0	0.05	6.2	14.4	0.18	1.9	0.6
		4.4	7.7	4.1	10.9	14.7	2.2	6.6	9.5	0.07	15.4	16.1	0.24	2.2	1.3
		5.9	7.8	3.4	7.5	22.2	2.6	7.8	11.1	0.05	10.1	15.7	0.23	2.3	0.8
		8.9	8.0	1.9	6.3	32.9	2.8	7.0	10.1	0.04	2.5	12.6	0.18	2.0	0.5
		11.9	8.2	1.0	7.5	30.7	2.5	7.6	10.7	0.05	1.4	10.7	0.17	1.7	0.6
		17.8	8.4	0.7	10.6	60.9	4.3	8.0	11.0	0.04	1.0	8.0	0.14	1.4	0.5
		23.7	8.5	0.6	16.1	72.4	4.6	9.3	11.0	0.04	0.8	7.0	0.13	1.2	0.6
Main effects															
Lime type (L)			0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
Lime rate (lime type) R(L)			0.0001	0.0001	0.0001	0.0001	0.0001	0.1931	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	
Harvest time (T)			0.0001	0.0001	0.0001	0.0001	0.7090	0.0001	0.0001	0.0001	0.0801	0.0001	0.0001	0.0001	
T*L			0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.4045	0.0001	0.0001	
T*R(L)			0.0003	0.0001	0.0001	0.0001	0.0017	0.0062	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	

^zExtractions performed 1 and 6 weeks after potting (WAP). Containers subjected to typical nursery irrigation (1.2 cm·d⁻¹).

^yWater and 0.005 M DTPA (diethylenetriaminepentaacetic acid) extractions were conducted over 24 h.

Table 2. Chemical characteristics of groundwater used for irrigation at the Oregon State University North Willamette Research and Extension Center.

pH	7.0
Alkalinity	87.3 mg·L ⁻¹
Hardness	86.8 mg·L ⁻¹
Total dissolved solids	129.5 mg·L ⁻¹
Electrical conductivity	0.2 dS·m
Sodium	6.9 mg·L ⁻¹
Chloride	1.5 mg·L ⁻¹
Potassium	2.1 mg·L ⁻¹
Calcium	19.8 mg·L ⁻¹
Magnesium	8.9 mg·L ⁻¹
Sulfate	12.9 mg·L ⁻¹
Iron	0.3 mg·L ⁻¹
Manganese	0.1 mg·L ⁻¹
Boron	0.0 mg·L ⁻¹
Copper	0.0 mg·L ⁻¹
Zinc	0.1 mg·L ⁻¹
Aluminum	0.2 mg·L ⁻¹

Expt. 2. On 8 Aug. 2006, a second repetition of the experiment was conducted similarly to the previous one. Data from both experiments were analyzed with repeated measures analysis of variance to determine significant main effects (Table 1). Nonlinear regression was used to determine the most appropriate model for pH response to lime rate as well as nutrient availability response to substrate pH. Procedures described by

Schabenberger and Pierce (2002) were used for nonlinear model selection and comparison. All statistical analyses were conducted with SAS 9.1 (SAS Institute, Cary, NC), although figures were constructed with SigmaPlot 10.0 (Systat Software, San Jose, CA).

Results and Discussion

Results from the two repetitions of the experiment were similar; thus, data were pooled in Table 1 and presented together in Figures 1 through 3. As a result of lime amendment, substrate pH ranged from 4.8 to 9.4 across both experiments (Figs. 2 and 3). Substrate pH increased exponentially with increasing CaCO₃ rate (Figs. 1A and 1C). Williams et al. (1988) reported that pH of sphagnum peat increases quadratically with increasing lime rates up to 7 kg·m⁻³. However, use of quadratic functions implies that pH would decline with increasing lime rate at some point beyond the observed range, which is unlikely. Exponential curves [pH = c + a(1 - b^x)] provide a more realistic fit to the data in that pH increases rapidly but then plateaus at some maximum value. Furthermore, parameter estimates of the exponential equation provide intuitive interpretation in that the solved estimate for c is pH of the substrate with 0 kg·m⁻³ (or nonamended pH), the

sum of c and a is equal to the maximum pH (or the point where the curve plateaus), and b (greater than 0) is a gauge of how rapidly pH increases from its minimum to maximum (steepness of the curve). Williams et al. (1988) suggest that as exchange sites become saturated with Ca²⁺ and Mg²⁺ ions, an increasing number of these ions were needed to further dissociate H⁺ into the substrate solution. This is a logical explanation considering the exponential relationship between lime rate and substrate pH. Substrate pH also increased exponentially with increasing Ca(OH)₂ rate (Figs. 1B and 1D).

Within a lime treatment, substrate pH increased from 1 to 6 WAP (Table 1). Across both lime types and experiments, adding more than 3 to 6 kg·m⁻³ lime had little or no effect on substrate pH. Williams et al. (1988) reported slightly higher dolomitic lime rates were necessary to maximize pH in peat substrates.

Macronutrients. Extractable NH₄-N and NO₃-N levels were low throughout the study regardless of lime treatment and did not respond to substrate pH (data not presented). Douglas fir bark contains very little N. Bollen (1969) documented the C:N ratio of DFB as 491:1 with only 0.11% total N on a dry weight basis. Others have documented the influence of substrate pH on N form (Niemiera and Wright, 1986; Ogden et al., 1987)




Fig. 1. Douglas fir bark pH response to CaCO_3 and $\text{Ca}(\text{OH})_2$ rates and 1 and 6 weeks after potting (WAP) in two experiments conducted in June (A and B) and Aug. (C and D) 2006.

in well-fertilized containers. However, considering low N levels inherent in DFB, it is not surprising that N in nonfertilized DFB substrate did not respond to pH.

Water-extractable P decreased with increasing pH at all collection dates (Fig. 2A). Lucas and Davis (1961) explained that P availability in an organic soil (greater than 50% organic matter) drops when pH is below 5.5 as a result of increased availability of Fe and Al, which form precipitates with soluble P. Our data show no such decrease in availability below pH 5.5. Bunt (1988) suggests differences in P uptake by plants results from changes in the form of water-soluble P from H_2PO_4^- to HPO_4^{2-} , the former being 10 times more available for plant uptake than the latter. Argo and Biernbaum (1996) report the equilibrium constant between these two phosphate forms is 7.2, so that when substrate pH is greater than 7.2, the less available HPO_4^{2-} predominates and plant-available P is greatly reduced. Argo and Biernbaum (1996) further demonstrate that when substrate pH is less than 7.2, plant P level is relatively unaffected regardless of the measurable decrease in water-extractable P with increasing pH. Our data show that total water-extractable P decreased (without in-

creasing) over the substrate pH range of 4.8 to 9.4, although our analytical methods do not distinguish between the two species of water-soluble P, H_2PO_4^- and HPO_4^{2-} . Peterson (1980) showed a similar trend to ours with his analysis of the effect of substrate pH on a peat-based mix.

Buamscha et al. (2007b) reported 10 to 28 $\text{mg}\cdot\text{L}^{-1}$ P in nonamended DFB analyzed immediately after collection from bark suppliers. However, by 1 WAP, water-extractable P in this experiment rarely exceeded 10 $\text{mg}\cdot\text{L}^{-1}$. Within a given lime treatment, water-extractable P decreased from 1 to 6 WAP ($P = 0.0001$). Warncke (1998) suggests that 3 to 5 $\text{mg}\cdot\text{L}^{-1}$ P is sufficient for growth of most container crops using the SME procedure. Although extractable P is initially high in DFB, it is quickly leached from the substrate. Yeager and Wright (1982) reported similarly high levels of P in pine bark, but 67% of that P was leached after one simulated irrigation event.

Water-extractable K did not respond to substrate pH (Fig. 2B). By 6 WAP, levels of water-extractable K were low regardless of substrate pH compared with the recommendation by Warncke (1998) (60 to 149 $\text{mg}\cdot\text{L}^{-1}$). Lucas and Davis (1961) reported

similar findings on organic peat soils in Michigan, but warns that conditions of high Ca (as would be typical if high lime rates are used) will competitively suppress uptake of K. Peterson (1980) showed no effect of pH on a well-fertilized greenhouse substrate. Buamscha et al. (2007b) reported 79 to 162 $\text{mg}\cdot\text{L}^{-1}$ K in a nonamended DFB. By 1 WAP, water-extractable K levels averaged just 39 $\text{mg}\cdot\text{L}^{-1}$ across lime rates and 15 $\text{mg}\cdot\text{L}^{-1}$ by 6 WAP (Table 1). Potassium is readily leached from containers (Broschat, 1995); thus, leaching is likely the cause of progressively lower amounts of extractable K over time. Water-extractable K increased with increasing $\text{Ca}(\text{OH})_2$ rates, but this is more likely the result of dislodging of K from cation exchange sites with increasing levels of soluble Ca.

Secondary nutrients. Levels of water-extractable Ca were more a function of Ca solubility of the lime products than of substrate pH. Within each harvest date, water-extractable Ca increased moderately with increasing CaCO_3 rates but increased more rapidly with increasing $\text{Ca}(\text{OH})_2$ rates (Table 1). Higher Ca levels would be expected from $\text{Ca}(\text{OH})_2$ amendments than CaCO_3 as a result of solubility and the percent Ca content of each product. Calcium hydroxide has a




Fig. 2. Water-extractable phosphorus (P), potassium (K), calcium (Ca), and magnesium (Mg) in Douglas fir bark using the saturated extraction method. Extractions were performed 1 and 6 weeks after potting (WAP) in two separate experiments. Containers were subjected to typical nursery irrigation ($1.2 \text{ cm}\cdot\text{d}^{-1}$) throughout experiments.

solubility product of $K_{sp} = 4.68 \times 10^{-6}$, whereas that for calcium carbonate is $K_{sp} = 8.7 \times 10^{-9}$ (Lide, 2007), indicating over a thousand-fold difference in solubility. Furthermore, CaCO_3 is $\approx 37\%$ Ca, whereas Ca(OH)_2 is $\approx 67\%$ Ca. Argo (1998) argued that low pH does not reduce Ca availability, but instead low pH is the result of insufficient Ca sources (lime or high-alkalinity water) applied to the substrate. Our data tend to support this notion in that Ca availability is a function of lime type and rate and not a function of pH.

Similar to Ca, water-extractable Mg increased with increasing pH but was more likely a function of increasing lime rate (Figs. 2C and 2D). Although both lime sources were Ca-based, both had low levels of Mg contamination (less than 2%). It is also possible that Ca–Mg competition for cation exchange sites caused an increase in extractable Mg at higher rates of Ca(OH)_2 . Peterson (1980) reported that Mg increased with increasing pH, in which pH was similarly adjusted with Ca(OH)_2 amendment.

Sulfate levels were affected by lime type and rate (Table 1); however, the differences in water-extractable sulfate levels across lime types and rates were minor (less than $5 \text{ mg}\cdot\text{L}^{-1}$) by 6 WAP. Sulfate anions are easily

leached from container substrates (Yeager and Barrett, 1985). Nonetheless, sulfate levels did not decrease between 1 and 6 WAP ($P = 0.709$), and they were similar to levels reported by Buamscha (2007b) for nonamended and nonirrigated DFB (10 to $20 \text{ mg}\cdot\text{L}^{-1}$). Browder (2004) concluded that 7 to $15 \text{ mg}\cdot\text{L}^{-1}$ sulfate (as measured in a pour-through procedure) should be present for optimum growth of pin oak (*Quercus palustris* Münchh.) and Japanese maple (*Acer palmatum* Thunb.). Handreck (1986) states that $6 \text{ mg}\cdot\text{L}^{-1}$ sulfate from a water extraction is sufficient for any crop. By this standard, sulfate levels are low to marginally sufficient. However, this is of little consequence because sulfate is provided in many fertilizer types such as ammonium sulfate [$(\text{NH}_4)_2\text{SO}_4$] used in controlled-release fertilizers, gypsum (CaSO_4), and sulfated micronutrients.

Micronutrients. Several studies have documented the micronutrient capacity of soilless substrates, each suggesting that the substrate alone may be a sufficient source of micronutrients to meet plant need without addition of supplemental micronutrient fertilizers (Buamscha et al., 2007a; Niemiera, 1992; Rose and Wang, 1999). Availability of micronutrients has been documented as pH-sensitive in organic soils and substrates

(Lucas and Davis, 1961; Peterson, 1980); thus, the degree to which a substrate can provide sufficient micronutrients to support crop growth should always be qualified with a target pH range.

DTPA-extractable B decreased exponentially with increasing substrate pH (Fig. 3A). Extractable B decreased slightly from 1 to 6 WAP; however, the relationship between substrate pH and extractable B remained consistent across experiments and harvest dates. Lucas and Davis (1961) warn that there is a strong Ca–B relationship in organic soils, and some peat soils high in Ca are more prone to B deficiency as a result of elevated Ca. In DFB, the relationship between DTPA-extractable B and pH is more consistent and predictable than the relationship between DTPA-extractable B and water-extractable Ca (data not shown). Warncke (1998) suggests DTPA-extractable B should range from 0.7 to $2.5 \text{ mg}\cdot\text{L}^{-1}$; thus, extractable B would be considered low in nonamended DFB. Buamscha et al. (2007a) demonstrated that DFB, even amended with a micronutrient fertilizer package (Micromax; The Scotts Co., Marysville, OH), had lower than recommended extractable B levels 5 and 8 WAP, but higher than recommended foliar B in annual vinca [*Catharanthus roseus* (L.) G.




Fig. 3. Relationship between DTPA-extractable boron (B), iron (Fe), copper (Cu), and aluminum (Al) in Douglas fir bark using the saturated extraction method. Extractions were performed 1 and 6 weeks after potting (WAP) in two separate experiments. Containers were subjected to typical nursery irrigation ($1.2 \text{ cm} \cdot \text{d}^{-1}$) throughout the experiment.

Don ‘Peppermint Cooler’]. The relationship between B requirements in plant foliage and substrate-extractable B requires further study to improve recommended sufficiency ranges.

DTPA-extractable Fe decreased exponentially with increasing substrate pH with Fe dropping below recommended levels at $\text{pH} \approx 7$ (Fig. 3B). Berghage et al. (1987) reported DTPA-extractable Fe decreased quadratically over the pH range of 3.5 to 7.5. Despite different mathematical models used to describe our data and that of Berghage et al. (1987), Fe response to pH in the two studies appears to be similar. Substrate Fe levels should range from 15 to $40 \text{ mg} \cdot \text{L}^{-1}$ (Warncke, 1998). Douglas fir bark has sufficient extractable Fe so long as pH is below 7. Lucas and Davis (1961) commented that Fe deficiency is rarely a problem in organic soils as a result of an abundance of humic acids maintaining Fe levels.

Neither Mn nor Zn responded to substrate pH (data not shown). Berghage et al. (1987) reported that DTPA-extractable Mn was slightly affected (not specific) and Zn was unaffected over a substrate pH range of 3.5 to 7.5. Extractable Mn levels in all substrates were above the recommended range (5 to $30 \text{ mg} \cdot \text{L}^{-1}$), whereas extractable Zn was below recommended levels (5 to $30 \text{ mg} \cdot \text{L}^{-1}$). Buamscha

et al. (2007a) demonstrated that fresh and aged DFB amended only with nitrogen–phosphorus–potassium fertilizers provided sufficient Zn for annual vinca through 8 weeks of production despite low levels of DTPA-extractable Zn in the substrate.

DTPA-extractable Cu decreased exponentially with increasing substrate pH (Fig. 3C). Handreck (1994) reported DTPA-extractable Cu in pine (*Pinus radiata* D. Don) bark was unaffected by substrate pH over the range 4.5 to 6.5. Either DFB differs substantially from pine bark evaluated by Handreck or the narrower range of pH observed in his study was insufficient to reveal a relationship. Handreck (1990) in a separate study reported that DTPA-extractable Cu in substrates should be greater than $0.25 \text{ mg} \cdot \text{L}^{-1}$ to avoid deficiency symptoms in flowering mums (*Chrysanthemum* \times *morifolium* Ramat. ‘Yellow Mandalay’), but levels should be at least $5.1 \text{ mg} \cdot \text{L}^{-1}$ to maximize plant growth.

Nonessential elements. Water-extractable Na did not respond to substrate pH (data not shown). Warncke (1998) recommends extractable Na levels in container substrates be less than $40 \text{ mg} \cdot \text{L}^{-1}$. After 6 weeks of irrigation, water-extractable Na ranged from 3 to $15 \text{ mg} \cdot \text{L}^{-1}$ across both experiments with

levels increasing slightly from 1 to 6 WAP. Increased Na levels over time were likely the result of accumulation from low levels of Na in irrigation water (Table 2).

Extractable Al in substrates is rarely considered. This is possibly the result of the general agreement that organic soils and soilless media contain low amounts of Al (Lucas and Davis, 1961; Yeager and Barrett, 1985). Significant amounts of Al (7.8 to $51.5 \text{ mg} \cdot \text{L}^{-1}$) were extracted from DFB in samples collected from bark suppliers (before amendment or irrigation in a nursery) (Buamscha et al., 2007b). DTPA-extractable Al decreased exponentially with increasing pH (Fig. 3D). Aluminum levels averaged $23.9 \text{ mg} \cdot \text{L}^{-1}$ in nonamended bark 1 WAP (Table 1), although levels decreased over time. Wright (1989) reviewed Al interactions with soils and crops, and described Al speciation as complex, dependent on soil pH and other mineralogical factors and difficult to predict. Wright (1989) also explains that Al forms complex ligands with sulfate and soluble organic compounds, which alleviate Al toxicity. In contrast to the notion that Al is lacking or only present in low concentration in bark substrates, it is likely that it is present in relatively high levels (pH-dependent) but in nontoxic forms as a result of a consistent

supply of sulfate from bark (Table 1) and fertilizer amendments as well as the presence of soluble organic compounds in DFB. It is unclear what levels of DTPA-extractable Al should be considered toxic in nursery substrates.

Water-extractable P and DTPA-extractable B, Fe, Cu, and Al are responsive to pH in nonamended DFB. Other nutrients were either nonresponsive to substrate pH or the observed response was deemed more likely caused by Ca competition on cation exchange sites. Although Ca and its competitive effects on cation exchange sites may be an undesirable artifact in a study such as this, applications of lime (and thus Ca) is the primary mechanism by which nursery growers adjust substrate pH. Irrigation water alkalinity (dissolved CO_3^{2-} and HCO_3^-) is another major factor affecting substrate pH, but across three sampling dates, Ca and alkalinity levels in Oregon nursery irrigation sources were highly correlated ($r = 0.80$, $n = 61$, unpublished data).

These results have several practical implications. Phosphorus levels across all lime treatments declined from 1 to 6 WAP, suggesting that DFB alone is not a sufficient source of P to sustain plant growth throughout a growing season. However, the common industry practice of incorporating a water-soluble source of P into the substrate as a starter fertilizer should be questioned. There is sufficient water-extractable P in the first few weeks of production contingent that substrate pH is low (less than 6). Water-extractable K declined steadily over time regardless of pH and thus should be supplemented in some sort of continuous manner throughout the production cycle of the crop. Like P, K is readily available from DFB soon after potting and thus its application in starter fertilizer packages is questionable. Calcium and Mg levels were dependent on lime rate. In typical nursery conditions, it would be difficult to appreciably raise substrate pH without the addition of calcium and magnesium carbonates (through lime application or irrigation water alkalinity). Sulfate levels were moderately low, but this is probably inconsequential considering the widespread use of sulfate-based fertilizers. Douglas fir bark has been reported to provide sufficient micronutrients for short-term container crops (Buamscha et al., 2007a); however, Buamscha warns that this observation may only be

valid at low pH (less than 5.7). This warning is justified considering the pH-dependent availability of Fe, B, and Cu in DFB.

Data herein provide a realistic model of how plant essential nutrients respond to substrate pH in DFB. These data suggest that even low lime rates have a profound effect on nutrient availability in DFB and that lime should be used in moderation. However, it is uncertain if substrate pH would similarly affect nutrient availability in substrates receiving supplemental fertilizers typical of nursery production. Future research will address the impact of lime additions and substrate pH on nutrient availability in well-fertilized DFB substrates.

Literature Cited

- Altland, J.E. 2006. Substrate pH, a tricky topic. *Digger* 50:42–47.
- Argo, W.R. 1998. Root medium chemical properties. *HortTechnology* 8:446–493.
- Argo, W.R. and J.A. Biernbaum. 1996. The effect of lime, irrigation-water source, and water-soluble fertilizer on root-zone pH, electrical conductivity, and macronutrient management of container root media with impatiens. *J. Amer. Soc. Hort. Sci.* 121:442–452.
- Berghage, R.D., D.M. Krauskopf, D.D. Warncke, and I. Widders. 1987. Micronutrient testing of plant growth media: Extractant identification and evaluation. *Commun. Soil Sci. Plant Anal.* 18:1089–1109.
- Bollen, W.B. 1969. Properties of tree barks in relation to their agricultural utilization. USDA For. Serv. Res. Paper PNW 77.
- Broschat, T.K. 1995. Nitrate, phosphate, and potassium leaching from container-grown plants fertilized by several methods. *HortScience* 30:74–77.
- Browder, J.F. 2004. Sulfur requirements of container-grown pin oak and Japanese maple. Va. Polytechnic Inst., Blacksburg, MS thesis.
- Buamscha, M.G., J.E. Altland, D.M. Sullivan, and D.A. Horneck. 2007a. Micronutrient availability in fresh and aged Douglas fir bark. *HortScience* 42:152–156.
- Buamscha, G.M., J.E. Altland, D.M. Sullivan, D.A. Horneck, and J. Cassidy. 2007b. Chemical and physical properties of Douglas fir bark relevant to the production of container plants. *HortScience* 42:1281–1286.
- Bunt, A.C. 1988. Media and mixes for container grown plants. Allen and Unwin, Inc., Winchester, MA.
- Gavlak, R., D. Horneck, R. Miller, and J. Kotuby-Amacher. 2003. Soil, plant, and water reference methods for the western region. 2nd Ed. WCC-103 Publication, Fort Collins, CO.
- Handreck, K.A. 1986. Critical concentrations of sulfur in liquid feeds for plants in containers. *Scientia Hort.* 30:1–17.
- Handreck, K.A. 1990. Extractants for assessing the availability of copper to *Chrysanthemum morifolium* cultivar 'Yellow Mandalay' growing in soil-less media. *Scientia Hort.* 44:323–334.
- Handreck, K.A. 1994. Effect of pH on the uptake of cadmium, copper, and zinc from soilless media containing sewage sludge. *Commun. Soil Sci. Plant Anal.* 25:1913–1927.
- Lide D.R. (ed.). 2007. CRC handbook of chemistry and physics. Internet version 2007 (87th Ed.). Dec. 2007. <<http://www.hbcpnetbase.com>>. Taylor and Francis, Boca Raton, FL.
- Lucas, R.E. and J.F. Davis. 1961. Relationships between pH values of organic soils and availabilities of 12 plant nutrients. *Soil Sci.* 92:171–182.
- Niemiera, A.X. 1992. Micronutrient supply from pine bark and micronutrient fertilizers. *HortScience* 27:272.
- Niemiera, A.X. and R.D. Wright. 1986. Effect of liming rate on nitrification in a pine bark medium. *J. Amer. Soc. Hort. Sci.* 111:713–715.
- Ogden, R.J., F.A. Pokorny, H.A. Mills, and M.G. Dunavent. 1987. Elemental status of pine bark-based potting media. *Hort. Rev. (Amer. Soc. Hort. Sci.)* 9:103–131.
- Peterson, J.C. 1980. Effects of pH upon nutrient availability in a commercial soilless root medium utilized for floral crop production. *Ohio Agr. Res. Dev. Ctr. Circ.* 268:16–19.
- Rose, M.A. and H. Wang. 1999. Comparison of micronutrient sources for container Rhododendron. *HortTechnology* 9:220–224.
- Schabenberger, O. and F.J. Pierce. 2002. Contemporary statistical models for the plant and soil sciences. CRC Press, Boca Raton, FL.
- Warncke, D. 1998. Recommended test procedure for greenhouse growth media, p. 34–37. In: Dahnke, W.C. (ed.). Recommended chemical soil test procedures for the north central region. North Central Reg. Res. Pub. No. 221. Miss. Agr. Expt. Stat. SB 1001.
- Williams, B.J., J.C. Peterson, and J.D. Utzinger. 1988. Liming reaction in sphagnum peat-based growing media. *J. Amer. Soc. Hort. Sci.* 113:210–214.
- Wright, R.D. and L.E. Hinsley. 1991. Growth of containerized eastern redcedar amended with dolomitic limestone and micronutrients. *HortScience* 26:143–145.
- Wright, R.J. 1989. Soil aluminum toxicity and plant growth. *Commun. Soil Sci. Plant Anal.* 20:1479–1497.
- Yeager, T.H. and J.E. Barrett. 1985. Phosphorous and sulfur leaching from an incubated superphosphate-amended soilless container medium. *HortScience* 20:671–672.
- Yeager, T.H. and R.D. Wright. 1982. Pine bark-phosphorous relationships. *Commun. Soil Sci. Plant Anal.* 13:57–66.