

## Longleaf Pine Photosynthetic Response to Soil Resource Availability and Elevated Atmospheric Carbon Dioxide

G. B. Runion,\* R. J. Mitchell, T. H. Green, S. A. Prior, H. H. Rogers, and D. H. Gjerstad

### ABSTRACT

Gas exchange responses during a drought cycle were studied in longleaf pine (*Pinus palustris* Mill.) seedlings after prolonged exposure to varying levels of atmospheric CO<sub>2</sub> ( $\approx 365$  or  $\approx 730$   $\mu\text{mol CO}_2 \text{ mol}^{-1}$ ), soil N (40 or 400 kg N ha<sup>-1</sup> yr<sup>-1</sup>), and water (“adequate” and “stressed”). Elevated atmospheric CO<sub>2</sub> concentration increased photosynthesis, tended to decrease stomatal conductance, and increased water-use efficiency (WUE). Although soil resource availability influenced gas exchange measurements, it generally did not affect the magnitude or direction of the response to CO<sub>2</sub> concentration. However, significant interactions among treatment variables were observed for plant xylem pressure potential. In seedlings grown with high N, a positive growth response to elevated atmospheric CO<sub>2</sub> increased whole-plant water use resulting in more severe plant water stress, despite increased leaf-level WUE; however, under low N conditions the lack of a growth response to elevated CO<sub>2</sub> reduced whole-plant water use, decreased water stress severity, and increased WUE. Photosynthetic response to CO<sub>2</sub> was greatest in the high N treatment at the beginning of the drought cycle, but diminished as water stress increased; however, plants grown with low N showed greater photosynthetic responses to CO<sub>2</sub> later in the drought cycle. Therefore, plant gas exchange rates interact with growth response in determining the severity of water stress under drought and, thus, the ability of elevated atmospheric CO<sub>2</sub> to ameliorate the effects of drought and allow plants to maintain increased rates of photosynthesis may be influenced by the availability of other resources, such as N and water.

THE CONCENTRATION of CO<sub>2</sub> in the atmosphere is increasing at an unprecedented rate (Houghton et al., 1990) and plants will be directly affected by this rise

G.B. Runion, International Paper, 719 Southlands Road, Bainbridge, GA 31717; R.J. Mitchell, J.W. Jones Ecological Research Center, Route 2, Box 2324, Newton, GA 31770; T.H. Green, Dep. of Plant and Soil Science, P.O. Box 1208, Alabama A&M Univ., Normal, AL 35762; S.A. Prior and H.H. Rogers, USDA-ARS, National Soil Dynamics Lab., 411 S. Donahue Drive, Auburn, AL 36832; and D.H. Gjerstad, School of Forestry, 108 M. White Smith Hall, Auburn Univ., AL 36849. Received 22 Dec. 1997. \*Corresponding author (brett.runion@ipaper.com).

Published in *J. Environ. Qual.* 28:880–887 (1999).

in CO<sub>2</sub>. In general, studies have shown positive effects of CO<sub>2</sub> enrichment on plants including increased growth and WUE (Rogers and Dahlman, 1993; Allen and Amthor, 1995; Wittwer, 1995) and, although more variable, increased rates of photosynthesis (Chaves and Pereira, 1992; Amthor, 1995). However, the majority of CO<sub>2</sub> research has focused on crops under optimal growth conditions, rather than on native plants under limiting soil resources (Ceulemans and Mousseau, 1994; Amthor, 1995), despite the fact that natural terrestrial ecosystems are often limited by suboptimal levels of soil resource availability (e.g., N and water).

Understanding how forest species respond to this CO<sub>2</sub> buildup, and to potential associated climatic changes, is critical for predicting changes in stand structure and growth (McGuire and Joyce, 1995), which is particularly important due to the long-lived nature of trees and the economic and ecological values of forests. Coastal plain forests of the southeastern USA were once dominated by nearly pure stands of longleaf pine with a diverse understory plant community (Peet and Allard, 1993). The longleaf pine savanna ecosystem now occupies only 2% of its former range, a loss comparable to or exceeding that of most endangered communities throughout the world (Noss, 1989).

Longleaf pine forests currently occupy sites at the more xeric end of the moisture continuum and are often found on soils with low N availability; therefore, understanding how availability of differing soil resources affect plant response to CO<sub>2</sub> is also critical in predicting future productivity of southern pine forests. Prior et al. (1997) reported that a growth response to elevated atmospheric CO<sub>2</sub> was observed for longleaf pine growing under high, but not low, N conditions; in contrast, however, plants exposed to water stress showed a

**Abbreviations:** R/S, root to shoot ratio; WUE, water-use efficiency; P<sub>n</sub>, photosynthesis; g<sub>s</sub>, stomatal conductance; XPP, xylem pressure potential.

greater relative growth response to elevated  $\text{CO}_2$  than did their adequately-watered counterparts. This may suggest fundamentally different mechanisms by which availability of soil resources interact with atmospheric  $\text{CO}_2$  concentration to affect plant growth, which may be reflected by plant photosynthetic responses.

Soil fertility may influence plant photosynthetic response to  $\text{CO}_2$  via control of sink/source relationships and photosynthetic acclimation through time (Groninger et al., 1996). However, N may also interact with water in regulating photosynthetic response to  $\text{CO}_2$ . Nitrogen fertility may also influence plant WUE (Green and Mitchell, 1992), but in a fundamentally different way than elevated  $\text{CO}_2$ . Elevated  $\text{CO}_2$  generally increases photosynthesis ( $P_n$ ) and tends to reduce stomatal conductance ( $g_s$ ) which increases leaf-level WUE (Tolley and Strain, 1984; Wray and Strain, 1986; Hollinger, 1987) and suggests elevated  $\text{CO}_2$  may provide a mechanism for moderation of drought stress. High N increases WUE by increasing  $P_n$  disproportionately more than  $g_s$  (Green and Mitchell, 1992); thus, high N may or may not moderate plant response to drought through increased WUE. Furthermore, high N may increase leaf area and decrease root to shoot ratio (R/S) such that plants are predisposed to drought stress, while elevated  $\text{CO}_2$  tends to increase R/S and have little effect on leaf area in pines (Prior et al., 1997).

Despite the commonality of soil resource limitations in natural terrestrial ecosystems, particularly longleaf pine forests of the southeastern USA, few investigations have tested the extent to which photosynthetic response to  $\text{CO}_2$  is regulated by resource limitations or examined the potential for differences among varying soil resources (e.g., N vs. water). The objective of this study was to evaluate gas exchange responses to elevated  $\text{CO}_2$  for longleaf pine seedlings grown under two levels of both N and water availability and determine the potential for these soil resources to vary in the manner in which they regulate photosynthetic response.

## MATERIALS AND METHODS

One-year-old, nursery-grown longleaf pine seedlings from an open-pollinated seed source were graded to a uniform size and planted (three per container) in 45-L plastic containers filled with a coarse sandy medium (pH 5.1) of very low fertility (Cope et al., 1980). Seedlings were exposed to ambient ( $\approx 365 \mu\text{mol CO}_2 \text{ mol}^{-1}$ ) or elevated ( $\approx 730 \mu\text{mol CO}_2 \text{ mol}^{-1}$ ) atmospheric  $\text{CO}_2$  beginning 30 Mar. 1993 in an open top chamber system (Rogers et al., 1983b). Open top chambers were fitted with clear Teflon film covers to exclude rainfall. The chambers,  $\text{CO}_2$  supply, and  $\text{CO}_2$  monitoring/dispersing systems have been previously described for this study site (Mitchell et al., 1995). Nitrogen treatments, slightly modified from those described by Bazzaz and Miao (1993), consisted of either 40 or 400 kg N  $\text{ha}^{-1} \text{yr}^{-1}$  applied at 3-mo intervals (sulfur coated urea; 38-0-0) initiated at planting. Foliar N levels resulting from these treatments have been previously reported (Runion et al., 1997). Other nutrients were maintained at nonlimiting levels (Prior et al., 1997). Two water stress treatments (target maximum values of  $-0.5$  and  $-1.5$  MPa pre-dawn xylem pressure potential (XPP) for each drought cycle) were initiated after a 19-wk establishment period. At initiation of water-stress treatments, all containers were flooded with water, allowed to drain overnight, and weighed; these weights were

taken as field capacity values. Gravimetric measurements were correlated with pre-dawn XPP, measured on excised needles (one per treatment per chamber) at each weighing for 1 mo using a pressure bomb (Scholander et al., 1965). Individual containers received enough water to return them to field capacity weight when gravimetric measurements indicated the appropriate stress level had been achieved; xylem pressure potentials continued to be taken a minimum of once a month as a check on gravimetric readings. New field capacity values were determined after the November 1993 harvest and midway through the second growing season (July 1994) to compensate for changes in plant mass; xylem pressure potential was measured at each weighing for 1 mo following determination of new field capacity weights. Average needle xylem pressure potentials immediately prior to watering were  $-0.6$  and  $-1.3$  MPa for seedlings in the adequately-watered and water-stressed treatments, respectively. Use of deionized water ensured that fertility treatments remained unaffected by water treatments. Treatments were arranged in a split-plot design;  $\text{CO}_2$  treatments (main plots) were randomly assigned to chambers within each of four replicates (total of eight chambers). Nitrogen and water stress treatments (subplots) were randomly assigned, in a 2 by 2 factorial design, to a total of 16 containers within each chamber. Container locations within each chamber were re-randomized monthly.

On 20 July 1994 (following 16 mo of  $\text{CO}_2$  exposure and 19 drought stress cycles) all containers were watered to field capacity; a water stress cycle was then initiated and continued until 5 August. Adequately-watered containers were watered four times (every 3–4 d) during this period; water-stressed containers received no water after 20 July. Gas exchange measurements were made twice weekly (total of 5) throughout this drought cycle on one randomly selected plant per treatment per chamber. Pre-dawn XPP was determined on excised needles (one per treatment per chamber) prior to each gas exchange measurement using a pressure bomb. Stomatal conductance ( $\text{mmol H}_2\text{O m}^{-2} \text{s}^{-1}$ ) and photosynthesis ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{s}^{-1}$ ) were measured as quickly as possible in the morning (on each date, measurements were concluded between 0900 and 1000 h EDT to avoid diurnal effects) using a CIRAS-1 portable infrared gas analyzer (PP Systems, Haverhill, MA). Measurement  $\text{CO}_2$  concentrations were equivalent to those in the open top chamber under assessment. Instantaneous leaf-level WUE ( $\mu\text{mol CO}_2 \text{ mmol}^{-1} \text{H}_2\text{O}$ ) was calculated.

One container from each treatment in each chamber was destructively harvested in November 1994, 20 mo after initiation of the study. Foliar surface area of harvested seedlings was determined photometrically using a LiCor LI-3100 area meter (Li-Cor, Lincoln, NE). Root to shoot ratio (R/S) was determined from dry weights (oven dried at  $55^\circ\text{C}$ ) of harvested seedlings. Destructive data were totaled for each seedling and averaged for each container prior to analysis.

Data were analyzed using the unstructured model in the repeated measures analysis with the Mixed procedure of the Statistical Analysis System (SAS Institute, 1996). Error terms appropriate to the split-plot design were used to test the significance of main effects and their interactions. Differences were considered significant at the  $P \leq 0.05$  level.

## RESULTS

Elevated atmospheric  $\text{CO}_2$  increased  $P_n$ , tended to reduce  $g_s$ , and, thus, dramatically increased WUE throughout the water stress cycle (Table 1); however, the reduction in  $g_s$  was significant only during the early phase of the water stress cycle. When averaged across the water stress cycle,  $P_n$  for pines grown with elevated  $\text{CO}_2$  was 46% higher than for those grown in ambient

**Table 1. Main treatment effects for photosynthesis, stomatal conductance, water-use efficiency, and xylem pressure potential of longleaf pine seedlings through a drought stress cycle.**

Variable† DOY‡	CO <sub>2</sub> Concentration			N Fertility regime			H <sub>2</sub> O Stress regime		
	365	720	Pr > F§	40	400	Pr > F	Adequate	Stress	Pr > F
	μmol mol <sup>-1</sup>			kg ha <sup>-1</sup> yr <sup>-1</sup>					
<b>P<sub>n</sub></b>									
203	6.18	8.17	0.046	5.94	8.41	<0.001	6.39	7.97	0.012
207	3.92	6.28	0.021	4.83	5.37	0.397	5.34	4.86	0.448
210	5.59	7.41	0.080	6.38	6.62	0.792	7.17	5.83	0.156
214	6.55	9.90	0.002	7.83	8.62	0.065	9.35	7.09	<0.001
217	4.39	7.06	0.004	6.28	5.18	0.112	7.25	4.21	<0.001
<b>g<sub>s</sub></b>									
203	105.2	75.2	0.022	104.2	76.3	0.031	82.8	97.6	0.231
207	98.2	64.5	0.007	101.6	61.2	0.002	82.8	80.0	0.800
210	76.8	66.1	0.442	91.6	51.2	0.010	84.7	58.1	0.071
214	74.6	72.1	0.666	92.3	54.4	<0.001	94.2	52.5	<0.001
217	53.8	45.4	0.525	70.2	29.0	0.004	72.5	26.8	0.002
<b>WUE</b>									
203	2.47	4.83	<0.001	2.44	4.86	<0.001	3.59	3.71	0.621
207	1.80	4.26	<0.001	2.08	3.98	<0.001	2.81	3.24	0.086
210	2.91	4.95	<0.001	2.96	4.90	<0.001	3.67	4.19	0.260
214	2.66	4.43	<0.001	2.47	4.62	<0.001	3.13	3.96	0.002
217	2.24	3.76	<0.001	2.39	3.61	<0.001	3.04	2.97	0.789
<b>XPP</b>									
203	-0.49	-0.47	0.936	-0.48	-0.48	0.880	-0.44	-0.52	0.020
207	-0.68	-0.69	0.952	-0.55	-0.83	0.006	-0.52	-0.86	0.001
210	-0.80	-0.91	0.687	-0.72	-0.99	0.017	-0.54	-1.17	<0.001
214	-0.84	-0.92	0.757	-0.79	-0.97	0.063	-0.61	-1.15	<0.001
217	-0.92	-1.09	0.635	-0.87	-1.14	0.008	-0.60	-1.41	<0.001


† Variables: P<sub>n</sub> = photosynthesis (μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>); g<sub>s</sub> = stomatal conductance (mmol H<sub>2</sub>O m<sup>-2</sup> s<sup>-1</sup>); WUE = water use efficiency (μmol CO<sub>2</sub> mmol H<sub>2</sub>O<sup>-1</sup>); XPP = xylem pressure potential (MPa); levels for the water stress regimes refer to target maximum xylem pressure potential values of -0.5 and -1.5 MPa for Adequate and Stress, respectively.

‡ DOY's (day of year) correspond to sample dates of 22, 26, 29, July and 2 and 5, August, respectively.

§ Probability of a greater *F* value calculated, using appropriate error terms for the split-plot design, under Mixed procedure of SAS. Values are from data averaged across a drought stress cycle.

CO<sub>2</sub> ( $P = 0.009$ ), g<sub>s</sub> was reduced from 81.7 to 64.7 mmol H<sub>2</sub>O m<sup>-2</sup> s<sup>-1</sup> ( $P = 0.072$ ), and WUE increased by more than a factor of 1.8 ( $P < 0.001$ ). Soil resource availability tended to influence gas exchange measurements but not the magnitude or direction of the response to CO<sub>2</sub> concentration (Fig. 1 and 2), thus, interactions of CO<sub>2</sub>

with other treatment variables were generally not significant. However, a significant CO<sub>2</sub> × N interaction for WUE was observed wherein the magnitude of increase due to elevated CO<sub>2</sub> was greater for plants grown in high, relative to low, N (Fig. 1). A significant CO<sub>2</sub> × water interaction for P<sub>n</sub> was also observed to be due to



**Fig. 1. Interactions of CO<sub>2</sub> with soil nitrogen (N) on photosynthesis (P<sub>n</sub>), stomatal conductance (g<sub>s</sub>), water-use efficiency (WUE), and xylem pressure potential (XPP). Data are averaged across five measurements made during a drought stress cycle. *P* values located above each set of bars refer to differences between CO<sub>2</sub> treatments.**




Fig. 2. Interactions of CO<sub>2</sub> with plant water stress on photosynthesis (P<sub>n</sub>), stomatal conductance (g<sub>s</sub>), water-use efficiency (WUE), and xylem pressure potential (XPP). Data are averaged across five measurements made during a drought stress cycle. P values located above each set of bars refer to differences between CO<sub>2</sub> treatments.

a difference in magnitude, that is, the increase due to elevated CO<sub>2</sub> was greater for plants grown with adequate water, relative to water stress (Fig. 2). Significant N × water and CO<sub>2</sub> × N × water interactions were also observed for XPP. These results are taken from a single water stress cycle within a long-term study and plants were not of equal size at initiation of the drought cycle; therefore, XPP responses are the result of both direct and indirect (those mediated through long-term changes in plant growth) effects of treatments and their interactions. In the water stress treatment, XPP was decreased in high N plants, possibly due to increases in leaf area (water demand) or decreases in R/S (water supply). In addition, the magnitude of the increase in leaf area or decrease in R/S was greater for plants grown under elevated, compared with ambient, CO<sub>2</sub> which resulted in a significantly lower XPP for high N plants under elevated CO<sub>2</sub> when subjected to water stress (Table 2).

Under adequately-watered conditions, XPP was unaffected by CO<sub>2</sub> and N (Fig. 3); g<sub>s</sub> also was not significantly affected by atmospheric CO<sub>2</sub> concentration, although it tended to be numerically lower under elevated CO<sub>2</sub>

throughout the water stress cycle (Fig. 4). In contrast, P<sub>n</sub> (Fig. 3) and WUE (Fig. 4) were generally higher under elevated CO<sub>2</sub> for plants grown with adequate water. Effects of CO<sub>2</sub> on gas exchange measurements tended to occur regardless of N level; however, differences between CO<sub>2</sub> treatments were generally of a larger magnitude and more frequently significant for plants grown with the high level of soil N.

Under water-stressed conditions, high N plants had lower XPP (i.e., more stress) than low N plants when grown under elevated CO<sub>2</sub>; for plants grown with ambient CO<sub>2</sub>, N treatment level did not affect XPP (Fig. 3). Although CO<sub>2</sub> concentration significantly affected XPP only in high N plants at the final measurement, the pattern of response to CO<sub>2</sub> varied dramatically between the two N treatments. In plants grown with high N, XPP was lower under elevated CO<sub>2</sub>, but XPP for low N plants was consistently higher under elevated CO<sub>2</sub> indicating that low N plants were never stressed to the extent of high N plants, particularly under elevated CO<sub>2</sub>. This differential pattern of XPP response in the water stress treatment influenced P<sub>n</sub> throughout the drought cycle

Table 2. Effects of treatment variables† on xylem pressure potential (XPP) of longleaf pine seedlings taken during a drought stress cycle after 16 mo of CO<sub>2</sub> exposure and on leaf area (LA) and root to shoot ratio (R/S) from plants destructively harvested after 20 mo of CO<sub>2</sub> exposure.

Soil N	Water stress	CO <sub>2</sub>	XPP	Pr > T‡	LA (m <sup>2</sup> )	Pr > T	R/S	Pr > T
40	Adequate	365	-0.55		0.41		1.20	
400	Adequate	365	-0.53	0.879	1.73	<0.001	0.81	0.004
40	Stress	365	-0.89		0.26		1.01	
400	Stress	365	-1.02	0.261	0.60	0.008	0.74	0.041
40	Adequate	730	-0.57		0.23		1.08	
400	Adequate	730	-0.52	0.637	1.67	<0.001	0.90	0.147
40	Stress	730	-0.72		0.19		1.39	
400	Stress	730	-1.46	<0.001	1.05	<0.001	1.08	0.021

† Treatment variables are: Soil N = soil nitrogen in kg ha<sup>-1</sup> yr<sup>-1</sup>; water stress levels refer to target maximum xylem pressure potential values of -0.05 and -1.5 MPa for Adequate and Stress, respectively; and atmospheric CO<sub>2</sub> concentration in µmol CO<sub>2</sub> mol<sup>-1</sup>. Data for XPP are averaged across five measurements made during a single drought stress cycle.

‡ Probability of a greater value for the T statistic calculated, with appropriate error terms for the split-plot design, using the unstructured model of the repeated measures analysis under the Mixed procedure of SAS for the difference between N treatments.






Fig. 3. Photosynthesis ( $P_n$ ) and pre-dawn xylem pressure potential (XPP) for longleaf pine seedlings at five measurement dates through a drought cycle as affected by interactions between  $\text{CO}_2$  and soil N. Asterisks above the graphed lines indicate significant differences ( $P \leq 0.05$ ) between  $\text{CO}_2$  treatments under high N (comparing filled symbols), while those below the lines indicated differences under low N (comparing open symbols). Elevated  $\text{CO}_2$  tended to increase  $P_n$  in high N plants only in the absence of water stress, while differences for low N plants became significant as stress increased. Elevated  $\text{CO}_2$  resulted in lower XPP for stressed plants grown with high N throughout most of the drought cycle, while this reduction was not observed for low N plants.

(Fig. 3). At the beginning of the drought cycle (i.e., no water stress), higher  $P_n$  in response to  $\text{CO}_2$  addition was greatest in the high N treatment; however, as water stress increased, differences between  $\text{CO}_2$  treatments decreased. In contrast, plants grown with low N showed increased  $P_n$  in response to added  $\text{CO}_2$  later in the drought cycle. For plants under water stress, elevated  $\text{CO}_2$  reduced  $g_s$  early in the cycle in both N treatments (Fig. 4). Similar to plants under adequately-watered conditions, elevated  $\text{CO}_2$  tended to increase WUE regardless of N conditions for plants under water stress and differences between  $\text{CO}_2$  treatments were larger for plants grown with high N. However, in contrast to the adequately-watered treatment, this increased WUE under elevated  $\text{CO}_2$  diminished at the end of the water stress cycle for plants grown with high, but not low, N (Fig. 4).

## DISCUSSION

Woodward (1992) states that the best documented and repeatable response to atmospheric  $\text{CO}_2$  enrichment is a significant increase in photosynthesis of  $\text{C}_3$  plants, a fact readily apparent from the  $\text{CO}_2$  literature (Rogers et al., 1983a; Wittwer, 1985; Long and Drake, 1992; Allen and Amthor, 1995; Amthor, 1995; Eamus et al., 1995; Tissue et al., 1995). In this experiment, longleaf pine seedlings were exposed to twice ambient

concentrations of  $\text{CO}_2$  for 16 mo and, in the high resource treatment (e.g., high N and adequate water),  $P_n$  increased 56% under elevated  $\text{CO}_2$  which is similar to increases reported for other tree species (see reviews by Eamus and Jarvis, 1989; Long and Drake, 1992; Mousseau and Saugier, 1992; Ceulemans and Mousseau, 1994).

Research has shown that elevated  $\text{CO}_2$  interacts with soil resource availability to influence plant  $P_n$ ; however, the manner in which these resources interact with  $\text{CO}_2$  may differ. In general, plants grown under ambient  $\text{CO}_2$  decrease  $P_n$  when placed under water stress, while elevated  $\text{CO}_2$  alleviates this stress (to some degree), allowing plants to maintain higher  $P_n$  (Huber et al., 1984; Conroy et al., 1986; Liang and Maruyama, 1995). Even in cases where water stress was severe enough to reduce  $P_n$  of elevated  $\text{CO}_2$ -grown plants, high  $\text{CO}_2$ /water-stressed plants maintained  $P_n$  rates equivalent to ambient  $\text{CO}_2$ /adequately-watered plants, again suggesting amelioration of water stress (Groninger et al., 1996). In contrast, elevated  $\text{CO}_2$  may not alleviate nutrient stress and both P (Conroy et al., 1986) and N (Tissue et al., 1993; Curtis et al., 1994) limitations have been reported to dampen or negate elevated  $\text{CO}_2$ -induced increases in  $P_n$ . However, some researchers have observed increased  $P_n$  under high  $\text{CO}_2$  even under conditions of low fertility (Norby and O'Neill, 1991); although, this may be due in part to  $P_n$  acclimation induced by small pot size (Arp,




Fig. 4. Stomatal conductance ( $g_s$ ) and water-use efficiency (WUE) for longleaf pine seedlings at five measurement dates through a drought cycle as affected by interactions between  $\text{CO}_2$  and soil N. Use of asterisks is the same as in Fig. 3. Elevated  $\text{CO}_2$  tended to decrease  $g_s$  under water stress early in the stress cycle. Elevated  $\text{CO}_2$  tended to increase WUE regardless of N and water conditions, although differences between  $\text{CO}_2$  treatments tended to be larger for plants grown with high N. This increase diminished at the end of the drought cycle for plants grown with high N.

1991; Thomas and Strain, 1991). Groninger et al. (1996) found that  $P_n$  increased in sweetgum (*Liquidambar styraciflua* L.) but decreased in loblolly pine in response to high N; however, response of  $P_n$  to  $\text{CO}_2$  was not influenced by N in either species. We observed that longleaf pine  $P_n$  response to  $\text{CO}_2$  was also rarely influenced by either N or water stress (elevated  $\text{CO}_2$ -grown seedlings tended to have higher  $P_n$  under both N and water treatments); so, although soil resource availability influenced gas exchange measurements, it rarely had a significant effect on the magnitude or direction of the response to  $\text{CO}_2$  concentration. Curtis et al. (1994) suggested that, although nutrient or water stress clearly influences the magnitude of  $\text{CO}_2$  response, the stimulation of  $P_n$  by elevated  $\text{CO}_2$  is rarely fully eliminated; results from this study support this suggestion. However, if N is insufficient for growth (Prior et al., 1997), additional C from increased  $P_n$  may accumulate and disrupt needle chloroplast integrity (Pritchard et al., 1997) and, thus, negatively impact plant health.

Decreased  $g_s$  under high  $\text{CO}_2$  (Rogers et al., 1983a; Morison, 1985; Eamus, 1991; Allen and Amthor, 1995) implies decreased water use, and possible amelioration of drought stress, which may become increasingly important under future climate change scenarios. However, increases in leaf area (i.e., under high N) might compensate for reduced leaf-level  $g_s$  and result in similar or greater, whole-plant water use. Therefore, this poten-

tial for alleviation of drought stress via decreased  $g_s$  will depend on plant growth response and on the severity and length of the stress. It has been suggested (Murray, 1995) that, while effects of elevated  $\text{CO}_2$  on stomata can override the effects of light, water stress triggers stomatal closure and can override both  $\text{CO}_2$  and light. This is supported by the present study in that elevated  $\text{CO}_2$  significantly decreased  $g_s$  until the middle of the water stress cycle, which was the same point at which the effect of water stress became significant.

Implications of altered  $g_s$ , in combination with increased  $P_n$ , are also of interest to plant scientists due to potential impacts on WUE. Increased instantaneous, leaf-level WUE under elevated  $\text{CO}_2$ , as we observed, has been commonly reported (Rogers et al., 1983a; Morison, 1985; Eamus and Jarvis, 1989; Norby and O'Neill, 1991; Liang and Maruyama, 1995); however, it has been suggested that drought stress and nutrient deficiency are likely to moderate this response (Mousseau and Saugier, 1992). Although previously reported  $\text{CO}_2$ -induced increases in WUE under high fertility (Norby and O'Neill, 1991) and water stress (Liang and Maruyama, 1995) were observed, the fact that interactions of these resources with  $\text{CO}_2$  on WUE tended to be of magnitude rather than rank suggests their moderating effect on WUE response to  $\text{CO}_2$  may be minimal.

In this study, high N as well as elevated  $\text{CO}_2$  increased WUE of longleaf pine seedlings and we suggested that

these increases may occur in fundamentally different ways (i.e., that elevated CO<sub>2</sub> increases WUE by increasing P<sub>n</sub> and reducing g<sub>s</sub>, while high N increases WUE by increasing P<sub>n</sub> disproportionately more than g<sub>s</sub>); this is not supported by our data. High N did increase P<sub>n</sub> (particularly under adequate water), but also tended to decrease g<sub>s</sub>; g<sub>s</sub> was only influenced by elevated CO<sub>2</sub> early in the water stress cycle, while CO<sub>2</sub> had a much stronger effect on P<sub>n</sub> (particularly under adequate water). However, the contention that elevated CO<sub>2</sub> ameliorates the effects of drought (Bazzaz, 1990) and allow plants to maintain increased P<sub>n</sub>, while high N does not, may still be valid. Increased N availability resulted in decreased XPP of water stressed plants, indicating an exacerbation of water stress. This increased stress was most likely due to the fact that growth of fertilized seedlings was sufficient to increase whole-plant water use. The effect of elevated atmospheric CO<sub>2</sub> concentration on water stress intensity was dependent on N availability. In seedlings grown with high N, a positive growth response to elevated CO<sub>2</sub> increased whole-plant water use resulting in more severe water stress, despite increased WUE. However, under low N conditions, the lack of a growth response to elevated CO<sub>2</sub> (Prior et al., 1997) combined with somewhat lower g<sub>s</sub>, reduced whole-plant water use and decreased water-stress severity.

Data from this study indicate that leaf-level g<sub>s</sub> interacts with growth response in determining the severity of water stress under drought conditions. Other things remaining constant, decreases in g<sub>s</sub> with elevated CO<sub>2</sub> will ameliorate water stress severity and allow increased P<sub>n</sub>; however, increased growth can easily overcome these positive effects. When growth responses to increased resource (N and/or CO<sub>2</sub>) availability occur, more water is used by the larger plant and more severe stress can occur, assuming no changes in resource availability or efficiency of acquisition. Therefore, our data indicate that, while increased P<sub>n</sub> and WUE may be maintained under low-N environments (such as those that longleaf pine normally inhabit), this might not be the case for longleaf pine growing under high N availability, when growth response to elevated atmospheric CO<sub>2</sub> may increase water stress due to greater whole-plant water use. However, plants grown in a more natural environment (as opposed to containers) might have a larger root system and, thus, increased access to soil water so that the increased water stress observed here would likely be diminished or delayed until water became more limiting; nonetheless, when water becomes limiting, larger plants will experience increased stress. Plants grown under high fertility tend to allocate less growth belowground, while those grown under elevated CO<sub>2</sub> tend to increase belowground allocation (Prior et al., 1997; Runion et al., 1997). Therefore, although the exact nature of the relationship of growth response with P<sub>n</sub> and WUE for plants in the field may vary from that observed in this container study, it will, again, depend on interactions of elevated CO<sub>2</sub> with soil resource availability.

Increased P<sub>n</sub> of longleaf pine seedlings under varying levels of both N and water may reflect the plasticity of

this species observed in nature and, when coupled with other potential benefits (e.g., resistance to insects, disease, and fire), may make longleaf pine an attractive species given rising CO<sub>2</sub> and changing climate. However, to predict plant responses to future climate change, leaf-level measurements of P<sub>n</sub>, g<sub>s</sub>, and WUE must be scaled up to the whole plant and canopy to take into account considerations of differences in plant size and, thus, the importance of reported CO<sub>2</sub> effects may be overestimated (Morison, 1985; Eamus, 1991). Although data from the present study do not address large trees or mature forests, they may be relevant for seedling establishment during regeneration which is of critical importance to forest health and productivity.

#### ACKNOWLEDGMENTS

This material is based upon work supported through the Southeastern Regional Center of the National Institute for Global Environmental Change by the U.S. Department of Energy under Cooperative Agreement DE-FC03-90ER61010 and through the Experimental Program to Stimulate Competitive Research by the U.S. Environmental Protection Agency under contract R821826-01-1. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the U.S. Department of Energy or the U.S. Environmental Protection Agency. The authors wish to thank Ms. Deborah Boykin, USDA-ARS, Stoneville, MS, for advice and assistance with statistical analyses, Ms. Trina Cagle-Cleveland for technical assistance during data collection, and Drs. William A. Carey, James L.J. Houppis, and Jeffrey L. Sibley for critical reviews of the manuscript.

#### REFERENCES

- Allen, L.H., Jr., and J.S. Amthor. 1995. Plant physiological responses to elevated CO<sub>2</sub>, temperature, air pollution, and UV-B radiation. p. 51-84. In G.M. Woodwell and F.T. Mackenzie (ed.) Biotic feedbacks in the global climatic system: Will the warming feed the warming? Oxford University Press, New York.
- Amthor, J.S. 1995. Terrestrial higher-plant response to increasing atmospheric [CO<sub>2</sub>] in relation to the global carbon cycle. *Global Change Biol.* 1:243-274.
- Arp, W.J. 1991. Effects of source-sink relations on photosynthetic acclimation to elevated CO<sub>2</sub>. *Plant Cell Environ.* 14:869-875.
- Bazzaz, F.A. 1990. The response of natural ecosystems to the rising global CO<sub>2</sub> levels. *Ann. Rev. Ecol. Syst.* 21:167-196.
- Bazzaz, F.A., and S.L. Miao. 1993. Successional status, seed size, and responses of tree seedlings to CO<sub>2</sub>, light, and nutrients. *Ecology* 74:104-112.
- Ceulemans, R., and M. Mousseau. 1994. Effects of elevated atmospheric CO<sub>2</sub> on woody plants. *New Phytol.* 127:425-446.
- Chaves, M.M., and J.S. Pereira. 1992. Water stress, CO<sub>2</sub> and climate change. *J. Exp. Bot.* 43:1131-1139.
- Conroy, J.P., R.M. Smillie, M. Küppers, D.I. Bevege, and E.W. Barlow. 1986. Chlorophyll *a* fluorescence and photosynthetic and growth responses of *Pinus radiata* to phosphorus deficiency, drought stress and high CO<sub>2</sub>. *Plant Physiol.* 81:423-429.
- Cope, J.T. Jr., C.E. Evans, and H.C. Williams. 1980. Soil test fertilizer recommendations for Alabama crops. Circular 251. Auburn University Agric. Exp. Stn., Auburn, AL.
- Curtis, P.S., D.R. Zak, K.S. Pregitzer, and J.A. Teeri. 1994. Above- and belowground response of *Populus grandidentata* to elevated atmospheric CO<sub>2</sub> and soil N availability. *Plant Soil* 165:45-51.
- Eamus, D. 1991. The interaction of rising CO<sub>2</sub> and temperatures with water use efficiency. *Plant Cell Environ.* 14:843-852.
- Eamus, D., and P.G. Jarvis. 1989. The direct effects of increase in the global atmospheric CO<sub>2</sub> concentration on natural and commercial temperate trees and forests. *Adv. Ecol. Res.* 19:1-55.

- Eamus, D., G.A. Duff, and C.A. Berryman. 1995. Photosynthetic responses to temperature, light flux-density, CO<sub>2</sub> concentration and vapour pressure deficit in *Eucalyptus tetradonta* grown under CO<sub>2</sub> enrichment. *Environ. Pollut.* 90:41-49.
- Green, T.H., and R.J. Mitchell. 1992. Effects of nitrogen on the response of loblolly pine to water stress. I. Photosynthesis and stomatal conductance. *New Phytol.* 122:627-633.
- Groninger, J.W., J.R. Seiler, S.M. Zedaker, and P.C. Berang. 1996. Photosynthetic response of loblolly pine and sweetgum seedlings stands to elevated carbon dioxide, water stress, and nitrogen level. *Can. J. For. Res.* 26:95-102.
- Hollinger, D.Y. 1987. Gas exchange and dry matter allocation responses to elevation of atmospheric CO<sub>2</sub> concentration in seedlings of three tree species. *Tree Physiol.* 3:193-202.
- Houghton, J.T., G.J. Jenkins, and J.J. Ephraums (ed.). 1990. Climate change, the IPCC scientific assessment. Cambridge Univ. Press, Cambridge, UK.
- Huber, S.C., H.H. Rogers, and F.L. Mowry. 1984. Effects of water stress on photosynthesis and carbon partitioning in soybean (*Glycine max* [L.] Merr.) plants grown in the field at different CO<sub>2</sub> levels. *Plant Physiol.* 76:244-249.
- Liang, N., and K. Maruyama. 1995. Interactive effects of CO<sub>2</sub> enrichment and drought stress on gas exchange and water-use efficiency in *Alnus firma*. *Environ. Exp. Bot.* 35:353-361.
- Long, S.P., and B.G. Drake. 1992. Photosynthetic CO<sub>2</sub> assimilation and rising atmospheric CO<sub>2</sub> concentrations. p. 69-107. *In* N.R. Baker and H. Thomas (ed.) *Crop photosynthesis: Spatial and temporal determinants*. Elsevier, New York.
- McGuire, A.D., and L.A. Joyce. 1995. Responses of net primary productivity to changes in CO<sub>2</sub> and climate. p. 9-45. *In* L.A. Joyce (ed.) *Productivity of America's forests and climate change*. General Tech. Rep. RM-271. USDA Forest Service, Rocky Mountain Forest and Range Exp. Stn., Fort Collins, CO.
- Mitchell, R.J., G.B. Runion, S.A. Prior, H.H. Rogers, J.S. Amthor, and F.P. Henning. 1995. Effects of nitrogen on *Pinus palustris* foliar respiratory responses to elevated atmospheric CO<sub>2</sub> concentration. *J. Exp. Bot.* 46:1561-1567.
- Morison, J.I.L. 1985. Sensitivity of stomata and water use efficiency to high CO<sub>2</sub>. *Plant Cell Environ.* 8:467-474.
- Mousseau, M., and B. Saugier. 1992. The direct effect of increased CO<sub>2</sub> on gas exchange and growth of forest tree species. *J. Exp. Bot.* 43:1121-1130.
- Murray, D.R. 1995. Plant responses to carbon dioxide. *Am. J. Bot.* 82:690-697.
- Norby, R.J., and E.G. O'Neill. 1991. Leaf area compensation and nutrient interactions in CO<sub>2</sub>-enriched seedlings of yellow-poplar (*Liriodendron tulipifera* L.). *New Phytol.* 117:515-528.
- Noss, R.F. 1989. Longleaf pine and wiregrass: Keystone components of an endangered ecosystem. *Nat. Areas J.* 9:211-213.
- Peet, R.K., and D.J. Allard. 1993. Longleaf pine vegetation of the southern Atlantic and eastern gulf coast regions: A preliminary classification. p. 45-81. *In* S.M. Hermann (ed.) *Proc. of the Tall Timbers Fire Ecology Conf.*, no. 18, The Longleaf Pine ecosystem: Ecology, restoration and management. Tall Timbers Res. Stn., Tallahassee, FL.
- Prior, S.A., G.B. Runion, R.J. Mitchell, H.H. Rogers, and J.S. Amthor. 1997. Effects of atmospheric CO<sub>2</sub> on longleaf pine: Productivity and allocation as influenced by nitrogen and water. *Tree Physiol.* 17:397-405.
- Pritchard, S.G., C.M. Peterson, S.A. Prior, and H.H. Rogers. 1997. Elevated atmospheric CO<sub>2</sub> differentially affects needle chloroplast ultrastructure and phloem anatomy in *Pinus palustris*: Interactions with soil resource availability. *Plant Cell Environ.* 20:461-471.
- Rogers, H.H., G.E. Bingham, J.D. Cure, J.M. Smith, and K.A. Surano. 1983a. Responses of selected plant species to elevated carbon dioxide in the field. *J. Environ. Qual.* 12:569-574.
- Rogers, H.H., and R.C. Dahlman. 1993. Crop responses to CO<sub>2</sub> enrichment. *Vegetatio* 104/105:117-131.
- Rogers, H.H., W.W. Heck, and A.S. Heagle. 1983b. A field technique for the study of plant responses to elevated carbon dioxide concentrations. *Air Pollut. Control Assoc. J.* 33:42-44.
- Runion, G.B., R.J. Mitchell, H.H. Rogers, S.A. Prior, and T.K. Counts. 1997. Effects of nitrogen and water limitation and elevated atmospheric CO<sub>2</sub> on ectomycorrhiza of longleaf pine. *New Phytol.* 137:681-689.
- SAS Institute. 1996. SAS system for mixed models. SAS Inst., Cary, NC.
- Scholander, P.F., H.T. Hammel, E.D. Bradstreet, and E.A. Hemmingsen. 1965. Sap pressure in vascular plants. *Science* (Washington, DC) 148:339-346.
- Thomas, R.B., and B.R. Strain. 1991. Root restriction as a factor in photosynthetic acclimation of cotton seedlings grown in elevated carbon dioxide. *Plant Physiol.* 96:627-634.
- Tissue, D.T., K.L. Griffin, R.B. Thomas, and B.R. Strain. 1995. Effects of low and elevated CO<sub>2</sub> on C<sub>3</sub> and C<sub>4</sub> annuals. II. Photosynthesis and leaf biochemistry. *Oecologia* 101:21-28.
- Tissue, D.T., R.B. Thomas, and B.R. Strain. 1993. Long-term effects of elevated CO<sub>2</sub> and nutrients on photosynthesis and rubisco in loblolly pine seedlings. *Plant Cell Environ.* 16:859-865.
- Tolley, L.C., and B.R. Strain. 1984. Effects of CO<sub>2</sub> enrichment and water stress on growth of *Liquidambar styraciflua* and *Pinus taeda* seedlings. *Can. J. Bot.* 62:2135-2139.
- Wittwer, S.H. 1985. Carbon dioxide levels in the biosphere: Effects on plant productivity. *CRC Crit. Rev. Plant Sci.* 2:171-198. CRC Press, Boca Raton, FL.
- Wittwer, S.H. 1995. Food, climate, and carbon dioxide: The global environment and world food production. CRC Press, Boca Raton, FL.
- Woodward, F.I. 1992. Tansley Review no. 41 Predicting plant responses to global environmental change. *New Phytol.* 122:239-251.
- Wray, S.M., and B.R. Strain. 1986. Response of two old field perennials to interactions of CO<sub>2</sub> enrichment and drought stress. *Am. J. Bot.* 73:1486-1491.