

# Prevalence of Self-Reported Nutrition-Related Health Problems in the Lower Mississippi Delta

## ABSTRACT

**Objectives.** The purpose of this study was to assess demographic and geographic differences in prevalence of self-reported nutrition-related health problems in Arkansas, Louisiana, and Mississippi.

**Methods.** The authors analyzed 1991 and 1993 Behavioral Risk Factor Surveillance System data for adults 18 years or older.

**Results.** Less educated African American women and women of other minority groups who were aged 35 to 64 years reported the highest prevalence of health problems. Geographic differences involved prevalence of hypertension, health status, and insurance status.

**Conclusions.** Specific demographic subgroups and geographic areas with a high risk of health problems are in particular need of targeted interventions. (*Am J Public Health*. 1999;89:1418-1421)

Juneal Smith, PhD, MPH, RD, Shelly Lensing, MS, Jacqueline A. Horton, ScD, Jennifer Lovejoy, PhD, Sahar Zaghoul, MD, PhD, Ivis Forrester, PhD, RD, Bernestine B. McGee, PhD, RD, and Margaret L. Bogle, PhD, RD

The 3 states that compose the Lower Mississippi Delta region—Arkansas, Louisiana, and Mississippi—rank among the nation's 5 poorest states.<sup>1</sup> The region is predominantly rural, and minorities (primarily African American) make up 37%, 33%, and 17% of the populations of Mississippi, Louisiana, and Arkansas, respectively.<sup>2</sup> While 20% to 26% of the area's residents have incomes below the federal poverty level, 56% of African American households have incomes below the poverty level.<sup>3</sup>

Others have documented the relationship between unhealthy dietary behaviors and chronic conditions.<sup>4,5</sup> Although it is well established that chronic health conditions are more prevalent in the Lower Mississippi Delta region than in the rest of the country, few studies have examined differences in prevalence among demographic and geographic subsets of the population.<sup>1,6</sup> The purpose of this study was to compare geographic regions and to identify subpopulations at higher risk for chronic nutrition-related disorders as an essential prelude to developing appropriate intervention strategies.

## Methods

The Behavioral Risk Factor Surveillance System (BRFSS) is an annual random-digit-dialed telephone survey of noninstitutionalized civilian adults 18 years or older that is conducted by state health departments in collaboration with the Centers for Disease Control and Prevention (CDC).<sup>7,8</sup> The BRFSS uses a multistage cluster sampling technique based on the Waksberg method.<sup>8,9</sup> The present analysis used Arkansas, Louisiana, and Mississippi data from 4586 respondents in 1991 and 5001 respondents in 1993 (Arkansas did not participate in 1992).

In 1994, Congress established the Lower Mississippi Delta Nutrition Intervention Research Initiative ("Delta NIRI"), a consortium of 6 academic institutions in the region (Alcorn State University [Lorman, Miss], Arkansas Children's Hospital Research Institute [Little Rock], Pennington Biomedical Research Center [Baton Rouge, La], Southern University and A&M College [Baton Rouge], University of Arkansas at Pine Bluff, Univer-

sity of Southern Mississippi [Hattiesburg]), a coordinating center (Westat, Inc [Rockville, Md]), and the Agricultural Research Service of the US Department of Agriculture, to conduct nutrition intervention research in the Lower Mississippi Delta region.<sup>1</sup> Of the 222 counties-parishes in Arkansas, Louisiana, and Mississippi, 36 were selected, on the basis of poverty and geographic criteria, as Delta Initiative counties for intervention research.<sup>10</sup> In the 1991 and 1993 versions of the BRFSS, there were 827 respondents from Delta NIRI counties-parishes.

Self-reported weight and height were used to derive body mass indexes (BMIs). Obesity was defined as a BMI of 27.3 kg/m<sup>2</sup> or higher for women and a BMI of 27.8 kg/m<sup>2</sup> or higher for men.<sup>8,11</sup> Respondents were asked whether (1) they had ever been told by a health professional that they had diabetes, high blood pressure, or high cholesterol; (2) they had ever had their cholesterol tested; (3) they had had their blood pressure taken in the previous year; and (4) they had health insurance. In 1993, respondents rated their health on a 5-point scale (excellent, very good, good, fair, or poor).

Prevalence rates and 95% confidence intervals were calculated, and *t* tests were used to compare proportions; analyses used Stata 5.0<sup>12,13</sup> and were based on large sample methods for weighted survey data. BRFSS data were weighted by the CDC to reflect the

Juneal Smith and Shelly Lensing are with Arkansas Children's Hospital Research Institute, Little Rock. Jacqueline A. Horton is with Westat, Inc, Rockville, Md. Jennifer Lovejoy is with the Pennington Biomedical Research Center, Baton Rouge, La. Sahar Zaghoul is with the Department of Human Sciences, University of Arkansas, Pine Bluff. Ivis Forrester is with the Department of Family and Consumer Sciences, Alcorn State University, Lorman, Miss. Bernestine B. McGee is with the Human Nutrition and Food Program, Southern University and A&M College, Baton Rouge, La. Margaret L. Bogle is with the US Dept of Agriculture, Agricultural Research Service, Little Rock.

Requests for reprints should be sent to Shelly Lensing, MS, Department of Pediatrics, University of Arkansas for Medical Sciences, Arkansas Children's Hospital, 800 Marshall St, Slot 512-26, Little Rock, AR 72202 (e-mail: slensing@care.ach.uams.edu).

This paper was accepted February 24, 1999.

**TABLE 1—Demographic Characteristics, Prevalence of Chronic Conditions, and Health Behaviors and Perceptions in Delta NIRI and non-Delta NIRI Counties—Parishes and the Rest of the United States: Behavioral Risk Factor Surveillance System, 1991 and 1993**

	Delta NIRI Counties, % (95% CI)	Non-Delta NIRI Counties, % (95% CI)	P <sup>a</sup>	Other States, <sup>p</sup> % (95% CI)
<b>Demographic characteristics</b>				
Female	53.9 (49.8, 58.1)	53.1 (52.0, 54.3)	.706	52.1 (51.7, 52.4)
African American—other minority <sup>c</sup>	44.5 (39.9, 49.0)	25.0 (23.8, 26.2)	<.001	14.4 (14.1, 14.7)
Less than high school education	34.8 (31.2, 38.5)	21.1 (20.1, 22.0)	<.001	15.0 (14.7, 15.3)
Aged 65+ years	19.9 (17.0, 22.8)	17.2 (16.3, 18.1)	.087	16.9 (16.7, 17.2)
<b>Chronic conditions</b>				
Obesity	31.3 (27.7, 34.9)	28.5 (27.4, 29.5)	.138	25.1 (24.8, 25.4)
Diabetes	6.9 (5.1, 8.8)	5.7 (5.2, 6.2)	.217	4.8 (4.7, 5.0)
High blood pressure	28.9 (25.4, 32.4)	24.3 (23.3, 25.3)	.014	21.3 (21.0, 21.6)
High cholesterol	28.0 (23.7, 32.2)	27.2 (25.9, 28.5)	.724	27.4 (27.0, 27.8)
<b>Behavioral characteristics</b>				
Tested for cholesterol	57.9 (53.8, 62.0)	63.7 (62.6, 64.9)	.007	68.9 (68.5, 69.2)
Recent blood pressure test	88.6 (85.8, 91.4)	89.3 (88.5, 90.0)	.651	88.4 (88.1, 88.6)
General health poor to fair	25.3 (20.2, 30.4)	18.6 (17.4, 19.9)	.013	13.2 (12.9, 13.5)
Covered by health insurance	76.0 (72.3, 79.7)	80.9 (79.9, 81.9)	.012	85.9 (85.6, 86.2)

Note. NIRI = Nutrition Intervention Research Initiative; CI = confidence interval.

<sup>a</sup>Delta NIRI counties vs non-Delta NIRI counties.

<sup>b</sup>In 1991, there were 88 021 respondents in 48 states and the District of Columbia; in 1993, there were 102 464 respondents in 49 states and the District of Columbia.

<sup>c</sup>All racial/ethnic groups, excluding non-Hispanic Whites.

unequal probabilities of selection and the demographic distribution of the population.<sup>7</sup> In the case of prevalence estimates for Delta NIRI and non-Delta NIRI counties-parishes within the 3 states, the data were reweighted with 1990 census data to better reflect the specific demographic distributions.

## Results

In comparison with non-Delta NIRI counties-parishes in Arkansas, Louisiana, and Mississippi, Delta NIRI counties-parishes had a significantly higher proportion of residents who were African American or were members of other racial/ethnic minority groups and residents who had less than a high school education (Table 1). In Delta NIRI counties-parishes, respondents were significantly more likely to report having hypertension and to rate their health as poor to fair, and they were less likely to have health insurance and to have had cholesterol screening.

For the 3 states, the proportion of respondents reporting that a health care professional had told them that they had diabetes increased with age for men and women regardless of educational level and race (Table 2). However, among African American women and women who were members of other racial/ethnic minority groups, the prevalence peaked earlier, at 35 to 64 years, for those with less education. In these groups, 1 in 4 women reported that they had

diabetes. Among all racial/ethnic and educational groups, men and women 65 years or older reported the highest prevalence of hypertension.

The prevalence of reported obesity peaked at midlife and declined among older persons in most demographic groups. In the 35- to 64-year age group, 70% of African American women and women of other racial/ethnic minority groups who had not graduated from high school were overweight. Similarly, the highest prevalence of elevated cholesterol was among African American women and women of other racial/ethnic minority groups aged 35 to 64 years who had not graduated from high school.

## Discussion

Arkansas, Louisiana, and Mississippi respondents in Delta NIRI counties reported higher prevalences of risk factors and chronic health conditions than respondents in non-Delta NIRI counties-parishes. For the 3 states, we identified demographic subgroups with the highest prevalences of chronic health conditions for potential nutrition intervention.

Hypertension, an important risk factor for cardiovascular disease,<sup>14</sup> was significantly more prevalent in Delta NIRI counties-parishes than in non-Delta NIRI counties-parishes. Our findings are consistent with other studies that have reported an associa-

tion between lack of health insurance and poor general health.<sup>15-18</sup> Furthermore, Shea et al. similarly demonstrated significant associations between educational attainment and obesity, knowledge about blood pressure and cholesterol, and elevated cholesterol.<sup>19</sup>

Several limitations should be considered. As a result of the small subgroup sample sizes, important differences may not have been identified. Those residing in households without telephones, who tend to be of lower socioeconomic status, were not included.<sup>8,20</sup> The prevalence of health problems has been shown to be higher in such populations, so prevalence estimates reported here may be low. In the Lower Mississippi Delta, where lack of health insurance is common, respondents may not know whether they have a health disorder. Moreover, the validity of self-reported cardiovascular risk factor data has been shown to be problematic in that prevalence is underreported.<sup>21</sup> Studies have revealed that respondents tend to underreport their weight and overestimate their height, which would lower prevalence estimates of obesity.<sup>22</sup> Furthermore, because body fat tends to be higher in older persons, measurements of weight underestimate the prevalence of "over-fatness."<sup>23,24</sup> Declines in the prevalence of obesity among older persons may be due to early mortality among obese individuals.<sup>25</sup>

These results indicate the need to further assess the factors that contribute to the high prevalence of reported risk factors and dis-

**TABLE 2—Prevalence of Diabetes, Obesity, High Blood Pressure, and High Cholesterol, by Sex, Race, Age, and Educational Level: Lower Mississippi Delta States, Behavioral Risk Factor Surveillance System, 1991 and 1993**

	White		African American—Other Minority	
	Less Than High School, % (95% CI)	High School or More, % (95% CI)	Less Than High School, % (95% CI)	High School or More, % (95% CI)
<b>Diabetes</b>				
Women, age, y				
<35	2.7 (0.0, 5.4)	2.5 (1.6, 3.5)	6.0 (0.0, 13.2)	1.9 (0.4, 3.3)
35–64	6.8 (4.0, 9.5)	4.4 (3.4, 5.4)	25.5 (18.5, 32.6)	9.5 (6.3, 12.6)
65+	16.5 (12.6, 20.5)	5.8 (3.8, 7.8)	25.3 (18.3, 32.4)	7.7 (1.4, 14.1)
Men, age, y				
<35	3.4 (0.0, 6.7)	1.4 (0.0, 2.6)	1.2 (0.0, 3.0)	0.3 (0.0, 0.8)
35–64	7.2 (3.5, 10.9)	5.3 (3.9, 6.6)	5.8 (0.6, 11.0)	5.2 (1.2, 9.2)
65+	13.6 (8.2, 19.0)	11.2 (7.5, 14.9)	16.6 (7.3, 25.8)	13.5 (0.0, 27.3)
<b>High blood pressure</b>				
Women, age, y				
<35	16.1 (9.3, 22.9)	8.1 (6.3, 10.0)	21.1 (10.8, 31.4)	15.6 (11.7, 19.6)
35–64	37.5 (31.2, 43.8)	21.4 (19.3, 23.6)	64.0 (56.0, 72.0)	36.5 (31.2, 41.9)
65+	50.1 (44.7, 55.6)	40.7 (36.5, 44.9)	65.8 (58.2, 73.3)	53.0 (38.4, 67.6)
Men, age, y				
<35	11.8 (6.0, 17.7)	8.3 (6.3, 10.3)	8.9 (1.8, 16.0)	13.9 (8.9, 18.9)
35–64	27.8 (21.5, 34.0)	24.2 (21.7, 26.8)	40.6 (29.1, 52.2)	25.1 (18.6, 31.6)
65+	41.8 (33.8, 49.9)	33.0 (27.4, 38.6)	51.6 (39.6, 63.6)	44.5 (20.9, 68.0)
<b>Obesity</b>				
Women, age, y				
<35	25.6 (17.1, 34.1)	15.3 (12.9, 17.7)	53.0 (40.0, 66.1)	28.5 (23.6, 33.4)
35–64	39.2 (33.2, 45.3)	25.6 (23.2, 28.1)	69.9 (62.1, 77.6)	49.0 (43.2, 54.9)
65+	32.6 (27.5, 37.7)	22.4 (18.6, 26.1)	53.6 (46.0, 61.2)	50.1 (35.3, 65.0)
Men, age, y				
<35	21.6 (13.6, 29.5)	21.6 (18.6, 24.7)	11.7 (3.8, 19.6)	26.1 (18.8, 33.4)
35–64	36.3 (29.3, 43.4)	34.0 (31.1, 36.9)	36.3 (24.6, 47.9)	34.7 (27.6, 41.8)
65+	19.0 (12.7, 25.3)	23.9 (19.0, 28.8)	21.0 (11.2, 30.8)	19.7 (0.0, 40.4)
<b>High cholesterol</b>				
Women, age, y				
<35	16.2 (4.5, 27.9)	14.7 (11.4, 18.1)	19.3 (3.4, 35.2)	13.6 (8.3, 18.9)
35–64	41.8 (33.7, 50.0)	29.2 (26.4, 32.0)	44.9 (34.3, 55.6)	27.2 (21.2, 33.2)
65+	40.3 (34.2, 46.4)	40.8 (36.2, 45.3)	35.4 (26.6, 44.2)	20.3 (9.0, 31.6)
Men, age, y				
<35	20.2 (6.8, 33.7)	13.0 (9.2, 16.7)	11.0 (0.0, 25.4)	12.7 (5.3, 20.1)
35–64	26.3 (17.9, 34.6)	31.7 (28.4, 35.0)	30.8 (14.7, 46.8)	25.3 (17.1, 33.6)
65+	31.8 (23.2, 40.5)	27.8 (22.1, 33.6)	17.2 (6.3, 28.1)	25.0 (0.1, 49.9)

Note. The sample sizes for diabetes, high blood pressure, and obesity ranged from 223 for Black men with less than a high school education to 3480 for White women with greater than a high school education. The sample sizes for high cholesterol were smaller because not all respondents had a cholesterol reading; they ranged from 108 for Black men with less than a high school education to 2367 for White women with greater than a high school education. White includes only non-Hispanic Whites. African American—Other Minority includes all racial/ethnic groups other than non-Hispanic Whites. CI = confidence interval.

ease among population subgroups in the Lower Mississippi Delta, where poverty, low education, and lack of health insurance are common. Interventions specifically designed to improve the health of high-risk individuals in this region are needed, especially for African American women at low educational levels. □

### Contributors

J. Smith conceived of and designed the study, interpreted data findings, and wrote the initial version of the paper. S. Lensing analyzed data, and J.A. Horton interpreted data findings. S. Lensing, J.A. Horton, J. Lovejoy, S. Zaghoul, I. Forrester, B. B. McGee, and M. L. Bogle all contributed to the writing of the paper.

### Acknowledgments

This study was funded in part by the United States Department of Agriculture, Agricultural Research Service.

We wish to thank the Centers for Disease Control and Prevention and William S. Garvin III for allowing us access to the data described in this article.

### References

1. Lower Mississippi Delta Nutrition Intervention Research Consortium. In: Harrison G, ed. *Nutrition and Health Status in the Lower Mississippi Delta of Arkansas, Louisiana, Mississippi: A Review of Existing Data*. Rockville, Md: Westat Inc; 1997.
2. McCloskey AH, Woolwich J, Holahan D. *Reforming the Health Care System: State Profiles 1995*. Washington, DC: American Association of Retired Persons; 1995:20–104.
3. 1990 Census of Population and Housing [STF-3C CD-ROM]. Washington, DC: US Dept of Commerce, Bureau of the Census; 1992.
4. Kuller LH. Dietary fat and chronic diseases: epidemiologic overview. *J Am Diet Assoc*. 1997;97(suppl):S9–S15.
5. Weisburger JH. Dietary fat and risk of chronic diseases: mechanistic insights from experimental studies. *J Am Diet Assoc*. 1997;97(suppl):S16–S23.
6. Anderson RN, Kochanek KD, Murphy SL. *Report on Final Mortality Statistics, 1995*. Hyattsville, Md: National Center for Health Statistics; June 12, 1997. Monthly Vital Statistics Report, vol. 45(11 Suppl 2):1–80.
7. Remington PL, Smith MY, Williamson DF, Anda RF, Gentry EM, Hogelin GC. Design, characteristics and usefulness of state-based

- behavioral risk factor surveillance: 1981-87. *Public Health Rep.* 1988;103:366-375.
8. Frazier EL, Okoro CA, Smith C, McQueen DV. State- and sex-specific prevalence of selected characteristics—Behavioral Risk Factor Surveillance System, 1992 and 1993. *MMWR CDC Surveill Summ.* 1996;45(SS-6):1-34.
  9. Waksberg JS. Sampling methods for random digit dialing. *J Am Stat Assoc.* 1978;73:40-46.
  10. Lower Mississippi Delta Development Commission. *Final Report: The Delta Initiatives.* Memphis, Tenn: Mercury Printing; 1990.
  11. Najjar MF, Rowland M. Anthropometric reference data and prevalence of overweight. *Vital Health Stat 11.* 1987;No. 238:1-73.
  12. *Stata, Release 5.0* [computer program]. College Station, Tex: Stata Corp; 1997.
  13. Cochran WG. *Sampling Techniques.* New York, NY: John Wiley & Sons Inc; 1977:39-44.
  14. Prevalence of cardiovascular disease risk-factor clustering among persons aged 45 years—Louisiana, 1991-95. *MMWR Morb Mortal Wkly Rep.* 1997;46:585-588.
  15. Hafner-Eaton C. Physician utilization disparities between the uninsured and insured: comparisons of the chronically ill, acutely ill, and well nonelderly populations. *JAMA.* 1993;269:787-792.
  16. Patrick DL, Madden CW, Diehr P, Martin DP, Cheadle A, Skillman SM. Health status and use of services among families with and without health insurance. *Med Care.* 1992;30:941-949.
  17. Hayward RA, Benard AM, Freeman HE, Corey CR. Regular source of ambulatory care and access to health services. *Am J Public Health.* 1991;81:434-438.
  18. Health insurance coverage and receipt of preventive health services—United States, 1993. *MMWR Morb Mortal Wkly Rep.* 1995;44:219-225.
  19. Shea S, Stein AD, Basch CE, et al. Independent associations of education attainment and ethnicity with behavioral risk factors for cardiovascular disease. *Am J Epidemiol.* 1991;134:567-582.
  20. Thornberry O, Massey JT. Trends in United States telephone coverage across time and subgroups. In: Groves RM, Biemer PP, Lyberg LE, Massey WL, eds. *Telephone Survey Methodology.* New York, NY: John Wiley & Sons Inc; 1988:25-49.
  21. Bowlin SJ, Morrill BD, Nafziger AN, Jenkins PL, Lewis C, Pearson TA. Validity of cardiovascular disease risk factors assessed by telephone survey: the Behavioral Risk Factor Survey. *J Clin Epidemiol.* 1993;46:561-571.
  22. Rowland ML. Self-reported weight and height. *Am J Clin Nutr.* 1990;52:1125-1133.
  23. Forbes GB, Reina JC. Adult lean body mass declines with age: some longitudinal observations. *Metabolism.* 1970;19:653-663.
  24. Kohrt WM, Malley MT, Dalsky GP, Holloszy JO. Body composition of healthy sedentary and trained, young and older men and women. *Med Sci Sports Exerc.* 1992;24:832-837.
  25. Bray GA, Gray DS. Obesity, part 1: pathogenesis. *West J Med.* 1988;149:429-441.