



This section of the report will provide a rolling three month update on a monthly basis of the state of the climatic and ecological indicators used in monitoring areas at risk to RVF activity. These indicators include, global SST anomalies patterns, Equatorial Western Indian Ocean (WIO) and Eastern Pacific Ocean (EPO: NINO 3.4) SST anomalies, Southern Oscillation Index (SOI) and Outgoing Longwave Radiation (OLR) anomalies, Rainfall and anomalies, Normalized Difference Vegetation index anomalies and RVF risk map for Africa and the Arabian Peninsula.

## September 2014



### 1. SOI and SST Indices



### Western Indian Ocean and NINO 3.4 SST Anomalies January 1982 - September 2014





The SOI index remained consistently negative with a value of -0.9 in September as was in August. However, convective cloudiness remained generally near average over most of the region, except for below average cloudiness observed across the central and western Pacific. These conditions generally reflect the prevalence of ENSO-neutral conditions. September monthly SST anomalies in the NINO3.4 SST have remained near normal at +0.30°C in September and so are the WIO SST (0.2°C) anomalies indicating the prevalence of normal conditions over these ocean basins. Even though above-average sea surface temperatures (SST) (below) in the NINO 1& 2 regions of the far eastern equatorial Pacific continued during September 2014, they remained near average in the central equatorial Pacific and in particular most of the Niño indices remained unchanged. The absence of a clear atmospheric response to the positive SSTs indicates ENSO-neutral conditions. As a result, nearly all model forecasts have further delayed the onset to the October - December 2014 period and to peak at weak strength during the winter (3-month values of the Niño 3.4 index between 0.5°C and 0.9°C) and last into the Northern Hemisphere spring 2015.



The entire equatorial Pacific Ocean shows a pattern of above normal SSTs ( $+0.5^{\circ}\text{C}$  to  $+2.0^{\circ}\text{C}$ ) except for the region from  $30^{\circ}\text{S}$  to  $1^{\circ}\text{S}$  (off the South American coast) with below-normal SSTs during the July 2014 to September 2014. Accordingly, the entire equatorial Indian Ocean between  $30^{\circ}\text{N-S}$  is now dominated by positive SST anomalies except for the region immediately off the East African coast. Other regions of significant anomalies include the north Pacific Ocean, north Atlantic, equatorial Atlantic off the West African coast, the Pacific Ocean off the California coast, and south Indian Ocean off the southern Africa landmass which show significant positive and negative anomalies on the order of  $-/+1.0^{\circ}\text{C}$  to  $-/+2.0^{\circ}\text{C}$ . Outgoing Longwave Radiation (OLR) anomalies are used here as a proxy for tropical deep convection (rainfall). Reduced convection is shown in yellow to light brown and brown shades and increased/intense convection is shown by shades of blue. Some impacts from the current SST anomaly patterns can be observed in the pattern of global convective activity illustrated by the OLR departure patterns here. During the July to September 2014 period, drier-than-average conditions ( $>+35\text{W/M}^2$ ) are observed over the equatorial south-western Pacific Ocean south of the equator between  $120^{\circ}\text{W}$  and  $150^{\circ}\text{E}$ , over the Caribbean region and central Indian Ocean and


enhanced cooler than average conditions ( $-40\text{W}/\text{M}^2$ ) over the equatorial central to eastern Pacific Ocean north of the equator between  $155^\circ\text{W}$  and  $900^\circ\text{E}$ . Convective activity continues to be prevalent over parts equatorial Africa particularly Chad and Sudan, central Canada and most of western half of the US, Mexico ( $-55\text{W}/\text{M}^2$ ) and Southern Africa. These patterns of depressed and enhanced convective activity coincide well with the patterns of SST departures. Monthly and weekly anomalies can be found [here](#). Rainfall and associated anomalies (below) for Africa from July 2014 to September 2014 show rainfall concentrated from the equator to  $15^\circ\text{N}$  with a maxima along  $5^\circ\text{N}$ . Areas of above normal rainfall ( $+50$  to  $250\text{mm}$ ) include parts of equatorial West Africa to central to eastern Sudan into northern Ethiopia.






Cumulative NDVI anomalies for Africa for July 2014 to September 2014 show positive anomalies concentrated over parts of central Sudan, north-western Ethiopia, Botswana and Namibia following the above normal rainfall in these areas in the last several months. The RVF risk map below was derived from thresholding NDVI anomaly data to detect areas persistent of above normal NDVI. Periods of widespread and prolonged heavy rainfall lead to flooding of dambos and anomalous green up in vegetation, creating ideal ecological conditions for the emergence RVF vectors. For the period July 2014 to September 2014, the RVF persistence model identifies isolated areas in Sudan and South Sudan where ecological conditions would support the emergence of RVF vectors. Enhanced surveillance is advised in these areas. These locations have reported cases Cholera over the last few months due to flooding and poor sanitary conditions.

# NDVI Anomaly July - September 2014



# RVF Potential September 2014



AFHSC/GEIS  
USDA/ARS-CMAVE

- RVF risk areas, humans and livestock present
- RVF risk areas, humans and livestock absent
- RVF potential epizootic areas

NASA/GSFC-GIMMS