

MIC Distribution and Percent Resistance among *Campylobacter coli* and *Campylobacter jejuni* isolates from poultry for 2009

Antimicrobial	Species	n	% Resistant	Distribution (%) of MICs (µg/ml) <sup>a</sup>														
				0.016	0.032	0.064	0.125	0.25	0.5	1	2	4	8	16	32	64	>64	
Azithromycin	<i>C. coli</i>	81	6.2		9.9	65.4	18.5										6.2	
	<i>C. jejuni</i>	117	0	3.4	57.3	36.8	2.6											
Ciprofloxacin	<i>C. coli</i>	81	22.2			18.5	50.6	8.6				8.6	13.6					
	<i>C. jejuni</i>	117	19.7		4.3	47.9	26.5	1.7			1.7	12	5.1	0.9				
Clindamycin	<i>C. coli</i>	81	0			1.2	48.1	37	3.7	2.5	1.2	6.2						
	<i>C. jejuni</i>	117	0		2.6	42.7	41	12	1.7									
Erythromycin	<i>C. coli</i>	81	6.2				1.2	16	37	33.3	6.2						6.2	
	<i>C. jejuni</i>	117	0				8.5	41	41.9	6.8	1.7							
Florfenicol	<i>C. coli</i>	81	0						7.4	86.4	6.2							
	<i>C. jejuni</i>	117	0					0.9	63.2	35	0.9							
Gentamicin	<i>C. coli</i>	81	2.5					32.1	65.4								2.5	
	<i>C. jejuni</i>	117	0.9				17.9	50.4	30.8								0.9	
Nalidixic Acid	<i>C. coli</i>	81	22.2									72.8	4.9				17.3	4.9
	<i>C. jejuni</i>	117	19.7									76.9	3.4				8.5	11.1
Telithromycin	<i>C. coli</i>	81	6.2				4.9	12.3	8.6	40.7	25.9	1.2		6.2				
	<i>C. jejuni</i>	117	0				1.7	20.5	51.3	22.2	4.3							
Tetracycline	<i>C. coli</i>	81	44.4				3.7	32.1	9.9	8.6		1.2			4.9	13.6	25.9	
	<i>C. jejuni</i>	117	49.6			3.4	23.1	13.7	6	2.6		0.9	0.9	1.7	8.5	22.2	17.1	

<sup>a</sup> The unshaded areas indicate the range of dilutions tested for each antimicrobial. Black vertical lines indicate susceptibility breakpoints while red vertical lines indicate resistance breakpoints. Values shown above the range denote the percentage of isolates with MIC values greater than the highest tested concentration. Isolates with MICs equal to or less than the lowest tested concentration are given as the lowest concentration.