

MIC Distribution and Percent Resistance among *Campylobacter coli* and *Campylobacter jejuni* isolates from poultry for 2006

Antimicrobial	Species	n	% Resistant	Distribution (%) of MICs ($\mu\text{g/ml}$) ^a														
				0.016	0.032	0.064	0.125	0.25	0.5	1	2	4	8	16	32	64	>64	
Azithromycin	<i>C. coli</i>	123	8.9	0.8	23.6	50.4	13.8	1.6		0.8								8.9
	<i>C. jejuni</i>	228	0.4	22.4	52.2	21.5	2.6			0.4	0.4							0.4
Ciprofloxacin	<i>C. coli</i>	123	15.4		2.4	29.3	43.9	8.9					4.9	10.6				
	<i>C. jejuni</i>	228	8.8		5.3	54.8	28.5	1.8		0.4	0.4		7.0	1.8				
Clindamycin	<i>C. coli</i>	123	1.6			6.5	39.8	36.6	4.1	0.8	4.9	5.7	1.6					
	<i>C. jejuni</i>	228	0		1.8	38.2	44.3	13.2	2.2			0.4						
Erythromycin	<i>C. coli</i>	123	8.9				6.5	30.9	10.6	30.9	8.9		3.3					8.9
	<i>C. jejuni</i>	228	0.4				15.4	47.4	28.1	6.6	1.8	0.4						0.4
Florfenicol	<i>C. coli</i>	123	0						5.7	90.2	4.1							
	<i>C. jejuni</i>	228	0					1.8	45.2	48.2	4.8							
Gentamicin	<i>C. coli</i>	123	0					30.9	66.7	2.4								
	<i>C. jejuni</i>	228	0				7.0	54.4	38.2	0.4								
Nalidixic Acid	<i>C. coli</i>	123	15.4									69.1	15.4				11.4	4.1
	<i>C. jejuni</i>	228	8.8									82.0	8.3	0.9			4.8	3.9
Telithromycin	<i>C. coli</i>	123	6.5				2.4	30.9	5.7	13.0	32.5	3.3	5.7	6.5				
	<i>C. jejuni</i>	228	0				2.6	21.9	51.8	18.9	3.9	0.4	0.4					
Tetracycline	<i>C. coli</i>	123	53.7				13.0	28.5	3.3	0.8		0.8		0.8	3.3	7.3		42.3
	<i>C. jejuni</i>	228	56.1			4.8	23.7	7.9	3.9	1.3	0.4		1.8	3.5	10.1	25.0		17.5

^a The unshaded areas indicate the range of dilutions tested for each antimicrobial. Single vertical lines indicate susceptibility breakpoints while double vertical lines indicate resistance breakpoints. Values shown above the range denote the percentage of isolates with MIC values greater than the highest tested concentration. Isolates with MICs equal to or less than the lowest tested concentration are given as the lowest concentration.