



NARMS-EB 2000

MICs for *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Chloramphenicol among *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Ciprofloxacin among *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Clindamycin among *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Erythromycin among *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Gentamicin among *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Nalidixic Acid among *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Tetracycline among *Campylobacter jejuni* isolates




NARMS-EB 2000

MICs for Azithromycin among *Campylobacter coli* isolates




NARMS-EB 2000

MICs for Chloramphenicol among *Campylobacter coli* isolates




NARMS-EB 2000

MICs for Ciprofloxacin among *Campylobacter coli* isolates




NARMS-EB 2000

MICs for Clindamycin among *Campylobacter coli* isolates




NARMS-EB 2000

MICs for Erythromycin among *Campylobacter coli* isolates




NARMS-EB 2000

MICs for Gentamicin among *Campylobacter coli* isolates



NARMS-EB 2000

MICs for Nalidixic Acid among *Campylobacter coli* isolates



NARMS-EB 2000

MICs for Tetracycline among *Campylobacter coli* isolates

