


NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**

Amoxicillin/Clavulanic Acid




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**




*including copenhagen

NARMS – EB 2002

Veterinary Isolates

**Fig. 22 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Cattle (Diagnostic)**

Trimethoprim/Sulfamethoxazole



*including copenhagen