

**TABLE 3: Total percent veterinary *Salmonella* isolates sensitive, intermediate, or resistant**

Antimicrobial	Sensitive		Intermediate		Resistant	
	n	%	n	%	N	%
Amikacin*	6950	100.0	0	0.0	0	0.0
Amoxicillin/ Clavulanic Acid	5528	79.2	368	5.3	1081	15.5
Ampicillin	4986	71.5	2	0.0	1989	28.5
Cefoxitin	5884	84.3	144	2.1	949	13.6
Ceftiofur	5935	85.1	4	0.1	1038	14.9
Ceftriaxone	6299	90.3	657	9.4	21	0.3
Cephalothin	5749	82.4	101	1.4	1127	16.2
Chloramphenicol	5646	80.9	63	0.9	1268	18.2
Ciprofloxacin	6975	100.0	0	0	2	0.0
Gentamicin	6337	90.8	92	1.3	548	7.9
Kanamycin	5950	85.3	34	0.5	993	14.2
Nalidixic Acid	6887	98.7	0	0	90	1.3
Streptomycin	4461	64.0	0	0	2514	36.0
Sulfamethoxazole	4994	71.6	0	0	1983	28.4
Tetracycline	4143	59.4	21	0.3	2813	40.3
Trimethoprim/ Sulfamethoxazole	6711	96.2	0	0	266	3.8

\*Due to concentrations tested for Amikacin (0.5 – 4.0 ug/ml), only susceptible isolates were able to be determined. Twenty-seven isolates presented an MIC of  $\geq 4.0$  ug/ml