


NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004 Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)




↓ Breakpoint

* Including var 5- formerly var Copenhagen

NARMS – EB 2004

Veterinary Isolates

Fig. 25 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Chicken (Diagnostic)



↓ Breakpoint

* Including var 5- formerly var Copenhagen