





NARMS – EB 2004 Veterinary Isolates

Fig. 4 Minimum Inhibitory Concentrations (µg/ml) by Antimicrobial Agent for All *Salmonella* Isolates from Chicken



↓ Breakpoint


Diagnostic n=76

Slaughter n=1280


NARMS – EB 2004 Veterinary Isolates

Fig. 4 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for All *Salmonella* Isolates from Chicken


Ceftiofur




Ceftriaxone



Chloramphenicol



Ciprofloxacin



↓ Breakpoint


Diagnostic n=76

Slaughter n=1280


NARMS – EB 2004 Veterinary Isolates

Fig. 4 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for All *Salmonella* Isolates from Chicken


Gentamicin




Kanamycin



Nalidixic Acid



Streptomycin



↓ Breakpoint

Diagnostic n=76

Slaughter n=1280

NARMS – EB 2004 Veterinary Isolates

Fig. 4 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for All *Salmonella* Isolates from Chicken

