





NARMS – EB 2004 Veterinary Isolates

Fig. 6 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for All *Salmonella* Isolates from Turkey




↓ Breakpoint

Diagnostic n=130 Slaughter n=236


NARMS – EB 2004 Veterinary Isolates

Fig. 6 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for All *Salmonella* Isolates from Turkey


Ceftiofur




Ceftriaxone



Chloramphenicol



Ciprofloxacin




↓ Breakpoint

Diagnostic n=130 Slaughter n=236


NARMS – EB 2004 Veterinary Isolates

Fig. 6 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for All *Salmonella* Isolates from Turkey


Gentamicin




Kanamycin



Nalidixic Acid



Streptomycin



↓ Breakpoint

Diagnostic n=130

Slaughter n=236

NARMS – EB 2004 Veterinary Isolates

Fig. 6 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for All *Salmonella* Isolates from Turkey

