

**Table 20. Confirmed *S. Typhimurium* DT104<sup>1,2</sup> Isolates from Food Animals, 1997-2005**

Year	Chicken			Turkey			Cattle			Swine		
	n	% ( <i>S.Typhimurium</i> )	% (Chicken)	n	% ( <i>S.Typhimurium</i> )	% (Turkey)	n	% ( <i>S.Typhimurium</i> )	% (Cattle)	n	% ( <i>S.Typhimurium</i> )	% (Swine)
1997	4	16.7	1.9	0	0.0	0.0	1	50.0	4.2	11	44.0	9.9
1998	11	16.7	2.0	0	0.0	0.0	2	6.1	0.7	48	45.7	6.1
1999	12	7.8	0.8	2	5.4	0.3	37	19.6	2.3	34	29.8	3.9
2000	18	12.4	1.5	3	16.7	0.6	46	24.6	3.3	25	30.9	5.5
2001	14	10.8	1.1	2	13.3	0.4	20	23.0	2.2	15	34.1	3.6
2002	16	10.7	1.1	1	11.1	0.4	21	21.4	2.1	13	27.1	3.4
2003	4	2.6	0.3	1	16.7	0.4	10	12.8	1.5	8	29.6	3.8
2004	3	1.8	0.2	0	0.0	0.0	14	29.2	2.3	11	20.8	3.6
2005	9	4.9	0.5	2	28.6	0.9	7	20.6	2.1	12	28.6	4.0

<sup>1</sup> Includes isolates that are DT104 complex (A or B)

<sup>2</sup> Includes *S. Typhimurium* and *S. Typhimurium* var. 5-