


**NARMS-EB 2003
Veterinary Isolates**

**Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)**

Amikacin




* Includes var copenhagen

NARMS-EB 2003
 Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
 Major Serotypes from Chicken (Slaughter)

Amoxicillin/Clavulanic Acid




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Ampicillin




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Cefoxitin



Breakpoint=32


Minimum Inhibitory Concentration

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Ceftiofur




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
 Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
 Major Serotypes from Chicken (Slaughter)

Ceftriaxone




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Cephalothin




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Chloramphenicol




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Ciprofloxacin




↓ Breakpoint

* Includes var copenhagen

**NARMS-EB 2003
Veterinary Isolates**

**Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)**

Gentamicin




↓ Breakpoint

* Includes var copenhagen

**NARMS-EB 2003
Veterinary Isolates**

**Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)**

Kanamycin




↓ Breakpoint

* Includes var copenhagen

**NARMS-EB 2003
Veterinary Isolates**

**Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)**

Nalidixic Acid




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Streptomycin




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
 Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
 Major Serotypes from Chicken (Slaughter)

Sulfamethoxazole




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Tetracycline




↓ Breakpoint

* Includes var copenhagen

NARMS-EB 2003
Veterinary Isolates

Fig. 26 Minimum Inhibitory Concentrations by Antimicrobial Agent
Major Serotypes from Chicken (Slaughter)

Trimethoprim/Sulfamethoxazole



↓ Breakpoint

* Includes var copenhagen