

Pectinolytic and Pectolytic Microorganisms

R. F. McFeeters, L. Hankin, and G. H. Lacy

18.1 INTRODUCTION

Pectic substances are important cell wall components of higher plants, particularly dicots. The specific functions that pectic materials perform in the cell wall are not understood. However, it appears that they are important in cementing plant cells together. BeMiller⁶ has reviewed pectin structure. Pectic substances are polymers of D-galacturonic acid residues glycosidically linked alpha-1,4. Pectin molecules contain occasional rhamnose units with 1,2 linkages in the main chain. They also have side chains on both the galacturonic acid and rhamnose residues that contain mainly galactose and arabinose residues. The carboxyl groups of galacturonic acid residues are usually methylesterified to a substantial degree. This has major effects on physical properties of pectin, such as gelation. The degree of methylation in plants has been found to range from about 40% to 90% of the carboxyl groups.² Data on the distribution of carboxyl groups are limited, but it appears most often to be random in pectin isolated to minimize enzymatic or chemical modification.² In some plants, substantial numbers of the hydroxyl groups of galacturonic acid residues are acetylated. This modification inhibits pectin gelation.⁶

The nomenclature for pectic substances has been somewhat variable and confused over the years. For the purpose of this chapter, pectic substances is an inclusive term for galacturonic acid-containing polymers from plant cell walls. Pectin is used for pectic substances with a substantial fraction of the galacturonic acid carboxyl groups esterified. Pectic acid refers to polymers with a negligible amount of the carboxyl groups esterified. Polypectate or pectate refers to pectic acid with carboxyl groups in the salt form. Pectinolytic refers to the degradation of pectin and pectolytic to the degradation of pectic acid or pectate. Commercially available pectin of the type used for microbiological or enzymatic assays generally has >60% methylation. Pectic acid or polypectate is <5% esterified.

18.11 Sources of Pectinolytic and Pectolytic Enzymes

Most pectin-degrading organisms are associated with raw agricultural products and with soil. Up to 10% of the organisms in soil have been shown to be pectinolytic.²³ These include, but are not limited to, bacteria in the genera *Achromobacter*, *Aeromonas*,

Arthrobacter, *Agrobacterium*, *Enterobacter*, *Bacillus*, *Clostridium*, *Erwinia*, *Flavobacterium*, *Pseudomonas*, and *Xanthomonas*,^{54,59} as well as in yeasts, molds, protozoa, and nematodes.³ Many of these organisms are plant pathogens.^{3,17} A survey of yeasts from 13 genera⁸ found pectinolytic activity produced by 69 of 207 strains tested. Pectolytic activity is not common in lactic acid bacteria. However, it has now been reported to occur in at least one strain of *Leuconostoc mesenteroides*,³⁰ *Lactobacillus acidophilus*,³⁴ *Lactobacillus casei*, *Lactobacillus plantarum*, and *Lactococcus lactis*.³¹ The molecular biology of pectolytic enzymes from plant pathogenic erwiniae has been extensively investigated.^{12,36,51} The discussion that follows refers to aerobic procedures. However, the detection of anaerobic pectinolytic bacteria also has been described.^{41,46}

18.2 DETECTING PECTINOLYTIC AND PECTOLYTIC ORGANISMS

The basic method used to detect pectinolytic or pectolytic organisms has been to grow the organisms on a gel medium that contains pectin or pectate substrates, respectively. Production of enzymes by a culture is detected either by observing depressions in the gel around the colony where the substrate has been degraded or by flooding the plate with a precipitant solution. Around producer colonies a clear zone will appear where the substrate has degraded to the point that precipitation does not occur, while non-producing colonies will be surrounded by opaque gel containing the non-degraded pectin or pectate substrate. Wieringa⁶⁰ reported the first medium of this type. Over the years, many variations of this theme have been developed to address particular problems of sample handling, enzyme specificity, sensitivity, or isolation of organisms from the plates. For example, a researcher can cut holes in agar gel plates with a cork borer in order to assay liquid samples, such as culture filtrates, for enzyme activity. This is the so-called well plate or cup plate technique for enzyme assays.

18.21 Pectate and Pectin Lyase Producers

Considerable research has been conducted on techniques to detect pectate lyase-producing organisms. This is because these enzymes from *Erwinia* species have been cloned into *Escherichia coli*. Since lyases have alkaline pH optima, while poly-galacturonases have acidic optima, a medium pH of 7.0 or above is the main

parameter used to distinguish pectate or pectin lyase producers from polygalacturonase producers. Durrands and Cooper¹⁸ provide an example of this approach, in which media were designed to detect polygalacturonase and pectin lyase production by *Verticillium albo-atrum* mutants. Roberts et al.^{52,53} and Allen et al.¹ used the pH 8.5 PEC-YA medium of Starr et al.⁵⁷ to clone pectate lyase genes from *Erwinia carotovora*.

18.211 Bacterial Pectate Lyases

Several media have been developed for the detection of bacteria that produce pectate lyases. King and Vaughn³⁵ developed a pectate medium with a pH of 7.0 that contained crystal violet to make it selective for gram-negative bacteria. Adding cycloheximide further inhibits the growth of yeasts and molds. Detection of pectolytic colonies is based on formation of depressions in the pectate gel because of enzymatic degradation.

Hankin et al.²⁴ used a mineral medium with 0.1% yeast extract, pectin, and agar to detect pectolytic colonies of *Erwinia* and *Pseudomonas*. A 1% aqueous solution of hexadecyltrimethylammonium bromide²⁹ was used to precipitate non-degraded substrate so that pectolytic colonies showed a clear zone on a white background. The researchers emphasized that a high phosphate level in the medium was needed to observe pectolytic activity. Sands et al.⁵⁵ modified the medium of King and Vaughn³⁵ by using 2% pectin and 1.5% agar instead of 7% polypectate. They then added a mixture of novobiocin, penicillin G, and cycloheximide to make the medium selective for fluorescent pseudomonads. Hexadecyltrimethylammonium bromide solution was used to precipitate the pectin for visualization of clear zones around pectolytic colonies.

Cuppels and Kelman¹⁵ did a detailed evaluation of the selectivity for and recovery of *Erwinia* from natural samples using another pectate medium containing crystal violet to prevent the growth of unwanted organisms. The medium gave an excellent recovery of pectolytic *Erwinia*, but did allow growth and enzyme production by some pseudomonads. This medium has also been used to isolate pectolytic strains of *Cytophaga johnsonae* from spoiled, fresh bell peppers and watermelon.³⁹ Woodward and Robinson⁶² modified crystal violet medium by addition of proteose peptone to reduce the time required to detect pectolytic *Erwinia* and improve pit formation. Pierce and McCain⁴⁸ modified the medium of Miller and Schroth⁴³ to selectively plate for pectolytic *Erwinia*. They found it to be selective for *Erwinia* and to improve recovery compared to the Cuppels and Kelman crystal violet medium. With all of these media, it has been reported that not all commercial polypectate preparations give suitable gelation. A procedure has been described to produce polypectate from orange peel and apple pulp that will give a good gel.¹³

An essential element of work to identify and characterize pectin-degrading genes from *Erwinia chrysanthemi* was the development of plating techniques to make it possible to identify clones that contained the genes of interest. Keen et al.³³ described the isolation of *E. coli* clones that contain pectate lyase genes. They used a pectate agar at pH 8.0. After incubating samples, they detected lyase activity by flooding the plates with 1 M CaCl₂. A white halo formed around positive clones. Kotoujansky et al.³⁷ developed a technique to isolate clones of lambda-L47-1 phage to which pectate lyase genes had been transferred. They used a medium for *E. coli*, the phage host, in one layer, and a pectate medium in the bottom layer. The two gels were separated by a nylon membrane that allowed enzymes to diffuse into the pectate

layer, but prevented transfer of the phage. Zones with pectate lyase activity were visualized by removing the nylon membrane and the upper gel and flooding the pectate medium with 1 M CaCl₂. The nylon membrane was then placed back on the plate so that phage clones could be isolated from the appropriate plaque.

18.212 Bacterial Pectin Lyases

Pectin lyases should give clearing zones on plates with pectin as the substrate. However, plates are not very sensitive for this group of pectic enzymes, and the zones produced can be indistinct.⁵⁶ Detection is accomplished by spectrophotometric assays at 235 nm on culture filtrate samples with high methoxyl pectin as substrate. Pectin lyases will give little or no measurable activity with polypectate as the substrate.

18.213 Fungal Pectate Lyases

Hankin and Anagnostakis²² describe a plate technique with a medium that contains 1% pectin, 0.2% yeast extract, mineral salts, and 3% agar adjusted to pH 7.0. After a 3- to 5-day incubation period, plates are flooded with 1% aqueous hexadecyltrimethylammonium bromide to precipitate non-degraded pectin. Clear zones occur around colonies that produce pectate lyase. If the precipitant is not allowed to remain in contact with fungal cells for more than 5 min, viable colonies can be isolated from the flooded plates. For fungal samples from natural isolations that contain bacteria, a mixture of the antibiotics neomycin and chloramphenicol provides control of bacterial growth with the least inhibition of growth or enzyme production by fungi. However, the authors emphasize that fungi should be purified and enzyme production checked in the absence of the antibiotics.

18.22 Fungal Polygalacturonase Producers

The detection of polygalacturonases by plate assays has generally been done simply by lowering the pH of a medium designed for detection of pectate lyase to 6 or below, so that polygalacturonases will be active and pectate lyases will be inactive.^{22,58}

18.23 Detecting Pectic Enzymes during Germination of Fungal Spores

Hagerman et al.²¹ described a plate procedure for detection of pectolytic enzymes, protease, and cellulase activity during germination of *Botrytis cinerea* spores. The method is very sensitive for the detection of lyases and pectinesterase, but it is considerably less sensitive for polygalacturonase.

18.24 Evaluating Macerating Activity of Pectic Enzymes

Mussell and Morre⁴⁴ analyzed in detail factors affecting the maceration of cucumber tissue by commercial polygalacturonase from *Aspergillus niger*, basing their procedure on the measurement of weight loss of the cucumber tissue after enzyme treatment. They pointed out that tissue maceration assay for polygalacturonase activity was about 500 times more sensitive than viscosity assays. Ishii²⁸ developed a procedure to evaluate the maceration of potato, onion, and radish tissues by measuring the volume of separated cells released from tissue samples. A polygalacturonase and pectate lyase, separately and in combination with *Aspergillus japonicus*, were used. Both enzymes caused tissue maceration, but the relative activity of the enzymes varied with the plant tissue.

18.3 VISUALIZATION AND ASSAY OF PECTIC ENZYMES

18.31 *Detecting Pectic Enzymes in Electrophoresis Gels*

The visualization of pectic enzymes in gels following electrophoresis has been described. Cruickshank and Wade¹⁴ incorporated citrus pectin into acrylamide slab gels. By using suitable incubation and staining procedures, they were able to differentiate pectate lyases, polygalacturonases, and pectinesterases in the gels. Bertheau et al.⁷ developed a sandwich technique in which suitable buffers and substrates could diffuse into acrylamide slab gels so that a variety of enzymes, including pectate lyases and polygalacturonases, could be detected in the gels.

18.32 *Detecting Pectic Enzymes on Isoelectric Focusing Gels*

A sensitive and rapid method for visualizing the isoelectric profiles of pectic enzymes using activity overlays has been devised.⁵⁰ Isoelectrically focused proteins in ultra-thin (0.35 mm) 5% acrylamide gels containing broad-range (pH 3.5 to 10) and high-pH (pH 9 to 11) ampholytes bonded to plastic support (Gel Bond PAG acrylamide support, FMC BioProducts, Rockland, MD 04841-2994) are overlaid with an ultra-thin (0.35 mm) layer of 1% agarose containing 0.1% polygalacturonic acid bonded on Gel Bond agarose support and also supported on plastic. After sufficient reaction time, the overlay film is stained with ruthenium red (0.05% wt/vol). The substrate gel stains a deep pink because of reaction of the ruthenium red with polygalacturonate, while the substrate in areas over the isoelectrically focused pectic enzymes has been degraded so that a clear band shows on the pink background.

18.33 *Pectinesterase*

Pectinesterase can be assayed based on the release of either free carboxyl groups or methanol from pectin. The most common technique is to measure the rate of release of free carboxyl groups from pectin using a pH stat.³⁶ A rapid, continuous spectrophotometric assay has been developed.³⁰ It is based on measuring the change in absorbance of a pH indicator as the pH decreases because of the formation of free carboxyl groups in pectin. Methanol may be measured colorimetrically⁶¹ or by gas chromatography.⁴² For colorimetric analysis, the modifications described by Hudson and Buescher²⁶ should be used because the incubation procedure of Wood and Siddiqui¹ results in nonlinear color development.

18.34 *Pectate and Pectin Lyase*

Pectate lyase activity can be assayed spectrophotometrically at 235 nm because of the formation of the 4,5-double bond in the non-reducing galacturonic acid residues.⁵⁷ For pectin lyase activity, pectin may be substituted for polypectate as the substrate. A pectin lyase should show little or no activity with polypectate as the substrate.

18.35 *Polygalacturonase*

The release of reducing groups because of hydrolysis of polypectate is the most common method for measurement of polygalacturonase activity. Lee and MacMillan³⁸ used 0.5% polypectate in 0.1 M sodium acetate buffer, pH 4.5 at a temperature of 30°C for tomato polygalacturonase. These are reasonable conditions for most microbial polygalacturonases, though 0.1% polypectate is

usually a saturating substrate level. For measurement of reducing groups, the procedure of Nelson⁴⁵ has been used for many years. However, this method requires the addition of one reagent solution before heating and a second after heating for color development. Also, the samples must be centrifuged prior to measurement of the absorbance because of cloudiness caused by precipitated substrate. A reducing sugar method using 2,2'-bichinonate¹⁶ is highly sensitive and avoids the inconveniences of the Nelson procedure. Gross¹⁹ demonstrated the use of 2-cyanoacetamide reagent, which also allows detection of reducing groups using a single reagent solution. Absorbance is measured at the UV region, 276 nm. Crude extracts with high levels of ultraviolet-absorbing material may cause interference with this reagent.

Sensitive, semi-quantitative assays to assess softening activity in pickle fermentation brines and spent brines that are recycled have been developed specifically for this industry. Bell et al.⁴ measured loss in viscosity of pH 5 polypectate solutions after a 20-hr or 44-hr incubation period. Buescher and Burgin⁹ have described a diffusion plate assay that is in common use to determine if there is polygalacturonase activity in brines before they are recycled.

Pectic enzymes are inhibited by phenolic compounds, indoleacetic acid, fatty acid, and endopolygalacturonase end-products.³ Tannins from certain plants inhibit both pectic enzymes and cellulase.⁵ Specific protein inhibitors of polygalacturonases have also been reported in various plants. Cervone et al.¹¹ purified an inhibitor protein from French bean hypocotyls and demonstrated that it inhibited polygalacturonases from *Colletotrichum lindemuthianum*, *Fusarium moniliforme*, and *Aspergillus niger*. Diethylpyrrolidone, which is used to chemically modify histidyl residues in proteins, has been shown to inhibit an endo-polygalacturonase from *Aspergillus ustus*.⁴

18.4 SOURCES OF PECTIC ENZYMES

Purified endo-splitting polygalacturonases and pectin lyases from *Aspergillus niger* and *Aspergillus japonicus* are available commercially. However, the preparations each contain low levels of the other enzyme as a contaminant. The ability to clone pectate lyase genes from *E. chrysanthemi* into *E. coli* makes available clones with sequenced genes that are good producers of individual *E. chrysanthemi* pectate lyases.³² In addition, *E. coli* clones containing *E. carotovora* genes for endo- and exo-pectate lyases are available.^{52,53} Since *E. coli* does not produce pectic enzymes, these clones will not produce pectinesterase or polygalacturonase. Phaff⁴⁷ described a yeast, *Kluyveromyces fragilis*, that reliably secretes large amounts of polygalacturonase into the growth medium. The polygalacturonase was estimated to be about 95% pure in the culture filtrate. The organism does not produce either pectinesterase or pectate lyase. Three or more polygalacturonases are present in the preparation.^{10,27,41} All of the enzymes appear to be endo-splitting with similar tissue-macerating properties.⁴⁰ Despite the presence of multiple enzymes, this is a source for easily produced polygalacturonases that are free of other classes of pectic enzymes. The culture can be obtained from Dr. Phaff for a nominal handling fee.

18.5 EQUIPMENT, MATERIALS, AND REAGENTS

Crystal violet pectate (CVP) medium
Medium to detect pectate lyase (MP-7 medium)
Medium to detect polygalacturonase (MP-5 medium)

Polypectate gel medium
 Polysaccharide precipitant
 Selective medium for fluorescent pectinolytic pseudomonads
 (FPA medium)

18.6 PROCEDURES

18.61 Sample Preparation

Prepare a homogenous suspension and appropriate dilutions of the food (Chapter 2). In food containing a large amount of glucose, sufficient dilution may be necessary to avoid possible catabolite repression of pectic enzyme synthesis.²⁵ Also, in high-sugar foods such as jams and jellies, exercise care in the dilution of the sample to prevent osmotic shock to the cells by the use of an appropriate buffer or isotonic solution.

18.62 Preparation and Incubation of Plates

Place not more than 0.25 mL quantities of appropriate dilutions (to obtain 20 to 30 colonies and to avoid a wet agar surface) on the surface of pre-poured plates. Distribute by a spread plate technique. Incubation of plates inoculated with a variety of plant materials has been carried out at 30°C for 48 hr. Incubation of plates inoculated with foods probably should be made at the temperature at which the food is stored. Some pectinolytic bacteria grow at 37°C, but do not produce pectate lyase until the temperature is 32°C or below. *E. carotovora* subsp. *atroseptica* often will not grow above 28°C.

18.63 Counting and Reporting

Colonies that cause depressions in polypectate gel medium or that are surrounded by a clear zone after flooding plates with a pectin precipitant are pectinolytic. Fluorescent bacteria are detected on FPA plates with long-wavelength UV light before adding pectin precipitant.

18.7 INTERPRETATION

The varieties of organisms that can degrade pectic substances are diverse. Thus, choice of the most suitable medium depends on the type of organism expected, as well as the type of food to be examined. Media and procedures are described for the detection, enumeration, and isolation of pectolytic and pectinolytic microorganisms. In addition, procedures are described for the detection of pectic enzymes. These techniques can be used to characterize degradation of fruits, vegetables, and processed foods, to screen for plant pathogens or soil organisms that degrade pectic substances, and to indicate the stability of raw plant products for storage and transport.

18.8 REFERENCES

- Allen, C., H. George, Z. Yang, G. H. Lacy, and M. S. Mount. 1987. Molecular cloning of an *endo*-pectate lyase gene from *Erwinia carotovora* subsp. *atroseptica*. *Physiol. Molec. Plant Pathol.* 31:325.
- Anger, H., and G. Dongowski. 1985. Distribution of free carboxyl groups in native pectins from fruits and vegetables. *Die Nahrung* 29:397.
- Bateman, D. F., and R. L. Millar. 1966. Pectic enzymes in tissue degradation. *Ann. Rev. Phytopath.* 4:119.
- Bell, T. A., J. L. Etchells, and I. D. Jones. 1955. A method for testing cucumber salt-stock brine for softening activity. *U.S. Dep. Agric. Agric. Res. Serv.* 72-5.
- Bell, T. A., J. L. Etchells, and W. W. G. Smart Jr. 1965. Pectinase and cellulase enzyme inhibitor from sericea and certain other plants. *Bot. Gaz. Chicago* 126:40.
- BeMiller, J. N. 1986. An introduction to pectins: Structure and properties, p. 2. *In* M. L. Fishman, J. J. Jen (ed.) *Chemistry and function of pectins*. Am. Chem. Soc., Washington, D. C.
- Bertheau, Y., E. Madgidi-Hervan, A. Kotoujansky, C. Nguyen-The, T. Andro, and A. Coleno. 1984. Detection of depolymerase isoenzymes after electrophoresis or electrofocusing, or in titration curves. *Anal. Biochem.* 139:383.
- Biely, P., and E. Slavikova. 1994. New search for pectolytic yeasts. *Folia Microbiol.* 39:485-488.
- Buescher, R. W., and C. Burgin. 1992. Diffusion plate assay for measurement of polygalacturonase activities in pickle brines. *J. Food Biochem.* 16:59-68.
- Call, H. P., and C. C. Emeis. 1983. Characterization of an "endopolygalacturonase" of the yeast *Kluyveromyces marxianus*. *J. Food Biochem.* 7:59.
- Cervone, F., G. De Lorenzo, L. Degra, G. Salvi, and M. Bergami, M. 1987. Purification and characterization of a polygalacturonase-inhibiting protein from *Phaseolus vulgaris* L. *Plant Physiol.* 85:631.
- Collmer, A., and N. T. Keen. 1986. The role of pectic enzymes in plant pathogenesis. *Ann. Rev. Phytopath.* 24:383.
- Cother, E. J., A. B. Blakeney, and S. J. Lamb. 1980. Laboratory-scale preparation of sodium polypectate for use in selective media for pectolytic *Erwinia* spp. *Plant Dis.* 64:1086.
- Cruikshank, R. H., and G. C. Wade. 1980. Detection of pectic enzymes in pectin-acrylamide gels. *Anal. Biochem.* 107:177.
- Cuppels, D., and A. Kelman. 1974. Evaluation of selective media for isolation of soft-rot bacteria from soil and plant tissue. *Phytopathology* 64:468.
- Doner, L. W., and P. L. Irwin. 1992. Assay of reducing end-groups in oligosaccharide homologues with 2,2'-bicinchoninate. *Anal. Biochem.* 202:50-53.
- Dowson, W. J. 1957. *Plant diseases due to bacteria*, 2nd ed. Cambridge University Press, New York.
- Durrands, P. K., and R. M. Cooper. 1988. Development and analysis of pectic screening media for use in the detection of pectinase mutants. *Appl. Microbiol. Biotechnol.* 28:463.
- Gross, K. C. 1982. A rapid and sensitive spectrophotometric method for assaying polygalacturonase using 2-cyanoacetamide. *HortScience* 17:933.
- Hagerman, A. E., and P. J. Austin. 1986. Continuous spectrophotometric assay for plant pectin methyl esterase. *J. Agric. Food Chem.* 34:440.
- Hagerman, A. E., D. M. Blau, and A. L. McClure. 1985. Plate assay for determining the time of production of protease, cellulase, and pectinases by germinating fungal spores. *Anal. Biochem.* 151:334.
- Hankin, L., and S. L. Anagnostakis. 1975. The use of solid media for detection of enzyme production by fungi. *Mycologia* 67:597.
- Hankin, L., D. C. Sands, and D. E. Hill. 1974. Relation of land use to some degradative enzymatic activities of soil bacteria. *Soil Sci.* 118:38.
- Hankin, L., M. Zucker, and D. C. Sands. 1971. Improved solid medium for the detection and enumeration of pectolytic bacteria. *Appl. Microbiol.* 22:205.
- Hsu, E. J., and R. H. Vaughn. 1969. Production and catabolite repression of the constitutive polygalacturonic acid *trans*-eliminase of *Aeromonas liquefaciens*. *J. Bacteriol.* 98:172.
- Hudson, J. M., and R. W. Buescher. 1986. Relationship between degree of pectin methylation and tissue firmness of cucumber pickles. *J. Food Sci.* 51:138.
- Inoue, S., Y. Nagamatsu, and C. Hatanaka. 1984. Preparation of cross-linked pectate and its application to the purification of endopolygalacturonase of *Kluyveromyces fragilis*. *Agric. Biol. Chem.* 48:633.

28. Ishii, S. 1976. Enzymatic maceration of plant tissues by endo-pectin lyase and endo-polygalacturonase from *Aspergillus japonicus*. *Phytopathology* 66:281.
29. Jayasankar, N. P., and P. H. Graham. 1970. An agar plate method for screening and enumerating pectinolytic microorganisms. *Can. J. Microbiol.* 16:1023.
30. Juven, B. J., P. Lindner, and H. Weisslowicz. 1985. Pectin degradation in plant material by *Leuconostoc mesenteroides*. *J. Appl. Bacteriol.* 58:533.
31. Karam, N. E., and A. Belarbi. 1995. Detection of polygalacturonases and pectin esterases in lactic acid bacteria. *World J. Microbiol. Biotech.* 11:559-563.
32. Keen, N. T., and S. Tamaki. 1986. Structure of two pectate lyase genes from *Erwinia chrysanthemi* EC16 and their high-level expression in *Escherichia coli*. *J. Bacteriol.* 168:595.
33. Keen, N. T., D. Dahlbeck, B. Staskawicz, and W. Belser. 1984. Molecular cloning of pectate lyase genes from *Erwinia chrysanthemi* and their expression in *Escherichia coli*. *J. Bacteriol.* 159:825.
34. Kim, S. D., K. S. Jang, Y. A. Oh, M. J. Kim, and Y. J. Jung. 1991. Characteristics of polygalacturonases produced from *Lactobacillus acidophilus*. *J. Korean Soc. Food Nutr.* 20:488-493.
35. King, A. D., Jr., and R. H. Vaughn. 1961. Media for detecting pectolytic gram-negative bacteria associated with the softening of cucumbers, olives, and other plant tissues. *J. Food Sci.* 26:635.
36. Kotoujansky, A. 1987. Molecular genetics of pathogenesis by soft-rot erwinias. *Ann. Rev. Phytopath.* 25:405.
37. Kotoujansky, A., A. Diolez, M. Boccarda, Y. Bertheau, T. Andro, and A. Coleno. 1985. Molecular cloning of *Erwinia chrysanthemi* pectinase and cellulase structural genes. *EMBO J.* 4:781.
38. Lee, M., and J. D. MacMillan. 1968. Mode of action of pectic enzymes. I. Purification and certain properties of tomato pectinesterase. *Biochemistry* 7:4005.
39. Liao, C.-H., and J. M. Wells. 1986. Properties of *Cytophaga johnsonae* strains causing spoilage of fresh produce at food markets. *Appl. Environ. Microbiol.* 52:1261.
40. Lim, J., Y. Yamasaki, Y. Suzuki, and J. Ozawa. 1980. Multiple forms of endo-polygalacturonase from *Saccharomyces fragilis*. *Agric. Biol. Chem.* 44:473.
41. Lund, B. M. 1972. Isolation of pectolytic clostridia from potatoes. *J. Appl. Bacteriol.* 35:609.
42. McFeeters, R. F., and S. A. Armstrong. 1984. Measurement of pectin methylation in plant cell walls. *Anal. Biochem.* 139:212.
43. Miller, T. D., and M. N. Schroth. 1972. Monitoring the epiphytic population of *Erwinia amylovora* on pear with a selective medium. *Phytopathology* 62:1175-1182.
44. Mussell, H. W., and D. L. Morre. 1969. A quantitative bioassay specific for polygalacturonases. *Anal. Biochem.* 28:353.
45. Nelson, N. 1944. A photometric adaptation of the Somogyi method for the determination of glucose. *J. Biol. Chem.* 153:375.
46. Ng, H., and R. H. Vaughn. 1963. *Clostridium rubrum* sp. N. and other pectinolytic clostridia from soil. *J. Bacteriol.* 85:1104.
47. Phaff, H. J. 1966. Alpha-2,4-polygalacturonide glycanohydrolase (endo-polygalacturonase) from *Saccharomyces fragilis*, p. 636. In E. F. Neufeld and V. Ginsburg (ed.) *Methods in enzymology, complex carbohydrates*, vol. 8. Academic, New York.
48. Pierce, L., and A. H. McCain. 1992. Selective medium for isolation of pectolytic *Erwinia* sp. *Plant Dis.* 76:382-384.
49. Rao, M. N., A. A. Kembhavi, and A. Pant. 1996. Implication of tryptophan and histidine in the active site of endo-polygalacturonase from *Aspergillus ustus*: Elucidation of the reaction mechanism. *Biochem. Biophys. Acta* 1296:167-173.
50. Reid, J. L., and A. Collmer. 1985. Activity stain for rapid characterization of pectic enzymes in isoelectric focusing and sodium dodecyl sulfate-polyacrylamide gels. *Appl. Environ. Microbiol.* 50:615.
51. Robert-Baudouy, J. 1991. Molecular biology of *Erwinia*: From soft-rot to antileukaemics. *Trends Biotechnol.* 9:325-329.
52. Roberts, D. P., P. M. Berman, C. Allen, V. K. Stromberg, G. H. Lacy, and M. S. Mount. 1986. *Erwinia carotovora*: Molecular cloning of a 3.4 kilobase DNA fragment mediating production of pectate lyases. *Can. J. Plant Pathol.* 8:17.
53. Roberts, D. P., P. M. Berman, C. Allen, V. K. Stromberg, G. H. Lacy, and M. S. Mount. 1986. Requirement for two or more *Erwinia carotovora* subsp. *carotovora* pectolytic gene products for maceration of potato tuber tissue by *Escherichia coli*. *J. Bacteriol.* 167:279.
54. Rombouts, F. M. 1972. Occurrence properties of bacterial pectate lyases. *Agric. Res. Report (Versl. Landbouwk. Onderz.)* 779. Cent. for Agric. Publ. and Doc., Wageningen, The Netherlands.
55. Sands, D. C., L. Hankin, and M. Zucker. 1972. A selective medium for pectolytic fluorescent pseudomonads. *Phytopathology* 62:998.
56. Schlemmer, A. F., C. F. Ware, and N. T. Keen. 1987. Purification and characterization of a pectin lyase produced by *Pseudomonas fluorescens* W51. *J. Bacteriol.* 169:4493.
57. Starr, M. P., A. K. Chatterjee, P. B. Starr, and G. E. Buchanan. 1977. Enzymatic degradation of polygalacturonic acid by *Yersinia* and *Klebsiella* species in relation to clinical laboratory procedures. *J. Clin. Microbiol.* 6:379.
58. Vaughn, R. H., G. D. Balatsouras, G. K. York II, and C. W. Nagel. 1957. Media for detection of pectinolytic microorganisms associated with softening of cucumbers, olives, and other plant tissues. *Food Res.* 22:597.
59. Voragen, A. G. J. 1972. Characterization of pectin lyases on pectins and methyl oligogalacturonates. *Agric. Res. Report (Versl. Landbouwk. Onderz.)* 780. Cent. for Agric. Publ. and Doc., Wageningen, The Netherlands.
60. Wieringa, K. T. 1949. A method for isolating and counting pectolytic microbes. 4th Int. Cong. Microbiol. Proc., p. 482.
61. Wood, P. J., and I. R. Siddiqui. 1971. Determination of methanol and its application to measurement of pectin ester content and pectin methyl esterase activity. *Anal. Biochem.* 39:418.
62. Woodward, E. J., and K. Robinson. 1990. An improved formulation and method of preparation of crystal violet pectate medium for detection of pectolytic *Erwinia*. *Lett. Appl. Microbiol.* 10:171-173.

COMPENDIUM OF METHODS FOR THE
**MICROBIOLOGICAL
EXAMINATION
OF FOODS**

FOURTH EDITION

EDITED BY
FRANCES POUCH DOWNES
KEITH ITO



AMERICAN PUBLIC HEALTH ASSOCIATION

The American Public Health Association is an Association of individuals and organizations working to improve the public's health. It promotes the scientific and professional foundation of public health practice and policy, advocates the conditions for a healthy global society, emphasizes prevention, and enhances the ability of members to promote and protect environmental and community health.

American Public Health Association
800 I St., NW
Washington, DC 20001-3710

Mohammad N. Akhter, MD, MPH
Executive Vice President

© 2001 by the American Public Health Association

All rights reserved. This book is protected by copyright. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Sections 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4744. Requests to the Publisher for permission should be addressed to the Permissions Department, American Public Health Association, 800 I Street, NW, Washington, DC 20001-3710, fax (202) 777-2531, e-mail: <permissions@apha.org>.

10 M 3/01
Library of Congress Catalog Card Number: 99-073697

ISBN: 0-87553-175-x

Printed and bound in the United States of America.

Cover Design: Sam Dixon Design

Typesetting: Susan Westrate, Joseph R. Loehle

Set in: Palatino and Universe

Printing and Binding: Sheridan Books, Inc.

NOTE: Any discussion of medical or legal issues in this publication is being provided for informational purposes only. Nothing in this publication is intended to constitute medical or legal advice, and it should not be construed as such. This book is not intended to be and should not be used as a substitute for specific medical or legal advice, since medical and legal opinions may only be given in response to inquiries regarding specific factual situations. If medical or legal advice is desired by the reader of this book, a medical doctor or attorney should be consulted.

The use of trade names and commercial sources by authors in certain chapters of this compendium does not imply endorsement by either the APHA or the editorial board of this volume.