

Protein and Amino Acids of Sweet Potato (*Ipomoea batatas* (L.) Lam.) Fractions

Albert E. Purcell,* William M. Walter, Jr., and Francis G. Giesbrecht

Sweet potatoes contain nutritionally significant amounts of high-quality protein but usually do not have enough to provide an adequate protein-calorie ratio. A method of fractionating sweet potatoes into fiber, starch, chromoplasts, syrup, and protein is presented. Nearly half of the nitrogen contained in the sweet potato may be recovered as a concentrate containing over 80% protein. Amino acid composition of the protein shows it to be limiting in total sulfur-containing amino acids. There is an excess of lysine, suggesting usefulness as a supplement to grain products.

Sweet potato protein is of good quality (Nagase, 1957; Purcell et al., 1972; Yamamoto, 1954), but the level which ranges from 2–11%, dry basis (Cooley, 1948; Crosby, 1964; Juritz, 1921; Murthy and Swaminathan, 1954; Purcell et al., 1972) usually does not provide an adequate protein-calorie ratio. Nevertheless, the potentially high yield of sweet potato (Hernandez, 1975) suggests that it could be an important source of protein. Extraction of the protein may be required, because its even distribution within the root precludes easy production of high protein sweet potato products (Purcell et al., 1976).

We report a method for fractionating sweet potato to obtain a protein concentrate. We also report the distribution of protein among the fractions and the amino acid composition of the proteins within each fraction.

MATERIALS AND METHODS

Sweet Potatoes. Cured roots of Centennial and Jewel cultivars were obtained from the North Carolina Agricultural Experiment Station farm near Clayton, North Carolina and from a commercial packer near Wake Forest.

Fractionation. Washed, unpeeled roots were cut into 2-cm cubes, and 500 g was blended for 30 s with 1500 mL of H₂O. The puree was filtered with two layers of cheesecloth. Fibrous material which was retained by the cheesecloth was again blended with 1000 mL of H₂O and filtered. Material retained by the cheesecloth was designated as "fiber". Filtrates were screened with a 120-mesh screen to retain cell walls not trapped by the cheesecloth. Starch grains in the filtrate were sedimented by centrifugation at 1000g for 10 min. The starch was resuspended in water and allowed to settle by gravity. Wash water was discarded.

The centrifugal supernatant was heated with stirring to 65 ± 1 °C. Upon addition of 0.5% CaCl₂·2H₂O the chromoplasts precipitated. After cooling to 30–40 °C the chromoplast paste was compacted by centrifugation at 10000g for 10 min. Heating the supernatant to 95 °C coagulated the soluble proteins. After cooling to 30–40 °C, proteins were compacted by centrifugation at 10000g for 10 min. The pellet was designated "protein" and the remaining liquid as "syrup".

Chromoplast and protein pellets were separately resuspended in acetone, and filtration removed residual water. Each fraction was extracted with a 1:1 mixture of acetone-ethyl ether until the extracts were colorless. Resulting powders were dried overnight in a forced draft

oven at 70 °C. Syrup was freeze-dried. Aliquots of each fraction were used for dry matter and protein determinations.

Nitrogen Analysis. Nitrogen content of each fraction was determined by the Kjeldahl method with copper and selenium catalysts. Protein was calculated as N × 6.25.

Amino Acid Analysis. Amino acid composition of each fraction was determined by automated amino acid analysis (Spackman et al., 1958). After hydrolysis and "clean-up", the samples were injected into a Beckman 119 Amino Acid Analyzer. Tryptophan and cystine-cysteine, however, were determined colorimetrically after enzyme hydrolysis.

Acid Hydrolysis. Samples containing 6–9 mg of protein were hydrolyzed with 6 N HCl in the presence of 4% thioglycolic acid (Purcell et al., 1972; Matsubara and Sasaki, 1969). After hydrolysis, the samples were dried over NaOH in a partially evacuated desiccator, resuspended in 4 mL of 1 N HCl, and passed through an ion exchange clean-up column.

Enzyme Hydrolysis. Samples containing 60 mg of protein were weighed into 50-mL Erlenmeyer flasks, suspended in 5 mL of 0.02 M phosphate buffer (pH 4.5) and mixed with 2 mL of the same buffer containing 4 mg of amyloglucosidase (Sigma Chemical Company) and 0.5 mL of toluene as a preservative. The mixtures were incubated 16 h at 34 °C, then neutralized with 0.1 mL of 0.4 N NaOH and 5 mL of 0.2 M phosphate buffer, pH 7.5. Proteins were hydrolyzed by addition of 3 mg of pronase (Sigma *Streptomyces griseus* protease) and incubation for 24 h at 34 °C. After incubation, samples were heated to 121 °C cooled, acidified to pH 2 with 0.5 mL of HOAc, and 0.1 mL of 3 N HCl. Samples were centrifuged and the supernatant passed through a "clean-up" column.

Ion-Exchange Cleanup. Acid washed Dowex 50W resin, 200–400 mesh, 0.35 g, was suspended in 1 mL of 1 N HCl and poured into a 6-mm tube plugged with glass wool. The sample was applied when the acid no longer covered the top of the resin. After the sample had passed into the column, the column was washed three times with 1 mL of 0.5 N HOAc and twice with 1-mL portions of water. Amino acids were eluted with a mixture of 4 mL of 20% triethylamine in 20% aqueous acetone. The amino acids were dried over H₂SO₄ at 0.5 atm and taken up in 4-mL sample dilution buffer.

Measurement of Tryptophan. Tryptophan in the enzyme hydrolyzate was determined according to Spies (1968). Blanks consisting of 1 mL samples in 9 mL 21 N H₂SO₄ were made for each sample. Absorbance at 590 nm was subtracted from absorbance of the respective samples. Samples were compared to a reference consisting of Diamino benzaldehyde in 21 N H₂SO₄. The amount of tryptophan was estimated from net absorbance and a standard curve.

Determination of Cystine-Cysteine. Total cystine and cysteine were determined by modification of the

Southern Region, United States Department of Agriculture, Agricultural Research Service (A.E.P., W.M.W.), the Department of Food Science, North Carolina State University, Raleigh, North Carolina 27607 (A.E.P., W.M.W.), and the Department of Statistics, North Carolina State University, Raleigh, North Carolina 27607 (F.G.G.).

Table I. Protein Content and Recovery of Protein and Dry Matter in Each Fraction (Means and Ranges for Two Cultivars from Two Locations)

	% protein dry basis	% total recovered in fraction	
		Protein	Dry matter
Fiber	3.0 (2.7-3.5)	9.3 (8.1-10.7)	25.8 (20.4-27.8)
Starch	0.2 (0.1-0.24)	0.4 (0.2-0.6)	24.5 (21.1-27.6)
Chromoplast	42.6 (39.3-66.0)	19.1 (15.7-23.8)	5.1 (3.8-7.0)
Protein	83.4 (80.7-88.0)	48.4 (44.4-50.5)	5.0 (3.8-6.4)
Syrup	4.7 (3.9-6.0)	22.7 (18.7-27.1)	39.5 (34.2-46.6)

method of Manning et al. (1971). A 2-mL aliquot of the enzyme hydrolyzate was mixed with 1 mL of 2.5% sodium borohydride in 0.2 M PO₄ buffer, pH 7.5, and held at 40 °C. After 30 min 1 mL of acidulant, 10.4 g of NaH₂PO₄ per 100 mL of 0.1 N HCl, and 1 mL of acetone were added, and the mixture was purged with N₂ for 6 min. The mixture was neutralized with 1 mL of 0.1 N NaOH and color was developed by addition of 1 mL of 0.2 M PO₄ buffer (pH 7.5) containing 2 mg of 5,5' dithiobis(2-nitrobenzoic acid) (DTNB). Volume was adjusted to 7.0 mL, samples were held at 25 °C for 40 min, and absorbance was measured at 420 nm. Controls were made for each sample, consisting of all reagents except DTNB. All samples were read against a blank made by using distilled water instead of sample. Absorbance of controls were subtracted from absorbance of samples and the amount of cysteine was estimated by use of a standard curve.

Statistical Procedures. Analysis of variance (AOV) for each amino acid was calculated to determine fraction effects upon content. AOV-derived least-significant differences (LSD) were used to compare fraction means.

RESULTS AND DISCUSSION

The fiber fraction contained an average of 25.8% of the total dry matter (Table I). Microscopic examination showed that the fraction consisted mainly of cell walls and vascular fibers. There were a few unruptured cells, starch grains, and chromoplasts. Further blending and washing caused only small changes in the composition of this fraction. The major source of protein in this fraction was probably the cell walls.

The starch sedimented by centrifugation contained some chromoplasts and nuclei. These were removed by subsequent suspension and sedimentation of the starch. These nonstarch components represented an insignificant part of the chromoplast fraction and were discarded due to great dilution. Starch could not be recovered quantitatively by the procedure used because some of the smaller grains would not settle before nuclei and chromoplasts. Average recovery of dry matter in the starch was 24.5% (Table I). Protein concentration of this fraction averaged 0.2%, some of which may have been from nonstarch particulates.

The chromoplast fraction contained all of the carotene not removed with the starch. All microstructure was destroyed, but occasional small starch grains which had not been removed by centrifugation were present. The amount and composition of this fraction were more variable than those of the other fractions (Table I). It contained an average of 19.1% protein and around 30% lipids (data not tabulated).

The protein concentrate, ranging from 80-88% protein contained about half of the total protein (Table I). It is believed to be mainly cytoplasmic proteins.

Table II. Amino Acid Composites of Various Sweet Potato Fractions as Means of Two Cultivars from Two Locations

	g amino acid/16 g of N recovered				
	Fiber	Chromo-plasts	Protein	Syrup	LSD _{0.05}
Tryptophan	0.6	1.1	1.4	0.3*	0.8
Lysine	4.7**	7.0	8.3	2.4*	2.0
Histidine	1.0*	2.1	2.7	0.8*	0.6
Arginine	1.2*	6.6	8.5	1.1*	2.2
Aspartic acid	22.6	15.6	12.3	27.0*	13.3
Threonine	6.8	6.0	4.5	3.2	N.S.
Serine	6.9*	6.7*	4.0	2.9	1.8
Glutamic acid	11.5	13.7	6.4	10.9	N.S.
Proline	5.7*	5.8*	3.3	1.8	1.5
Glycine	5.5*	6.2*	3.4	2.2	1.2
Alanine	6.5*	7.1*	3.7	2.5	1.4
Half-cystine	.03*	.06*	0.3	.07*	.11
Valine	9.1*	8.4*	5.6	2.6*	2.6
Methionine	0.7*	2.8*	1.7	0.5*	0.9*
Isoleucine	6.0	7.5*	3.8	1.8	2.5
Leucine	9.2*	9.8*	5.0	2.6	2.8
Tyrosine	2.9*	5.4*	4.0	0.8*	1.0
Phenylalanine	6.6	7.3	5.8	2.4*	2.8
% of N recov.	82.0	99.3	79.0	48.4	

* (*) significantly different (0.05) than concentrate.

Table III. Lysine and Histidine Contents of Fractions from Centennial and Jewel Cultivars

	Lysine, g/16 g of N		Histidine, g/16 g of N	
	Centennial	Jewel	Centennial	Jewel
Fiber	3.37	6.06	0.84	1.11
Chromoplast	8.48	5.56	2.79	1.47
Protein	6.75	9.77	2.54	2.84
Syrup	2.72	2.01	0.73	0.81
LSD _{0.05}		2.88		0.79

The syrup fraction contained 34-47% of the total dry matter (Table I) and 19-27% of the protein. Concentration of protein (N × 6.25) appeared to be less in Centennial than in Jewel.

Amino acid composition of the fractions varied significantly (Table II), but differences due to location were not significant. There were significant cultivar-fraction interactions for histidine and lysine (Table III). The chromoplast fraction of Centennial contained significantly more histidine and lysine than the same fraction from Jewel. The protein concentrate fraction of Centennial contained significantly less lysine than that of Jewel.

The syrup fraction contained the lowest amount of essential amino acids. Aspartic and glutamic acid accounted for 38% of this fraction (Table II). All essential amino acid levels were below those of the FAO reference protein (Burton, 1965). The fiber fraction contained more amino acids with lipophilic groups, i.e., alanine, valine, isoleucine, leucine, and phenylalanine, than the protein concentrate; this suggests that much of the protein in the fiber fraction is contained in cell walls. For nutritional purposes, the fiber fraction is deficient in tryptophan and sulfur-containing amino acids. Amino acid composition of the chromoplast fraction suggests that the protein is even more lipophilic than the protein of the fiber fraction. Tryptophan and sulfur-containing amino acids limit the nutritional value of this fraction.

Nutritional value of the protein concentrate is more limited due to deficiency of sulfur-containing amino acids than soy protein. The abundance of lysine suggests that this fraction may be more valuable in supplementing grain protein than soy.

With steadily increasing yield of sweet potatoes and trends toward mechanization, sweet potato may eventually compete with corn as a source of commercial starch. The protein could be retained and used in human diets.

LITERATURE CITED

- Burton, B. T., "The Heinz Handbook of Nutrition", 2nd ed, McGraw-Hill, New York, N.Y., 1965, p 144.
Cooley, J. S., *Econ. Botan.* 2, 83 (1948).
Crosby, D. G., *J. Food Sci.* 29, 187 (1964).
Hernandez, T., National Sweet Potato Collaborators Groups Progress Report, material assembled at Louisiana State University, 1975.
Juritz, C. I., *Dep. Agric. S. Afr.* 2, 340 (1921).
Manning, P. B., Jenness, R., Morr, C. V., Presentation Annual Meeting Dairy Science Association, 1971.
Matsubara, H., Sasaki, R. M., *Biochem. Biophys. Res. Commun.* 35, 175 (1969).

- Murthy, H. B. N., Swaminathan, M., *Curr. Sci.* 23, 14 (1954).
Nagase, T., *Fukuoka Igaku Zasshi* 48, 1828 (1957).
Purcell, A. E., Swaisgood, H. E., Pope, D. T., Jr., *Am. Soc. Hortic. Sci.* 97, 30 (1972).
Purcell, A. E., Walter, W. M., Jr., Giesbrecht, F. G., *J. Agric. Food Chem.* 24, 64 (1976).
Spackman, D. H., Stein, W. H., Moore, S., *Anal. Chem.* 30, 1190 (1958).
Spies, J. R., *J. Agric. Food Chem.* 16, 1412 (1968).
Yamamoto, Y., *Eiyo To Shokuryo* 7, 112 (1954).

Received for review November 10, 1977. Accepted January 6, 1978. Paper No. 5414, Journal Series of the North Carolina Agricultural Experiment Station, Raleigh, N.C. Mention of a trademark or proprietary product does not constitute a guarantee or warranty of the product by the U.S. Department of Agriculture of North Carolina Agricultural Experiment Station, nor does it imply approval to the exclusion of other products that may be suitable.