

## Analysis of genetic diversity in an invasive population of Asian long-horned beetles in Ontario, Canada

Maureen E. Carter<sup>1</sup>

Department of Entomology, Comstock Hall, Cornell University, Ithaca, New York 14853-2601, United States of America

Michael T. Smith

United States Department of Agriculture, Agricultural Research Service, Beneficial Insect Introduction Research Unit, Newark, Delaware 19713, United States of America

Jean J. Turgeon

Great Lakes Forestry Centre, Canadian Forest Service, Natural Resources Canada, Sault Ste. Marie, Ontario, Canada P6A 2E5

Richard G. Harrison

Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York 14853, United States of America

**Abstract**—Adult Asian long-horned beetles, *Anoplophora glabripennis* (Motschulsky) (Coleoptera: Cerambycidae: Lamiinae), were discovered in Ontario, Canada, in 2003 in the vicinity of a commercial warehouse. Trees were heavily scarred with signs of attack and larvae and adult beetles were common, suggesting that there had been multiple generations at the site. We amplified 16 microsatellite loci from 326 beetles to examine genetic diversity in this population. Based on Hardy–Weinberg equilibrium, 6 of 16 loci were monomorphic and 8 were not, indicating nonrandom mating. Measures of microsatellite genetic diversity and mitochondrial DNA haplotype diversity were significantly lower than those in *A. glabripennis* from China and Korea but were not significantly different from those in the New York City population. The proportion of different multilocus genotypes in the Ontario population was lower than in the populations in New York City and Linden, New Jersey. These results suggest that limited genetic diversity in the Ontario population has not hampered reproduction of this invasive insect. This genetic signature is common in other invasive species, likely because a population is founded by a few closely related individuals, or a large founding population suffers subsequent genetic bottlenecks.

**Résumé**—Des adultes du longicorne asiatique, *Anoplophora glabripennis* (Motschulsky) (Coleoptera : Cerambycidae : Lamiinae), ont été découverts en Ontario, Canada, en 2003 près d'un entrepôt commercial. Les arbres portaient de fortes lacérations dues à l'attaque des coléoptères et les larves et les adultes y étaient nombreux, ce qui laisse croire qu'il y avait eu plusieurs générations à ce site. Nous avons examiné la diversité génétique de la population en amplifiant 16 locus microsatellites chez 326 coléoptères. Six des 16 locus sont monomorphes et 8 locus ne suivent pas l'équilibre Hardy–Weinberg, ce qui indique que les accouplements ne sont pas aléatoires. Les mesures de diversité génétique des microsatellites et de diversité des haplotypes d'ADN mitochondrial sont significativement plus faibles que celles d'*A. glabripennis* en Chine et en Corée, mais elles ne diffèrent pas significativement des mesures de la population de la ville de New York. La proportion de génotypes différents à locus multiples est plus faible dans la population d'Ontario que dans les populations de la ville de New York et de Linden (New Jersey). Ces résultats indiquent que la diversité génétique restreinte de la population d'Ontario n'a pas nui à la reproduction de cet insecte envahissant. Une telle signature génétique est commune chez d'autres espèces envahissantes, vraisemblablement parce que la population est fondée sur un petit nombre d'individus fortement

Received 5 February 2009. Accepted 6 July 2009.

<sup>1</sup> Corresponding author (e-mail: mec8@cornell.edu).  
doi: 10.4039/n09-026

apparentés ou alors parce qu'une population fondatrice de grande taille subit plus tard des goulots d'étranglement génétiques.

[Traduit par la Rédaction]

## Introduction

The Asian long-horned beetle, *Anoplophora glabripennis* (Motschulsky) (Coleoptera: Cerambycidae: Lamiinae), is a wood-borer native to China and Korea (Cavey *et al.* 1998; Lingafelter and Hoebeke 2002). In China *A. glabripennis* is a destructive pest of man-made forests (plantations) and windrows of one to several tree species planted to shelter urban, rural, and agricultural areas from sandstorms and provide a local resource for timber and paper products (Moore and Russell 1990; Uchida *et al.* 2005). In contrast, *A. glabripennis* is rare in South Korea, occurring only in sparse populations along edges of species-rich deciduous forest stands (Williams *et al.* 2004).


Female beetles chew through bark on trunks and branches and lay eggs singly in the inner bark (Xiao 1992; Cavey *et al.* 1998). Young larvae feed at the inner bark – sapwood interface for 14–28 days (Keena 2005; Hunter *et al.* 2009), slowly destroying the vascular system and cutting off translocation (Ric *et al.* 2007). Older larvae tunnel into the heartwood, causing branch dieback, structural deterioration, and often tree mortality (Haack *et al.* 1997; Ric *et al.* 2007). Pupation takes place in the wood, where teneral adults remain for up to 7 days before chewing an exit hole and emerging (Xiao 1992; Lingafelter and Hoebeke 2002). Adult beetles feed on twigs and leaf petioles and primary leaf veins (Keena 2002; Smith *et al.* 2002; Ric *et al.* 2007). Newly eclosed females become sexually mature in 9–15 days (Keena 2002; Smith *et al.* 2002); development from egg to adult requires approximately 12–24 months. In China, *A. glabripennis* overwinter as eggs or larvae, with egg development and larval feeding resuming in the spring. Pupation occurs from May through July, with adult emergence peaking from mid-June to late July (Pan 2005). Larvae that are still young in the fall overwinter a second time,

*i.e.*, a 2-year life cycle occurs in colder climates. Although high levels of infestation occur in weak trees grown in monoculture (Yin and Lu 2005), the beetle also attacks apparently healthy host trees (Haack *et al.* 1997). Adult males and females mated repeatedly in field (Xiao 1992) and greenhouse studies (Morewood *et al.* 2004). In the laboratory, females lay between 30 and 127 eggs, but the number varies among host tree species (Smith *et al.* 2002).

Several established populations have been discovered in the United States of America, the first in Brooklyn, New York, in 1996 (Haack *et al.* 1997). Subsequently, between 1998 and 2008, the beetle was found at other sites in New York City and elsewhere in New York State (Long Island, Prall's Island, Staten Island), New Jersey (Jersey City, Carteret, Linden), Illinois (Chicago; Poland *et al.* 1998), Massachusetts (Worcester), and California (Sacramento). This species has also been intercepted or introduced multiple times in Europe (Austria (Tomiczek *et al.* 2002), France, Germany (Hérard *et al.* 2006), the Czech Republic (Sabol 2006), and Italy (Maspero *et al.* 2007)). *Anoplophora glabripennis* is usually introduced through the transportation of solid-wood packing material from Asia to North American warehouses, from where adult beetles escape to nearby trees (Haack *et al.* 1997).

In its native range the primary hosts of *A. glabripennis* are poplars and willows (*Populus* L. and *Salix* L., Salicaceae), maples (*Acer* L., Aceraceae), and elms (*Ulmus* L., Ulmaceae), but hosts also include lindens (*Tilia* L., Tiliaceae), Russian olive and buckthorns (*Eleagnus angustifolia* L. and *Hippophae* L., Elaeagnaceae), birches (*Betula* L., Betulaceae), horse chestnuts (*Aesculus* L., Hippocastanaceae), the London planetree (*Platanus hybrida* Brot., Platanaceae), and the goldenrain tree (*Koelreuteria paniculata* Laxm., Sapindaceae) (Wang 2004). Outside its native range, hosts include maples, willows,

**Fig. 1.** Map of the Asian long-horned beetle (ALHB) regulated area in Vaughan and Toronto, Ontario, showing the location and number of beetles per tree that were sampled for genetic analysis.



elms, birches, horse chestnuts, mountain ashes (*Sorbus* L., Rosaceae), the London planetree, and species that are not attacked in China (Nowak *et al.* 2001; Morewood *et al.* 2003; Sawyer 2003; Hajek and Kalb 2007; Ric *et al.* 2007). Various potential hosts found in the currently regulated area in Canada have not been attacked: *Prunus* L., *Crataegus* L., and *Malus* Mill. (Rosaceae) among others (Turgeon *et al.* 2007).

*Anoplophora glabripennis* was first intercepted in Canada in 1998 in packing material sent from China to a warehouse in Waterloo, Ontario (Sellers 2004). In Ontario, an infestation was discovered in the Greater Toronto Area in September 2003 in an industrial area on the boundary between Toronto and Vaughan, near the intersection of Weston Road and Steeles Avenue (Fig. 1). By the time it was discovered, hundreds of trees had been infested. In February 2004, a regulated area was

established to prevent the movement of infested wood from the region (Canadian Food Inspection Agency 2003). As of October 2007, approximately 27 400 host trees had been removed, including more than 600 known to be infested (North American Plant Protection Organization 2007). Additional infested trees in Toronto are being removed when found, along with any potential hosts within 200–400 m (Canadian Food Inspection Agency 2008).

Molecular markers can be used to characterize patterns of genetic variation in invasive species, revealing the signature of an expanding population and providing data about founder-population size and subsequent population bottlenecks, factors that may be important in population growth and spread (Wares *et al.* 2005; Carroll 2007; Eales *et al.* 2008; Ficetola *et al.* 2008). Additive genetic variance may increase after a population bottleneck, allowing a rapid response to

selection (Wade *et al.* 1996; Cheverud *et al.* 1999; Naciri-Graven and Goudet 2008). The projected annual economic impact of *A. glabripennis* on Canadian resources has been estimated to be over \$331 million (Colautti *et al.* 2006). It is important to look for evidence of multiple founding events or population bottlenecks, as these different scenarios could influence the choice of prevention and eradication programs.

We evaluated the amount of genetic diversity in the *A. glabripennis* population in Canada and compared it with the known genetic diversity in populations of this beetle in the United States of America, China, and Korea. To estimate variation, we amplified microsatellite DNA markers consisting of tandem repetitive units 1–6 base pairs (bp) long (Oliveira *et al.* 2006). High rates of mutation (Jarne and Lagoda 1996) of these presumably neutral markers make them particularly useful for inferring current population structure and recent population history (Roderick and Navajas 2003; Schlötterer 2004). In a previous study (Carter *et al.* 2009b) we amplified and sequenced DNA from a subset of the *A. glabripennis* specimens from Ontario for mitochondrial DNA COI and COII genes, using species-specific primers to determine the number and source of haplotypes. Here we also consider mitochondrial DNA haplotype diversity in the Ontario *A. glabripennis* population in evaluating the history of the introduction.

## Methods and materials


Emergence period and seasonal occurrence and abundance of *A. glabripennis* in Ontario are currently unknown. However, a day-degree model predicting emergence of adult *A. glabripennis* in China and associated data were used to create an emergence model for the Ontario population, based on accumulated day-degrees: emergence rate (%) =  $\exp(-\exp(-0.004 \times \text{day-degrees} + 3.5)) \times 100$  (Smith *et al.* 2004). The Historical Weather Database for Toronto, Ontario (1978–2008), based on data obtained from Toronto Lester B. Pearson International Airport, was used to find the mean date

corresponding to accumulated day-degrees (Fig. 2). To find the date of a specific day-degree accumulation (temperature),  $X$ , the accumulated day-degrees between  $(X - 5)$  and  $(X + 5)$  were rounded to the specific day-degree. For example, to find the corresponding mean dates for 400 day-degrees, all observations between  $395 \leq \text{day-degree} \leq 405$  for each year from 1978 to 2008 were pooled to create a new variable, “rounded-day-degree” (= 400). From the resulting data set of “rounded-day-degrees” the mean dates were calculated. Using the same method, the mean dates for rounded-day-degrees for 450–1900 were also calculated. These data were used to generate a plot of accumulated day-degrees ( $^{\circ}\text{C}$ ) (first  $x$  axis) versus percent emergence ( $y$  axis) for Toronto. A curve-smoothing function was used to generate an arc connecting all predicted points. The date corresponding to each rounded-day-degree was applied to this plot as a second  $x$  axis.

Between 26 September 2003 and 2 September 2005, live adult and larval *A. glabripennis* and eggs were collected in the regulated area (Fig. 1) and placed in 95% ethanol, and their host was recorded (Table 1). Using DNeasy tissue kits (QIAGEN), total DNA was extracted from adult thoracic muscle tissue, from a segment of the larval abdomen, or by crushing a whole egg. A subset of samples has been deposited as vouchers with the Cornell University Insect Collection (lot No. 1262).

DNA samples from 57 *A. glabripennis* were amplified and sequenced for mitochondrial DNA using six species-specific primers at optimized annealing temperatures (Carter *et al.* 2009c). We sequenced 1317 bp of cytochrome *c* oxidase 1, 65 bp of tRNA-leucine, and 224 bp of cytochrome *c* oxidase 2 mitochondrial DNA genes. Each PCR was run in a volume of 14  $\mu\text{L}$  containing 6.7  $\mu\text{L}$  of water, 1.5  $\mu\text{L}$  of  $10 \times$  PCR buffer (20 mmol/L Tris-HCl (pH 8.4), 500 mmol/L KCl (Invitrogen, Carlsbad, California)), 3  $\mu\text{L}$  of 2.5 mmol/L dNTP, 0.75  $\mu\text{L}$  of 50 mmol/L  $\text{MgCl}_2$ , 0.06  $\mu\text{L}$  of *Taq* DNA polymerase (Invitrogen), 0.6  $\mu\text{L}$  of primers at 10 mmol/L, and 20 ng of DNA. PCR products were cleaned with PCR mini-elute kits (QIAGEN).

**Fig. 2.** Population abundance of Asian long-horned beetles, *Anoplophora glabripennis*, over accumulated day-degrees in Toronto, based on the following estimated model: emergence rate (%) =  $\exp(-\exp(-0.004 \times \text{day-degrees} + 3.5)) \times 100$  (see Smith *et al.* 2004); SE = 0.0002 for  $r$  and 0.2 for  $b$ ;  $R^2 = 0.88$ . The model was used to calculate the predicted values on the  $y$  axis for accumulated proportions of adult beetle population emergence (%) during the corresponding day-degree interval between two points. The  $x$  axis shows the accumulated day-degrees and the corresponding mean date (see Methods).



**Table 1.** Numbers of trees and numbers of Asian long-horned beetles, *Anoplophora glabripennis*, collected for genetic analysis between 26 September 2003 and 2 September 2005 in Vaughan and Toronto, Ontario, where the host trees were known to be present.

| Tree genus     | No. of trees | No. of beetles |
|----------------|--------------|----------------|
| <i>Acer</i>    | 170          | 292            |
| <i>Betula</i>  | 4            | 7              |
| <i>Populus</i> | 1            | 1              |
| <i>Salix</i>   | 3            | 3              |
| <i>Ulmus</i>   | 2            | 2              |

Five-microlitre sequencing reactions contained 1.88  $\mu\text{L}$  of water, 0.25  $\mu\text{L}$  of 5M Betaine (Sigma), 1  $\mu\text{L}$  of Ready Reaction and 5 $\times$  buffer (Applied Biosystems), 0.12  $\mu\text{L}$  of 10 mmol/L primer, and 1  $\mu\text{L}$  of PCR products. Sequencing reactions were cleaned with Sephadex (Sigma) and directly sequenced

on a ABI 3730 sequencer. Sequences were assembled into contiguous arrays, edited and trimmed using Seqman, and aligned using Megalign (Lasergene 7). Approximately 1750 bp of sequence were trimmed to 1606 bp without gaps. Using a  $z$  test the proportion of different haplotypes in the population from Ontario, Canada was tested against the proportions found in populations in China, Korea, and the United States of America.

We genotyped 326 *A. glabripennis* at 16 microsatellite loci. Reaction mixtures consisted of universal fluorescent sequence primers with reverse-specific primers at 3.2  $\mu\text{mol/L}$  combined with nonfluorescent forward-specific primers at 0.8  $\mu\text{mol/L}$ , 20 ng of DNA, 1 $\times$  PCR buffer (20 mmol/L Tris-HCl (pH 8.4), 500 mmol/L KCl (Invitrogen)), 1.5 mmol/L  $\text{MgCl}_2$ , 1M betaine, 0.2 mmol/L each dNTP, and 0.5 U of *Taq* (Promega). Primer sequences for 15 primers and PCR

conditions are listed in Carter *et al.* (2009a). Primers for an additional locus, locus 15, were GGCCTATTTTGATGCGAGTG (forward) and GGCCTACCTGCTACACAGC (reverse) fluorescently labeled with PET (Applied Biosystems) and amplified following the methodology of Carter *et al.* (2009c).

Diversity information was calculated in PowerMarker (Liu and Muse 1995) and included the number of alleles and genotypes and frequency of the most common (major) allele. Allelic richness was calculated in FSTAT (Goudet 1995). The observed number of alleles in a sample is highly dependent on sample size, therefore it is more accurate to estimate allelic richness (number of alleles at a locus independent of sample size). Allelic richness is estimated from  $n$  samples genotyped at all loci (no missing data). Also calculated were observed heterozygosity, fraction of individuals in a population with two different alleles at a locus, and expected heterozygosity (an estimate of all individuals heterozygous at a locus, calculated from allele frequencies at that locus). A lower than expected observed heterozygosity suggests a deviation from Hardy–Weinberg equilibrium due to an external factor such as migration, selection, inbreeding, or drift. To test the null hypothesis of a random union of gametes, we calculated deviations from Hardy–Weinberg equilibrium ( $F_{IS}$ ; exact Hardy–Weinberg test) for each locus, and genotypic disequilibrium (Fisher's method) for each locus pair, using Genepop (Raymond and Rousset 1995). Two group comparisons of mean allelic richness and mean observed and expected heterozygosities were calculated in FSTAT (Goudet 1995) to compare data sets from China plus Korea, United States of America, and Canada. We counted the different genotypes, *i.e.*, that differ by one or more alleles across all 16 loci, in the Canadian population and in populations in the United States of America using Microsoft tools (Park 2001). With a  $z$  test we compared the proportions of two different groups for the number of different genotypes between New York City and Linden, New Jersey, between New York City and Toronto, and between Linden and Toronto.

To make inferences about the source of the Ontario population, we compared the multi-locus genotypes of beetles from Ontario with those of reference populations from provinces of China and Korea (each province was labeled a population), using GeneClass2 (Piry *et al.* 2004). Using a resampling method (Rannala and Mountain 1997), the likelihood of assignment to a reference population was calculated for each individual. Individuals were assigned to reference populations based on the largest likelihood score. For each individual beetle, an assignment was made only if the probability of assignment was 90% with a statistical confidence of 95% (threshold 0.05). Only beetles with all 32 alleles (no missing data) were used. We repeated the assignment test for each individual beetle from Canada, with the reference data set partitioned into four clusters determined by means of a model-based clustering method for inferring population structure (*structure*; Falush *et al.* 2003).

## Results

We report here the predicted seasonal phenology of adult *A. glabripennis* in Toronto (Fig. 2) and the results from two independent genetic data sets to determine the genetic diversity of the Toronto population of *A. glabripennis* and the implications of population establishment and spread.

The day-degree model predicted peak abundance of adult *A. glabripennis* in Toronto to be between approximately 850 and 950 accumulated day-degrees, coinciding with the 3rd and 4th weeks of August.

For the microsatellite markers, allele number ranged from 1 to 4, with allelic richness from 1 to 3.76 (Table 2). Of 44 pairwise comparisons for linkage disequilibrium, 5 were significant ( $P \leq 0.05$  with Bonferroni correction). All other pairwise comparisons (76) involved monomorphic loci. For 9 of the 10 non-monomorphic loci, observed heterozygosity was less than expected heterozygosity (Table 2). For 8 of these 10 loci, the calculated  $F_{IS}$  value indicated a significant probability that the observed sample was not in Hardy–Weinberg equilibrium, *i.e.*, a non-random union of gametes.

**Table 2.** Number of alleles ( $N_A$ ), number of genotypes ( $N_G$ ), major allele frequencies (MAF), expected heterozygosity ( $H_E$ ), observed heterozygosity ( $H_O$ ), and allelic richness (AR) for 16 microsatellite loci in *Anoplophora glabripennis* collected in Vaughan and Toronto.

| Locus | $N_A^*$ | $N_G^*$ | MAF* | $H_E^*$ | $H_O^*$ | AR <sup>†</sup> | $F_{IS}^{\ddagger}$ | $P^{\S}$       |
|-------|---------|---------|------|---------|---------|-----------------|---------------------|----------------|
| alb10 | 1 (326) | 1       | 1    | 0       | 0       | 1               | —                   | —              |
| alb53 | 2 (326) | 3       | 0.64 | 0.45    | 0.38    | 2               | 0.156               | <b>0.005</b>   |
| alb21 | 1 (324) | 1       | 1    | 0       | 0       | 1               | —                   | —              |
| alb59 | 1 (326) | 1       | 1    | 0       | 0       | 1               | —                   | —              |
| alb38 | 3 (325) | 4       | 0.64 | 0.46    | 0.4     | 2.96            | 0.119               | <b>0.003</b>   |
| alb77 | 4 (261) | 4       | 0.98 | 0.34    | 0.01    | 4               | 1                   | < <b>0.001</b> |
| alb14 | 2 (303) | 3       | 0.69 | 0.49    | 0.26    | 2               | 0.319               | < <b>0.001</b> |
| alb30 | 2 (324) | 3       | 0.62 | 0.47    | 0.44    | 2               | 0.048               | 0.425          |
| alb15 | 1 (326) | 1       | 1    | 0       | 0       | 1               | —                   | —              |
| alb24 | 1 (325) | 1       | 1    | 0       | 0       | 1               | —                   | —              |
| alb9  | 4 (325) | 4       | 0.58 | 0.49    | 0.4     | 3.76            | 0.176               | < <b>0.001</b> |
| alb19 | 2 (325) | 2       | 0.99 | 0.01    | 0       | 1.96            | 1                   | <b>0.002</b>   |
| alb35 | 4 (325) | 5       | 0.55 | 0.5     | 0.01    | 3.61            | 0.988               | < <b>0.001</b> |
| alb40 | 1 (325) | 1       | 1    | 0       | 0       | 1               | —                   | —              |
| alb43 | 2 (310) | 3       | 0.56 | 0.53    | 0.53    | 2               | 0.019               | 0.823          |
| alb44 | 3 (324) | 6       | 0.47 | 0.62    | 0.18    | 3               | 0.136               | <b>0.011</b>   |

\*The total number of beetles sampled was 326. The number in parentheses is the number of beetles that amplified per locus.

<sup>†</sup>The minimum number of beetles sampled was 261 (locus 77).

<sup>‡</sup>According to Weir and Cockerham (1984).

<sup>§</sup>Exact Hardy–Weinberg test; values in boldface type are significantly different.

For the 230 individuals from Ontario with no missing data, there were 13 different microsatellite genotypes (allele combinations) across all 16 loci (Table 3). We refer to nonamplification of alleles at any locus as missing data. Common reasons for missing data include mutations at primer sites or PCR inhibition due to contaminants found in DNA or reagents, or DNA shearing at the amplification site. The Ontario population had fewer different genotypes relative to the number of *A. glabripennis* sampled than other invasive populations (Table 3). Pairwise comparisons of the proportions of different genotypes yielded the following results: the New York City population differed from the Toronto ( $z = 16.6$ ,  $P \leq 0.01$ ) and Linden ( $z = 11.038$ ,  $P \leq 0.01$ ) populations, and the Toronto population differed from the Linden population ( $z = 3.125$ ,  $P \leq 0.01$ ) (Table 3). However, allelic richness and observed and expected heterozygosities were not significantly lower in the *A. glabripennis* population from Toronto than in populations in the United States of America, but were significantly

**Table 3.** Numbers and proportions of different genotypes found in North American populations of *Anoplophora glabripennis*.

|                                 | No. of genotypes tested* | No. of different genotypes | Percentage <sup>†</sup> |
|---------------------------------|--------------------------|----------------------------|-------------------------|
| <b>United States of America</b> |                          |                            |                         |
| Long Island, New York           | 33                       | 33                         | 100                     |
| New York City, New York         | 103                      | 103                        | 100 <sup>a</sup>        |
| Jersey City, New Jersey         | 23                       | 23                         | 100                     |
| Carteret, New Jersey            | 22                       | 22                         | 100                     |
| Linden, New Jersey              | 72                       | 14                         | 19.4 <sup>b</sup>       |
| <b>Canada</b>                   |                          |                            |                         |
| Toronto, Ontario                | 230                      | 13                         | 5.7 <sup>c</sup>        |

\*Only beetles with no missing genotype data were tested.

<sup>†</sup>The number of different genotypes as a percentage of the number tested. Values followed by a different letter show significantly different two by two group proportions ( $z$  test,  $P \leq 0.01$ ).

**Table 4.** Proportions of different mitochondrial DNA haplotypes and indices of microsatellite polymorphism in populations of *Anoplophora glabripennis* from Asia, the United States of America, and Toronto, Ontario.

| Collecting locality      | <i>H/N</i> <sup>*</sup> | Percentage <sup>†</sup> | AR <sup>‡</sup> | <i>H</i> <sub>O</sub> <sup>§</sup> | <i>H</i> <sub>E</sub> <sup>¶</sup> |
|--------------------------|-------------------------|-------------------------|-----------------|------------------------------------|------------------------------------|
| China and Korea          | 37/124                  | 29.8 <i>a</i>           | 3.28 <i>a</i>   | 0.29 <i>a</i>                      | 0.47 <i>a</i>                      |
| United States of America | 11/258                  | 0.04 <i>b</i>           | 2.21 <i>b</i>   | 0.15 <i>b</i>                      | 0.27 <i>b</i>                      |
| Toronto                  | 1/57                    | 0.02 <i>b</i>           | 2.17 <i>b</i>   | 0.16 <i>b</i>                      | 0.27 <i>b</i>                      |

**Note:** Values within a category followed by a different letter are significantly different (FSTAT:  $P \leq 0.05$ , one-sided test).

<sup>\*</sup>Number of different haplotypes (*H*) out of the total number sequenced (*N*).

<sup>†</sup>Proportion of different haplotypes in the population. Values followed by a different letter show significantly different two by two group proportions (*z* test,  $P \leq 0.01$ ).

<sup>‡</sup>Mean allelic richness over all 16 loci.

<sup>§</sup>Mean observed heterozygosity over all 16 loci.

<sup>¶</sup>Mean expected heterozygosity over all 16 loci.

lower than in populations in China plus Korea (Table 4).

Using a threshold confidence level of 90%, 36 beetles from the Toronto population were assigned to Shanxi Province, 14 to Liaoning Province, and 2 to Hubei Province. No other beetles could be assigned with 90% confidence. When the assignment test was repeated using the four clusters of beetles from China and Korea documented in a previous *structure* analysis as reference populations (Carter *et al.* (2009c), 165 beetles (72%) from Toronto were assigned to a single cluster characterized by beetles from the contiguous provinces of Hebei, Inner Mongolia, Liaoning, Shanxi, and Tianjin in north-central China. Four beetles (<1%) were assigned to a cluster characterized by beetles from the contiguous provinces of Anhui, Henan, Hubei, Jiangxi, and Shandong in eastern and central China.

Mitochondrial DNA sequences of 1606 bp of COI and COII genes were identical for the 57 *A. glabripennis* collected in the regulated area, *i.e.*, there was a complete absence of mitochondrial DNA sequence variation (Carter *et al.* (2009b). *Anoplophora glabripennis* collected from Canada exhibited less variation in mitochondrial DNA than invasive populations in the United States of America, which themselves varied far less than *A. glabripennis* populations collected in China and Korea (Table 4). A test of proportions of different haplotypes from the three collection locations showed that China plus Korea populations differ from those in the United

States of America ( $z = 6.9$ ,  $P \leq 0.01$ ), China plus Korea populations differ from the population in Toronto ( $z = 4.1$ ,  $P \leq 0.01$ ), and Toronto populations did not differ from those in the United States of America ( $z = 0.51$ , ns).

## Discussion

Emergence of adult *A. glabripennis* peaked at similar accumulated day-degrees in Toronto (latitude 43.4°N) as in Liu Hua, Gansu Province, China (latitude 35°N; 800–900 day-degrees; Smith *et al.* 2004). This day-degree accumulation occurs in mid-August in Toronto, whereas the same day-degree accumulation occurs in mid-July in Gansu Province. This agrees with the assignment of Toronto beetle genotypes to genotypes of beetles from the cluster of more northerly provinces of China (at latitudes more similar to that of Toronto) rather than to genotypes from Jing tai, Gansu Province (latitude 37.2°N; Carter *et al.* (2009c).

Allelic variation is often critical for adaptation to environmental changes. Genetic variation in the Ontario population of *A. glabripennis* is limited; the greatest microsatellite polymorphism observed was four alleles. This variation could be derived from a minimum sample of two individuals. The Ontario population is also fixed for a single mitochondrial DNA haplotype. Our data suggest that the Toronto population is derived from a small number of beetles, possibly one multiply inseminated female.

Polymorphism indices from tests for the 16 microsatellite markers for the Toronto population were not significantly different from similar measures for populations in the United States of America, but were significantly different from those for populations in China and Korea (Table 4). This suggests a general pattern of low genetic diversity of nuclear markers for invasive populations of *A. glabripennis*. However, the Toronto population of *A. glabripennis* exhibited reduced multilocus genotypic diversity compared with populations in the United States of America. In the Long Island, New York City, Jersey City, and Carteret populations, all individuals sampled had different microsatellite genotypes, a different pattern from that seen in Toronto, where many beetles share the same genotype. This could indicate that the Toronto population had fewer founding individuals and (or) suffered more severe population bottlenecks than did populations in the United States of America.

The mitochondrial DNA haplotype of the Toronto population has not yet been found in samples from China, Korea, or the United States of America, but has been found in two beetles collected in Neukirchen, Germany (Carter *et al.* (2009b). This further confirms that the Toronto population is an introduction independent of any of those in the United States of America.

Patterns similar to the limited genetic diversity we report here for *A. glabripennis* have been recorded for other invasive species (Dlugosch and Parker 2008). For example, populations of Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae), introduced into Europe were found to have significantly reduced variation in nuclear markers (AFLPs) compared with source populations in the United States of America, and European populations are fixed for single mitochondrial DNA haplotypes (Grapputo *et al.* 2005). Six invasive populations of cabbage seedpod weevil, *Ceutorhynchus obstrictus* (Marsham) (Coleoptera: Curculionidae), in North America are each characterized by a single mitochondrial DNA haplotype, in contrast to native European populations, which have more haplotypes

(Laffin *et al.* 2005). Likewise, invasive populations of red turpentine beetle, *Dendroctonus valens* LeConte (Coleoptera: Curculionidae: Scolytinae), in China have fewer mitochondrial DNA haplotypes for the number of beetles sampled than do potential source populations in North America (Cai *et al.* 2008). Population-genetic analyses of mitochondrial DNA sequences, microsatellite markers, and a sex-determining locus showed that a solitary bee, *Lasioglossum leucozonium* (Schrank), experienced a single but severe bottleneck during its introduction from Europe into North America (Zayed *et al.* 2007).

An alternative explanation for the reduced genetic variation in the Toronto *A. glabripennis* population is that a larger founder event was followed by one or more genetic bottlenecks resulting from intense local selection or random survival and reproduction (Nei *et al.* 1975). It is probable that *A. glabripennis* first arrived at the Ontario site in 1996 or 1997 (J. Turgeon, unpublished data). Following Sawyer (2006), we dissected all infested trees to determine the year when exit holes were created. The analyzed beetles were collected between 2003 (the year of discovery) and 2005 (when we collected hundreds of eggs, larvae, and adults). In Toronto the shortest life cycle for *A. glabripennis* is usually 1 year, which suggests that there had been seven to nine generations prior to our assessment of genetic diversity. There are no biological data to suggest that the *A. glabripennis* population in Toronto is not mating at random. It appears that *A. glabripennis* in Toronto has followed the three main steps common to successful invasions: colonization, followed by a lag phase, and then rapid population expansion (Sakai *et al.* 2001).

Our inability to assign most beetles from Toronto to a single source population (province) in China is likely due to extensive genetic admixture within Chinese populations (any single multilocus genotype can be found in many provinces). This genetic admixture is due to reforestation efforts in China, where plantations of preferred hosts of *A. glabripennis* became infested with beetles and then beetles were moved to other areas in transported cut wood (Zhang *et al.* 2000; Uchida *et al.* 2005;

Yin and Lu 2005). This was evident when we divided the reference population into four clusters, with three clusters containing beetles from more than one province in China. Using these clusters as reference, we were able to assign 165 individuals to a single cluster.

Our data show that *A. glabripennis* can multiply to outbreak proportions in urban areas even when genetic diversity is extremely low. This suggests that the best protection against future invasions of this and similar forest pests will be regulatory actions that ensure the mortality of beetles transported in solid-wood packing materials from Asia.

### Acknowledgements

We thank A. Liang for technical assistance. R. Fournier created Figure 1. D. Borchert and W. Jinquan created Figure 2. P. de Groot and J. Pedlar provided valuable input during the study. We thank B. Gasman for making this research possible, many individuals from the Canadian Food Inspection Agency for collecting and preserving specimens, and two anonymous reviewers and the Editor for critically reviewing the manuscript and making helpful suggestions for revision. This work was supported by a gift from Howard P. Milstein and Leonard Litwin to Cornell University and by United States Department of Agriculture Specific Cooperative Agreement No. 58–1926–6–606 to Richard Harrison and Michael Smith.

### References

- Cai, Y.-W., Cheng, X.-Y., Xu, R.-M., Duan, D.-H., and Kirkendall, L.R. 2008. Genetic diversity and biogeography of red turpentine beetle *Dendroctonus valens* in its native and invasive regions. *Insect Science* **15**(4): 291–301 doi:10.1111/j.1744-7917.2008.00213.x.
- Canadian Food Inspection Agency. 2003. Toronto–Vaughan, Ontario \_ regulated area [online]. Canadian Food Inspection Agency. Available from <http://www.inspection.gc.ca/english/plaveg/pestrava/anogla/mc/mapcarrege.shtml> [accessed 4 February 2009].
- Canadian Food Inspection Agency. 2008. Asian longhorned beetle \_ latest information [online]. Canadian Food Inspection Agency. Available from <http://www.inspection.gc.ca/english/plaveg/pestrava/anogla/situationone.shtml> [accessed 20 October 2008].
- Carroll, S.P. 2007. Brave New World: the epistatic foundations of natives adapting to invaders. *Genetica*, **129**(2): 193–204 PMID:16924403 doi:10.1007/s10709-006-9014-8.
- Carter, M., Casa, A.M., Zeid, M., Mitchell, S.E., and Kresovich, S. 2009a. Isolation and characterization of microsatellite loci for the Asian longhorned beetle, *Anoplophora glabripennis*. *Molecular Ecology Resources*, **9**(3): 925–928 doi:10.1111/j.1755-0998.2008.02485.x.
- Carter, M., Smith, M., and Harrison, R. 2009b. Genetic analyses of the Asian longhorned beetle (Coleoptera, Cerambycidae, *Anoplophora glabripennis*), in North America, Europe and Asia. *Biological Invasions*. In press. doi:10.1007/s10530-009-9538-9.
- Carter, M.E., Smith, M.T., and Harrison, R.G. 2009c. Patterns of genetic variation among populations of the Asian longhorned beetle (Coleoptera: Cerambycidae) in China and Korea. *Annals of the Entomological Society of America*, **102**(5): 985–905.
- Cavey, J.F., Hoebeke, E.R., Passoa, S., and Lingafelter, S.W. 1998. A new exotic threat to North American hardwood forests: an Asian longhorned beetle, *Anoplophora glabripennis* (Motschulsky) (Coleoptera: Cerambycidae). I. Larval description and diagnosis. *Proceedings of the Entomological Society of Washington*, **100**: 373–381.
- Cheverud, J.M., Vaughn, T.T., Pletscher, L.S., King-Ellison, K., Bailiff, J., Adams, E., *et al.* 1999. Epistasis and the evolution of additive genetic variance in populations that pass through a bottleneck. *Evolution*, **53**(4): 1009–1018 doi:10.2307/2640806.
- Colautti, R.I., Bailey, S.A., van Overdijk, C.D.A., Amundsen, K., and MacIsaac, H.J. 2006. Characterised and projected costs of nonindigenous species in Canada. *Biological Invasions*, **8**(1): 45–59 doi:10.1007/s10530-005-0236-y.
- Dlugosch, K.M., and Parker, I.M. 2008. Founding events in species invasions: genetic variation, adaptive evolution, and the role of multiple introductions. *Molecular Ecology*, **17**(1): 431–449 PMID:17908213 doi:10.1111/j.1365-294X.2007.03538.x.
- Eales, J., Thorpe, R.S., and Malhotra, A. 2008. Weak founder effect signal in a recent introduction of Caribbean *Anolis*. *Molecular Ecology*, **17**(6): 1416–1426 PMID:18266633 doi:10.1111/j.1365-294X.2007.03684.x.
- Falush, D., Stephens, M., and Pritchard, J.K. 2003. Inference of population structure using multi-locus genotype data: linked loci and correlated allele frequencies. *Genetics*, **164**(4): 1567–1587 PMID:12930761.
- Ficetola, G.F., Bonin, A., and Miaud, C. 2008. Population genetics reveals origin and number of founders in a biological invasion. *Molecular Ecology*, **17**(3): 773–782 PMID:18194168.

- Goudet, J. 1995. FSTAT (version 1.2): a computer program to calculate *F*-statistics. *Journal of Heredity*, **86**: 485–486.
- Grapputo, A., Boman, S., Lindström, L., Lyytinen, A., and Mappes, J. 2005. The voyage of an invasive species across continents: genetic diversity of North American and European Colorado potato beetle populations. *Molecular Ecology*, **14**(14): 4207–4219 PMID:16313587.
- Haack, R.A., Law, K.R., Mastro, V.C., Ossenbruggen, H.S., and Raimo, B.J. 1997. New York's battle with the Asian longhorned beetle. *Journal of Forestry*, **95**(1): 11–15.
- Hajek, A.E., and Kalb, D.M. 2007. Suitability of *Acer saccharum* and *Acer pensylvanicum* (Aceraceae) for rearing *Anoplophora glabripennis* (Coleoptera: Cerambycidae). *The Canadian Entomologist*, **139**(5): 751–755 doi:10.4039/N06-107.
- Hérard, F., Ciampitti, M., Maspero, M., Krehan, H., Benker, U., Boegel, C., *et al.* 2006. *Anoplophora* species in Europe: infestations and management processes. EPPO (European and Mediterranean Plant Protection Organization) Bulletin, **36**: 470–474.
- Hunter, W.B., Smith, M.T., and Hunnicutt, L.E. 2009. Analysis and functional annotation of expressed sequence tags from the Asian longhorned beetle, *Anoplophora glabripennis*. *Journal of Insect Science*, **9**(21): 1–13 doi:10.1673/031.009.2101.
- Jarne, P., and Lagoda, P.J.L. 1996. Microsatellites, from molecules to populations and back. *Trends in Ecology and Evolution*, **11**(10): 424–429 doi:10.1016/0169-5347(96)10049-5.
- Keena, M.A. 2002. *Anoplophora glabripennis* (Coleoptera: Cerambycidae) fecundity and longevity under laboratory conditions: comparisons of populations from New York and Illinois on *Acer saccharum*. *Environmental Entomology*, **31**: 490–498.
- Keena, M.A. 2005. Pourable artificial diet for rearing *Anoplophora glabripennis* (Coleoptera: Cerambycidae) and methods to optimize larval survival and synchronize development. *Annals of the Entomological Society of America*, **98**(4): 536–547 doi:10.1603/0013-8746(2005)098[0536:PADFRA]2.0.CO;2.
- Laffin, R.D., Dosdall, L.M., and Sperling, F.A.H. 2005. Population structure of the cabbage seedpod weevil, *Ceutorhynchus obstrictus* (Marsham) (Coleoptera Curculionidae): origins of North American introductions. *Environmental Entomology*, **34**(2): 504–510.
- Lingafelter, S.W., and Hoebeke, E.R. 2002. Revision of *Anoplophora* (Coleoptera: Cerambycidae) Entomological Society of Washington, Washington, D.C.
- Liu, K., and Muse, S.V. 2005. PowerMarker: an integrated analysis environment for genetic marker analysis. *Bioinformatics*, **21**(9): 2128–2129 PMID:15705655 doi:10.1093/bioinformatics/bti282.
- Maspero, M., Jucker, C., and Colombo, M. 2007. First record of *Anoplophora glabripennis* (Motschulsky) (Coleoptera Cerambycidae Lamiinae Lamiini) in Italy. *Bollettino di Zoologia Agraria e di Bachicoltura*, **39**(2): 161–164.
- Moore, R., and Russell, R. 1990. The “Three North” forest protection system — China. *Agroforestry Systems*, **10**(1): 71–88 doi:10.1007/BF00118728.
- Morewood, W.D., Neiner, P.R., McNeil, J.R., Sellmer, J.C., and Hoover, K. 2003. Oviposition preference and larval performance of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) in four eastern North American hardwood tree species. *Environmental Entomology*, **32**(5): 1028–1034.
- Morewood, W.D., Neiner, P.R., Sellmer, J.C., and Hoover, K. 2004. Behavior of adult *Anoplophora glabripennis* on different tree species under greenhouse conditions. *Journal of Insect Behavior*, **17**(2): 215–226 doi:10.1023/B:JOIR.0000028571.52739.50.
- Naciri-Graven, Y., and Goudet, J. 2008. The additive genetic variance after bottlenecks is affected by the number of loci involved in epistatic interactions. *Evolution*, **7**: 706–716.
- Nei, M., Maruyama, T., and Chakraborty, R. 1975. The bottleneck effect and genetic variability in populations. *Evolution*, **29**(1): 1–10 doi:10.2307/2407137.
- North American Plant Protection Organization. 2007. Update on the Asian longhorned beetle (*Anoplophora glabripennis*) — Canada. Phytosanitary Alert Official Pest Report 10/12/2007 [online]. Available from <http://www.pestalert.org/oprDetail.cfm?oprID=288> [accessed 19 October 2008].
- Nowak, D.J., Pasek, J.E., Sequeira, R.A., Crane, D.E., and Mastro, V.C. 2001. Potential effect of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) on urban trees in the United States. *Journal of Economic Entomology*, **94**(1): 116–122 PMID:11233100.
- Oliveira, E.J., Padua, J.G., Zucchi, M.I., Vencovsky, R., and Vieira, M.L.C. 2006. Origin, evolution and genome distribution of microsatellites. *Genetics and Molecular Biology*, **29**(2): 294–307 doi:10.1590/S1415-47572006000200018.
- Pan, H.Y. 2005. Review of the Asian longhorned beetle: research, biology, distribution and management in China [online]. Working Paper FBS/6E, Forestry Department, Food and Agriculture Organization, Rome, Italy. Available from <http://www.fao.org/forestry/media/66891/1/102/> [accessed 27 May 2008].
- Park, S.D.E. 2001. Trypanotolerance in West African cattle and the population genetic effects of selection [online]. Ph.D. thesis, University of Dublin, Dublin, Eire. Available from

- <http://animalgenomics.ucd.ie/sdepark/ms-toolkit/> [accessed 12 June 2009].
- Piry, S., Alapetite, A., Cornuet, J.-M., Paetkau, D., Baudouin, L., and Estoup, A. 2004. GENECLASS2: a software for genetic assignment and first-generation migrant detection. *Journal of Heredity*, **95**(6): 536–539 PMID:15475402 doi:10.1093/jhered/esh074.
- Poland, T.M., Haack, R.A., and Petrice, T.R. 1998. Chicago joins New York in battle with the Asian longhorned beetle. *Newsletter of the Michigan Entomological Society*, **43**: 15–17.
- Rannala, B., and Mountain, J.L. 1997. Detecting immigration by using multilocus genotypes. *Proceedings of the National Academy of Sciences of the United States of America*, **94**(17): 9197–9201 PMID:9256459 doi:10.1073/pnas.94.17.9197.
- Raymond, M., and Rousset, F. 1995. Genepop (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity*, **86**(3): 248–249.
- Ric, J., de Groot, P., Gasman, B., Orr, M., Doyle, J., Smith, M.T., *et al.* 2007. Detecting signs and symptoms of Asian longhorned beetle injury: a training guide [online]. Canadian Forest Service, Sault Ste. Marie, Ontario. Available from <http://warehouse.pfc.forestry.ca/glfc/26860.pdf> [accessed 19 November 2008].
- Roderick, G.K., and Navajas, M. 2003. Genes in new environments: genetics and evolution in biological control. *Nature Reviews Genetics*, **4**(11): 889–899 PMID:14634636 doi:10.1038/nrg1201.
- Sabol, O. 2006. Faunistic records from the Czech Republic — 196. *Klapalekiana*, **42**: 78.
- Sakai, A.K., Allendorf, F.W., Holt, J.S., Lodge, D.M., Molofsky, J., With, K.A., *et al.* 2001. The population biology of invasive species. *Annual Review of Ecology and Systematics*, **32**(1): 305–332 doi:10.1146/annurev.ecolsys.32.081501.114037.
- Sawyer, A.J. 2003. Annotated categorization of ALB host trees [online]. United States Department of Agriculture, Animal and Plant Health Protection Service, Plant Protection and Quarantine, Otis Plant Protection Laboratory, Otis Air National Guard Base, Otis, Massachusetts. Available from <http://www.uvm.edu/albeetle/hosts.htm> [accessed 4 October 2008].
- Sawyer, A.J. 2006. A controlled study of the healing response of host trees to simulated Asian longhorned beetle damage. *In Emerald Ash Borer and Asian Longhorned Beetle Research and Development Review Meeting*, Cincinnati, Ohio, 29 October – 2 November 2006. *Compiled by V. Mastro, D. Lance, R. Reardon, and G. Parra. FHTET-2007-04*, United States Department of Agriculture, Animal and Plant Health Protection Service, Plant Protection and Quarantine, Otis Air National Guard Base, Otis, Massachusetts pp. 126–128.
- Schlötterer, C. 2004. The evolution of molecular markers — just a matter of fashion? *Nat. Rev. Genet.* **5**(1): 63–69 PMID:14666112 doi:10.1038/nrg1249.
- Sellers, C. 2004. The Asian longhorned beetle in Ontario. *Ontario Insects*, **9**: 21.
- Smith, M.T., Bancroft, J., and Tropp, J. 2002. Age-specific fecundity of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) on three tree species infested in the United States. *Environmental Entomology*, **31**(1): 76–83.
- Smith, M.T., Tobin, P., Bancroft, J.S., Li, G., and Gao, R. 2004. Dispersal and spatiotemporal dynamics of Asian longhorned beetle (Coleoptera: Cerambycidae) in China. *Environmental Entomology*, **33**(2): 435–442.
- Tomiczek, C., Krehan, H., and Menschhorn, P. 2002. Dangerous Asiatic longicorn beetle found in Austria: new danger for our trees?. *AFZ/der Wald: allgemeine Forst Zeitschrift für Waldwirtschaft und Umweltversorge*, **57**(2): 52–54.
- Turgeon, J.J., Ric, J., De Groot, P., Gasman, B., Orr, M., Doyle, J., *et al.* 2007. Détection des signes et des symptômes d'attaque par le longicorne étoilé : guide de formation [online]. Agence canadienne d'inspection des aliments, Sault Ste. Marie, Ontario. Available from <http://warehouse.pfc.forestry.ca/glfc/26861.pdf> [accessed 16 June 2009].
- Uchida, E., Xu, J., and Rozelle, S. 2005. Grain for green: cost-effectiveness and sustainability of China's conservation set-aside program. *Land Economics* **81**(2): 247–264.
- Wade, M.J., Shuster, S.M., and Stevens, L. 1996. Inbreeding: its effect on response to selection for pupal weight and the heritable variance in fitness in the flour beetle, *Tribolium castaneum*. *Evolution*, **50**(2): 723–733 doi:10.2307/2410845.
- Wang, Z. 2004. Occurrence of *Anoplophora glabripennis* (Motsch.) and control in China. Ph.D. thesis, Northeast Forestry University of China, Harbin, Heilongjiang Province, People's Republic of China.
- Wares, J.P., Hughes, A.R., and Grosberg, R.K. 2005. Mechanisms that drive evolutionary change. *In Species invasions: insights into ecology, evolution, and biogeography. Edited by D.F. Sax, J.J. Stachowicz, and S.D. Gaines.* Sinauer Associates Inc., Sunderland, Massachusetts. pp. 229–257.
- Weir, B.S., and Cockerham, C.C. 1984. Estimating F-statistics for the analysis of population structure. *Evolution*, **38**(6): 1358–1370 doi:10.2307/2408641.
- Williams, D.W., Lee, H.P., and Kim, I.K. 2004. Distribution and abundance of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) in natural *Acer* stands in South Korea. *Environmental Entomology*, **33**(3): 540–545.
- Xiao, G. 1992. Forest insects of China. China Forestry Publishing House, Beijing, People's Republic of China.
- Yin, W., and Lu, W. 2005. Review of tree selection and afforestation for control of Asian longhorned beetle in north China [online]. Forestry Department Working Paper FBS/7E, Food and

- Agriculture Organization, Rome, Italy. Available from <http://www.fao.org/forestry/media/7175/102/> [accessed 27 May 2008].
- Zayed, A., Constantin, S.A., and Packer, L. 2007. Successful biological invasion despite a severe genetic load. *PLoS One*, **9**: 1–6.
- Zhang, P., Shao, G., Zhao, G., Le Master, D.C., Parker, G.R., Dunning, J.B., Jr., Li, Q. 2000. China's forest policy for the 21st century. *Science* (Washington, D.C.), **288**(5474): 2135–2136. PMID:10896587 doi:10.1126/science.288.5474.2135.