Table 28. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Family Income (as % of Poverty Level³) and Age, in the United States, 2015-2016

Family income as % of poverty level Percent and age reporting ⁴				Energy		otein	Carbo- hydrate		Total sugars		Dietary fiber		Total fat		Saturated fat		Mono- unsaturated fat		unsat	oly- urated at
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Under 131% poverty:																				
2 - 5	90	(2.3)	25	(1.8)	15	(1.5)	28	(1.8)	36	(2.1)	22	(1.8)	23	(2.3)	24	(2.5)	23	(2.2)	23	(2.4)
6 - 11	89	(2.5)	24	(1.6)	15	(1.3)	28	(1.7)	33	(2.1)	21	(1.0)	24	(1.8)	24	(1.9)	23	(1.8)	26	(2.4)
12 - 19	85	(2.1)	22	(1.6)	14	(1.2)	25	(1.7)	33	(2.0)	19	(1.4)	21	(2.0)	22	(2.4)	21	(2.0)	21	(2.2)
20 and over	89	(1.2)	22	(1.0)	12	(0.8)	27	(1.3)	38	(1.7)	17	(0.8)	19	(1.0)	20	(1.2)	19	(0.9)	19	(1.1)
2 and over	89	(0.9)	23	(0.7)	13	(0.6)	27	(1.0)	36	(1.3)	18	(0.6)	20	(0.6)	21	(0.9)	19	(0.6)	20	(0.6)
131-350% poverty:																				
2 - 5	96*	(1.5)	27	(1.5)	18	(1.4)	30	(1.5)	38	(2.1)	23	(1.7)	26	(1.7)	27	(1.6)	25	(2.0)	24	(1.9)
6 - 11	97*	(1.2)	24	(1.6)	12	(1.2)	28	(1.6)	33	(1.6)	22	(1.4)	23	(1.9)	23	(2.4)	22	(1.8)	25	(1.9)
12 - 19	87	(2.6)	23	(1.3)	12	(0.8)	26	(1.4)	33	(2.2)	21	(1.2)	23	(1.7)	24	(2.3)	23	(1.6)	24	(1.4)
20 and over	94	(0.7)	21	(0.6)	11	(0.4)	25	(0.6)	35	(1.1)	19	(0.8)	18	(0.7)	18	(0.8)	18	(0.9)	17	(0.7)
2 and over	93	(0.7)	22	(0.6)	12	(0.4)	26	(0.6)	35	(0.9)	20	(0.7)	19	(0.7)	20	(0.9)	19	(0.8)	18	(0.6)
Over 350% poverty:																				
2 - 5	97*	(2.4)	28	(1.8)	20	(1.5)	33	(2.1)	41	(3.4)	22	(2.2)	26	(2.2)	25	(2.6)	27	(2.7)	25	(3.2)
6 - 11	97*	(1.4)	26	(1.5)	15	(1.0)	30	(1.7)	39	(2.2)	26	(2.0)	26	(1.8)	27	(2.6)	25	(1.7)	26	(1.9)
12 - 19	93*	(1.8)	19	(1.9)	10	(0.9)	23	(2.4)	32	(3.3)	18	(1.6)	19	(1.8)	19	(2.2)	19	(1.5)	19	(1.7)
20 and over	97	(0.4)	22	(1.0)	11	(0.7)	25	(1.0)	35	(1.4)	18	(0.9)	19	(1.2)	19	(1.3)	20	(1.3)	18	(1.1)
2 and over	96	(0.4)	22	(0.9)	12	(0.7)	25	(1.0)	35	(1.4)	18	(0.9)	19	(1.1)	20	(1.1)	20	(1.2)	18	(1.0)
All Individuals ⁵ :																				
2 - 5	95	(1.2)	27	(0.8)	18	(0.8)	30	(0.9)	39	(1.2)	23	(0.9)	25	(1.1)	26	(1.1)	25	(1.2)	25	(1.4)
6 - 11	94	(1.0)	25	(0.9)	14	(0.7)	29	(0.9)	35	(1.1)	23	(0.9)	24	(0.9)	24	(1.1)	23	(0.8)	26	(1.0)
12 - 19	88	(1.6)	21	(1.0)	12	(0.6)	24	(1.2)	33	(1.5)	20	(0.7)	21	(1.1)	22	(1.4)	21	(1.1)	21	(1.2)
20 and over	94	(0.4)	22	(0.7)	12	(0.5)	25	(0.7)	36	(0.9)	18	(0.5)	19	(0.7)	19	(0.8)	19	(0.8)	18	(0.7)
2 and over	93	(0.5)	22	(0.6)	12	(0.4)	26	(0.6)	35	(0.8)	19	(0.4)	19	(0.6)	20	(0.7)	20	(0.7)	19	(0.6)

Table 28. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Family Income (as % of Poverty Level³) and Age, in the United States, 2015-2016 *(continued)*

Family income as % of poverty level and age (years)	Choles- terol % (SE)		terol (RAE)		•		Lyco %	Lycopene % (SE)		Thiamin % (SE)		Ribo- flavin % (SE)		Niacin % (SE)		Vitamin B6 % (SE)		olate FE) (SE)
Under 131% poverty:																		
2 - 5	12	(1.4)	19	(2.0)	24*	(8.9)	9*	(3.1)	17	(2.0)	20	(1.6)	14	(1.8)	15	(1.7)	16	(2.3)
6 - 11	12	(1.5)	19	(1.9)	20	(4.2)	13	(2.3)	19	(1.6)	19	(1.5)	17	(1.2)	17	(1.3)	21	(2.2)
12 - 19	11	(1.6)	21	(4.3)	9	(1.8)	8	(2.2)	17	(2.1)	20	(2.3)	15	(1.7)	18	(2.1)	18	(2.6)
20 and over	9	(0.8)	16	(1.2)	11	(1.3)	9	(1.2)	15	(0.8)	21	(1.1)	17	(1.1)	19	(1.5)	15	(0.9)
2 and over	9	(0.6)	17	(1.3)	12	(1.1)	9	(1.2)	16	(0.7)	21	(0.7)	16	(0.7)	18	(1.1)	16	(0.8)
131-350% poverty:																		
2 - 5	13	(1.1)	22	(1.9)	19	(3.2)	6*	(2.2)	18	(1.0)	24	(1.6)	15	(1.4)	19	(1.5)	17	(1.3)
6 - 11	9	(1.3)	20	(5.5)	30*	(11.2)	6*	(2.8)	16	(1.5)	18	(2.2)	14	(1.6)	15	(2.3)	15	(1.2)
12 - 19	11	(1.4)	14	(1.3)	6	(1.0)	8	(1.3)	15	(1.0)	18	(1.3)	14	(1.0)	16	(2.2)	15	(1.5)
20 and over	7	(0.4)	20	(2.6)	20*	(7.5)	7	(1.2)	14	(0.6)	21	(0.8)	17	(0.7)	20	(0.9)	16	(1.0)
2 and over	8	(0.5)	19	(2.0)	20*	(6.3)	7	(1.0)	15	(0.5)	21	(0.7)	16	(0.5)	19	(0.8)	16	(0.8)
Over 350% poverty:																		
2 - 5	16*	(2.1)	20	(2.8)	6*	(1.9)	10*	(4.6)	21	(0.8)	27	(2.3)	15*	(1.1)	22	(2.1)	22	(1.8)
6 - 11	11	(1.8)	15	(1.7)	11*	(3.5)	7*	(3.3)	16	(1.4)	18	(1.8)	15	(1.1)	16	(1.2)	16	(1.7)
12 - 19	7*	(0.9)	17	(2.9)	17*	(5.1)	9*	(4.3)	12	(1.1)	17	(1.5)	12	(1.2)	16	(2.2)	12	(1.7)
20 and over	8	(0.9)	15	(0.9)	7	(1.4)	6	(0.7)	13	(0.6)	21	(1.0)	15	(1.1)	17	(1.6)	14	(0.5)
2 and over	8	(0.8)	15	(0.8)	8	(1.4)	6	(0.7)	14	(0.5)	20	(0.9)	15	(0.9)	17	(1.3)	14	(0.5)
All Individuals ⁵ :																		
2 - 5	13	(0.7)	20	(1.1)	13	(3.0)	8	(1.4)	19	(0.9)	23	(1.0)	15	(0.9)	18	(1.0)	17	(1.0)
6 - 11	10	(0.7)	18	(2.5)	20*	(6.1)	10	(1.3)	17	(1.0)	19	(1.1)	15	(0.9)	16	(1.2)	17	(1.1)
12 - 19	10	(0.9)	16	(1.6)	11	(2.0)	8	(1.2)	14	(0.9)	18	(0.9)	13	(0.7)	16	(0.7)	14	(1.4)
20 and over	8	(0.5)	17	(1.0)	12	(3.0)	7	(0.7)	14	(0.4)	21	(0.6)	16	(0.7)	18	(0.7)	15	(0.5)
2 and over	8	(0.4)	17	(0.8)	13	(2.5)	8	(0.5)	15	(0.3)	21	(0.5)	16	(0.5)	18	(0.5)	15	(0.4)

Table 28. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Family Income (as % of Poverty Level³) and Age, in the United States, 2015-2016 *(continued)*

Family income as % of poverty level										min E pha-								
and age	Ch	oline	Vitan	nin B12	Vita	min C	Vita	min D	tocop	herol)	Vita	Vitamin K		Calcium		Phosphorus		nesium
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Under 131% poverty:																		
2 - 5	16	(1.3)	16	(1.6)	37	(4.0)	21	(1.9)	26	(2.5)	15	(2.3)	26	(1.6)	20	(1.6)	23	(1.4)
6 - 11	14	(1.1)	15	(1.6)	24	(2.5)	17	(2.0)	28	(1.9)	16	(1.5)	20	(2.1)	17	(1.4)	22	(1.4)
12 - 19	15	(1.5)	17	(2.5)	30	(2.7)	22	(4.3)	24	(3.4)	16	(2.1)	23	(2.2)	17	(1.8)	25	(2.2)
20 and over	13	(0.6)	16	(1.7)	22	(2.0)	16	(1.5)	22	(1.1)	9	(0.8)	25	(1.5)	17	(0.9)	25	(1.1)
2 and over	14	(0.4)	16	(1.2)	24	(1.5)	18	(1.2)	23	(0.9)	11	(0.6)	24	(1.0)	17	(0.6)	24	(0.9)
131-350% poverty:																		
2 - 5	18	(1.5)	22	(2.0)	36	(3.9)	26	(2.7)	27	(2.8)	20	(2.3)	29	(1.6)	23	(1.7)	25	(2.0)
6 - 11	13	(1.0)	11	(1.7)	25	(2.8)	13	(1.4)	27	(1.4)	19	(2.3)	20	(2.8)	17	(2.1)	22	(1.2)
12 - 19	15	(1.1)	15	(2.0)	24	(1.9)	14	(2.5)	25	(1.7)	14	(1.5)	19	(1.2)	16	(0.9)	24	(1.0)
20 and over	14	(0.6)	15	(0.8)	26	(1.6)	16	(1.4)	23	(1.3)	12	(1.7)	23	(1.0)	16	(0.5)	25	(0.6)
2 and over	14	(0.6)	15	(0.7)	26	(1.3)	16	(1.0)	24	(1.1)	12	(1.3)	22	(0.9)	17	(0.5)	25	(0.6)
Over 350% poverty:																		
2 - 5	21	(2.2)	25	(2.9)	31	(5.9)	32	(4.4)	26	(2.6)	18	(3.1)	30	(3.1)	25	(2.0)	28	(2.0)
6 - 11	15	(1.1)	17	(1.4)	31	(3.7)	16	(1.9)	27	(2.1)	16	(1.1)	20	(2.2)	18	(1.3)	26	(2.0)
12 - 19	11	(1.2)	14	(1.2)	29	(3.5)	15	(3.3)	24	(2.6)	12	(1.9)	19	(2.3)	14	(1.5)	22	(1.4)
20 and over	13	(0.9)	14	(1.7)	19	(2.0)	12	(1.3)	23	(1.5)	6	(0.5)	22	(1.0)	16	(0.7)	25	(0.9)
2 and over	13	(0.8)	15	(1.5)	21	(1.8)	13	(1.0)	23	(1.4)	7	(0.5)	22	(0.8)	16	(0.7)	25	(0.8)
All Individuals ⁵ :																		
2 - 5	18	(0.9)	20	(0.9)	36	(2.9)	25	(1.5)	27	(1.6)	18	(1.6)	28	(1.1)	22	(1.0)	26	(1.1)
6 - 11	14	(0.5)	14	(0.8)	27	(2.0)	15	(1.1)	28	(0.8)	17	(1.2)	20	(1.3)	18	(1.0)	24	(1.0)
12 - 19	13	(0.7)	15	(0.8)	27	(1.6)	16	(1.7)	24	(1.2)	13	(1.0)	20	(1.0)	16	(0.9)	24	(0.6)
20 and over	13	(0.6)	15	(0.9)	22	(0.9)	14	(1.0)	23	(0.9)	9	(0.6)	23	(0.7)	16	(0.5)	25	(0.5)
2 and over	14	(0.5)	15	(0.7)	24	(0.8)	15	(0.7)	23	(0.7)	10	(0.6)	23	(0.5)	17	(0.4)	25	(0.5)

Table 28. Snacks¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Family Income (as % of Poverty Level³) and Age, in the United States, 2015-2016 *(continued)*

Family income as % of poverty level																
and age	I	ron	Z	inc	Co	pper	Sele	nium	Pota	ssium	So	dium	Caf	feine	Alo	cohol
(years)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Under 131% poverty:																
2 - 5	18	(2.4)	16	(1.2)	25	(1.9)	13	(1.5)	23	(1.5)	17	(1.8)	37	(8.9)		
6 - 11	23	(2.1)	15	(1.4)	25	(1.5)	13	(1.2)	20	(1.0)	18	(1.7)	42	(8.7)		
12 - 19	20	(2.2)	16	(1.9)	26	(2.8)	12	(1.4)	20	(1.3)	16	(1.5)	33	(6.8)		
20 and over	15	(1.1)	14	(0.9)	25	(1.1)	10	(0.7)	20	(0.9)	13	(0.7)	41	(2.6)	60	(5.2)
2 and over	17	(0.9)	15	(0.7)	25	(0.8)	11	(0.5)	20	(0.6)	14	(0.5)	40	(2.5)		
131-350% poverty:																
2 - 5	18	(1.6)	19	(1.5)	28	(2.6)	15	(1.4)	26	(1.8)	18	(1.1)	27	(6.5)		
6 - 11	17	(1.4)	13	(1.2)	24	(1.0)	11	(1.1)	19	(1.2)	17	(1.6)	36	(2.8)		
12 - 19	15	(1.4)	15	(0.9)	27	(1.5)	11	(0.8)	19	(0.9)	13	(0.7)	41	(3.8)		
20 and over	15	(1.0)	14	(0.7)	26	(0.7)	9	(0.3)	19	(0.5)	11	(0.4)	35	(2.4)	63	(5.8)
2 and over	16	(0.9)	14	(0.6)	26	(0.7)	10	(0.3)	19	(0.5)	12	(0.4)	35	(2.4)		
Over 350% poverty:																
2 - 5	22	(2.3)	22	(2.2)	30	(1.3)	16*	(2.0)	28	(3.0)	18	(1.4)	41*	(6.5)		
6 - 11	18	(1.3)	18	(1.2)	30	(2.1)	12	(1.3)	23	(1.5)	16	(1.0)	57	(6.9)		
12 - 19	13	(2.1)	12	(1.1)	24	(1.7)	8*	(1.0)	18	(1.6)	10	(1.2)	44	(6.7)		
20 and over	15	(0.5)	14	(0.7)	28	(1.1)	10	(0.7)	18	(0.8)	11	(0.6)	38	(2.1)	62	(3.7)
2 and over	15	(0.6)	15	(0.6)	27	(1.0)	10	(0.7)	18	(0.8)	11	(0.5)	39	(2.1)		
All Individuals ⁵ :																
2 - 5	19	(1.3)	19	(0.9)	28	(1.2)	14	(0.9)	25	(1.2)	18	(0.9)	33	(3.8)		
6 - 11	20	(1.2)	15	(0.8)	26	(1.0)	12	(0.8)	21	(0.8)	17	(0.8)	42	(3.9)		
12 - 19	16	(1.2)	14	(0.9)	26	(0.7)	10	(0.7)	19	(0.7)	13	(0.8)	39	(3.0)		
20 and over	15	(0.5)	14	(0.5)	27	(0.6)	10	(0.4)	19	(0.6)	12	(0.4)	38	(1.6)	61	(3.5)
2 and over	16	(0.5)	14	(0.4)	27	(0.5)	10	(0.4)	19	(0.5)	12	(0.3)	38	(1.6)		

Symbol Legend

* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages and ratios are as follows:

Percent reporting: An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 2.41.

Nutrient ratios expressed as percentages: An estimated ratio between 25 and 75 percent is flagged when based on a sample size n^* of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect and n^* is the number of individuals in the sample reporting non-zero intake of the respective nutrient. An estimated ratio less than or equal to 25 percent or greater than or equal to 75 percent, is flagged when the smaller of n^*p and n^* (1-p) is less than 8 times the VIF, where p is the percentage expressed as a fraction. Additionally, an estimated ratio is flagged when either the relative standard error or p/(1-p) times the relative standard error is greater than 30 percent. The VIF used in this table is 2.41.

Footnotes

- ¹ Snack occasions were reported as distinct eating occasions during the dietary interview and consisted of one or more food and beverage items, including plain water. Water was the only item reported in approximately 27 percent of the snack occasions. Survey respondents selected the name of all eating occasions from a fixed list that was provided during the interview. All reports of "snack", "drink" or "extended consumption" (items that were consumed over a long period of time) were included as snack occasions. Spanish language interviewers used Spanish language snack occasion names: "merienda", "entre comida", "botana", "botana", "tentempie", and "bebida".
- ² Percentages are estimated as a ratio of total nutrients from snacks for all individuals to total daily nutrient intakes for all individuals. Total daily nutrient intakes are available from: www.ars.usda.gov/nea/bhnrc/fsrg. See Table 4. Nutrient Intakes from Food and Beverages: Mean Amounts Consumed per Individual, by Family Income (as % of Poverty Level) and Age, in the United States, 2015-2016.
- ³ The ratio of family income to poverty expressed as a percentage. The Department of Health and Human Services' poverty guidelines were used as the poverty measure to calculate the ratio, http://aspe.hhs.gov/poverty/.
- ⁴ The percentage of respondents in the income/age group who reported consuming at least one item at an eating occasion designated as snack.
- ⁵ Includes persons of all income levels or with unknown family income.

Abbreviations

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2015-2016

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2015-2016. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2015-2016 www.ars.usda.gov/nea/bhnrc/fsrg.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

Suggested Citation

U.S. Department of Agriculture, Agricultural Research Service. 2018. Snacks: Percentages of Selected Nutrients Contributed by Food and Beverages Consumed at Snack Occasions, by Family Income (as % of Poverty Level) and Age, What We Eat in America, NHANES 2015-2016.