Table 29. Snacks: Distribution of Snack Occasions¹, by Gender and Age, in the United States, 2017-March 2020 Prepandemic

Gender and age (years)																	
	Sample size	Zero		One		Two		Three		Four		Five		Six		Seven or more	
		%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Males:	1																
2 - 5	468	5	(1.0)	9	(1.5)	17	(1.7)	24	(2.2)	24	(2.3)	11	(2.2)	5	(1.2)	4*	` /
6 - 11	701	4	(0.7)	18	(2.3)	30	(2.4)	23	(2.9)	15	(2.3)	6	(1.6)	2*	(0.9)	1*	()
12 - 19	899	10	(1.2)	24	(2.3)	28	(1.9)	18	(2.0)	13	(1.7)	5	(1.1)	1*	(0.4)	1*	(0.4)
20 - 29	551	8	(1.6)	18	(2.2)	25	(2.2)	24	(3.4)	11	(2.6)	7	(2.1)	5	(2.1)	2*	(1.1)
30 - 39	535	7	(1.2)	16	(1.8)	25	(2.6)	22	(1.9)	13	(2.5)	6	(1.2)	6	(1.5)	5	(0.7)
40 - 49	613	6	(1.1)	15	(2.3)	21	(2.1)	21	(2.7)	15	(2.4)	11	(2.7)	7	(2.1)	3*	(2.1)
50 - 59	608	4	(1.1)	19	(2.8)	23	(2.7)	24	(2.6)	16	(2.5)	7	(1.4)	3*	(1.0)	3*	(1.0)
60 - 69	776	4	(0.7)	17	(2.2)	29	(3.6)	22	(2.9)	13	(3.7)	7	(1.6)	4	(1.7)	4	(1.1)
70 and over	662	7	(1.3)	21	(2.0)	26	(2.6)	17	(2.7)	14	(2.5)	9	(1.7)	2*	(0.8)	4	(1.3)
2 - 19	2068	7	(0.7)	19	(1.5)	26	(1.4)	21	(1.4)	16	(1.1)	7	(0.8)	2	(0.5)	2	(0.3)
20 and over	3745	6	(0.7)	18	(0.9)	25	(1.1)	22	(0.7)	14	(1.0)	8	(0.9)	5	(0.6)	3	(0.5)
2 and over	5813	6	(0.6)	18	(0.8)	25	(0.9)	22	(0.7)	14	(0.8)	8	(0.7)	4	(0.5)	3	(0.4)
Females:																	
2 - 5	472	4*	(1.2)	8	(1.5)	22	(2.1)	26	(2.9)	21	(3.0)	9	(1.5)	7	(2.5)	3*	(1.1)
6 - 11	710	5	(1.0)	17	(2.3)	25	(2.7)	25	(2.0)	15	(2.1)	8	(1.2)	2*	(0.6)	2*	(0.8)
12 - 19	841	10	(1.9)	22	(1.7)	28	(2.4)	20	(1.5)	11	(1.3)	5	(1.1)	3	(1.4)	#	
20 - 29	605	8	(1.5)	17	(1.6)	23	(2.3)	25	(2.5)	12	(1.2)	9	(1.4)	4	(1.4)	2*	(0.9)
30 - 39	667	5	(1.1)	17	(2.1)	19	(2.3)	27	(2.7)	19	(3.5)	7	(1.0)	3*	(0.8)	2*	
40 - 49	621	4	(1.0)	15	(2.5)	23	(2.7)	25	(3.9)	18	(3.0)	7	(1.5)	4	(0.9)	4	(1.6)
50 - 59	706	4	(0.8)	14	(1.4)	26	(3.9)	23	(2.0)	14	(1.6)	12	(2.7)	4	(1.3)	4	(1.2)
60 - 69	744	4	(1.0)	15	(3.1)	21	(2.1)	22	(2.9)	18	(2.3)	9	(2.4)	6	(1.6)	4	(1.0)
70 and over	619	5	(1.0)	15	(1.7)	26	(2.6)	23	(2.0)	16	(1.7)	9	(1.8)	3*	(0.7)	2*	(0.9)
2 - 19	2023	7	(1.0)	17	(1.0)	26	(1.4)	23	(1.3)	15	(1.2)	7	(0.7)	4	(0.9)	2	(0.4)
20 and over	3962	5	(0.4)	16	(0.9)	23	(1.0)	24	(1.2)	16	(0.9)	9	(0.9)	4	(0.5)	3	(0.4)
2 and over	5985	6	(0.5)	16	(0.8)	24	(0.9)	24	(1.0)	16	(0.8)	9	(0.7)	4	(0.5)	3	(0.4)
Males and females:																	
2 - 19	4091	7	(0.7)	18	(1.1)	26	(0.9)	22	(1.1)	15	(0.9)	7	(0.6)	3	(0.5)	2	(0.3)
20 and over	7707	5	(0.7) (0.4)	17	(0.7)	24	(0.8)	23	(0.8)	15	(0.7)	8	(0.0)	4	(0.3) (0.4)	3	(0.3)
2 and over	11798	6	(0.4)	17	(0.7)	24	(0.7)	23	(0.7)	15	(0.6)	8	(0.6)	4	(0.4)	3	(0.2)

Symbol Legend

* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages are as follows:

Percent reporting: An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 2.76.

Indicates a non-zero value too small to report.

Footnotes

¹ Snack occasions were reported as distinct eating occasions during the dietary interview and consisted of one or more food and beverage items, including plain water. Water was the only item reported in approximately 27 percent of the snack occasions. Survey respondents selected the name of all eating occasions from a fixed list that was provided during the interview. All reports of "snack", "drink" or "extended consumption" (items that were consumed over a long period of time) were included as snack occasions. Spanish language interviewers used Spanish language snack occasion names: "merienda", "entre comida", "botana", "botana", "tentempie", and "bebida".

Abbreviations

SE = standard error.

Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2017-March 2020 Prepandemic

The 2017–March 2020 files represent a unique public-use data release from NHANES. The coronavirus disease 2019 (COVID-19) pandemic required suspension of data collection in March 2020. As a result, the partially completed NHANES 2019–2020 cycle was not nationally representative. Therefore, the 2019–March 2020 data were combined with the data from the 2017–2018 cycle to create the nationally representative 2017–March 2020 prepandemic data files.

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2017-March 2020 Prepandemic. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2017-2018 and 2019-2020 as applicable www.ars.usda.gov/nea/bhnrc/fsrg.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

Suggested Citation

U.S. Department of Agriculture, Agricultural Research Service. 2022. Snacks: Distribution of Snack Occasions, by Gender and Age, What We Eat in America, NHANES 2017-March 2020 Prepandemic.