Table 5. Energy Intakes: Percentages¹ of Energy from Protein, Carbohydrate, Fat, and Alcohol, by Male/Female and Age, in the United States, 2017-March 2020 Prepandemic

Males/Females and age	Sample size ²	Energy		Protein		Carbohydrate		Total fat		Saturated fat		Mono- unsaturated fat		Poly- unsaturated fat		Alcohol	
(years)		kcal	(SE)	% kcal	(SE)	% kcal	(SE)	% kcal	(SE)	% kcal	(SE)	% kcal	(SE)	% kcal	(SE)	% kcal	(SE)
Males:																	
2 - 5	468	1667	(41.7)	14	(0.2)	53	(0.7)	34	(0.5)	12	(0.3)	11	(0.2)	8	(0.2)		
6 - 11	701	1989	(38.4)	14	(0.3)	52	(0.6)	35	(0.5)	13	(0.3)	12	(0.2)	8	(0.2)		
12 - 19	899	2299	(36.4)	15	(0.2)	49	(0.6)	36	(0.4)	12	(0.2)	12	(0.2)	8	(0.2)		
20 - 29	551	2420	(80.9)	16	(0.3)	46	(0.7)	35	(0.6)	11	(0.2)	12	(0.2)	8	(0.4)		
30 - 39	535	2688	(70.3)	16	(0.3)	46	(0.7)	35	(0.6)	11	(0.3)	12	(0.3)	8	(0.2)		
40 - 49	613	2579	(86.3)	16	(0.4)	44	(0.7)	37	(0.6)	12	(0.2)	13	(0.4)	8	(0.3)		
50 - 59	608	2545	(42.8)	15	(0.4)	45	(0.9)	36	(0.5)	11	(0.2)	12	(0.3)	8	(0.2)		
60 - 69	776	2448	(57.9)	16	(0.4)	45	(0.6)	38	(0.5)	12	(0.3)	13	(0.2)	9	(0.4)		
70 and over	662	2145	(43.0)	15	(0.2)	46	(0.6)	37	(0.4)	12	(0.3)	13	(0.2)	9	(0.2)		
2 - 19	2068	2061	(20.9)	15	(0.2)	51	(0.3)	35	(0.3)	12	(0.1)	12	(0.1)	8	(0.1)		
20 and over	3745	2483	(24.2)	16	(0.1)	45	(0.3)	36	(0.2)	12	(0.1)	12	(0.1)	8	(0.1)	4	(0.2)
2 and over	5813	2379	(20.3)	15	(0.1)	47	(0.2)	36	(0.2)	12	(0.1)	12	(0.1)	8	(0.1)		
Females:																	
2 - 5	472	1417	(24.8)	14	(0.2)	54	(0.5)	33	(0.4)	12	(0.2)	11	(0.2)	7	(0.2)		
6 - 11	710	1902	(37.9)	13	(0.3)	53	(0.5)	35	(0.4)	12	(0.2)	11	(0.1)	8	(0.2)		
12 - 19	841	1812	(57.3)	14	(0.3)	51	(0.8)	36	(0.5)	12	(0.2)	12	(0.2)	9	(0.2)		
20 - 29	605	1963	(30.3)	15	(0.3)	46	(0.7)	38	(0.7)	13	(0.2)	12	(0.3)	9	(0.3)		
30 - 39	667	1916	(27.2)	15	(0.2)	47	(0.6)	37	(0.5)	12	(0.2)	13	(0.2)	9	(0.2)		
40 - 49	621	1805	(44.9)	15	(0.2)	48	(0.7)	36	(0.7)	12	(0.4)	12	(0.3)	9	(0.2)		
50 - 59	706	1859	(48.5)	16	(0.4)	47	(0.7)	36	(0.4)	12	(0.2)	13	(0.2)	8	(0.2)		
60 - 69	744	1745	(46.9)	16	(0.3)	45	(0.8)	38	(0.6)	12	(0.3)	13	(0.3)	9	(0.3)		
70 and over	619	1648	(32.0)	15	(0.3)	47	(0.5)	37	(0.6)	12	(0.4)	13	(0.2)	9	(0.1)		
2 - 19	2023	1757	(31.5)	14	(0.2)	52	(0.4)	35	(0.3)	12	(0.1)	11	(0.1)	8	(0.1)		
20 and over	3962	1829	(17.7)	15	(0.1)	47	(0.3)	37	(0.3)	12	(0.2)	13	(0.1)	9	(0.1)	2	(0.2)
2 and over	5985	1812	(12.1)	15	(0.1)	48	(0.3)	37	(0.2)	12	(0.1)	12	(0.1)	9	(0.1)		
Males and females:																	
2 - 19	4091	1911	(19.7)	14	(0.2)	52	(0.3)	35	(0.2)	12	(0.1)	12	(0.1)	8	(0.1)		
20 and over	7707	2144	(14.9)	16	(0.1)	46	(0.2)	37	(0.2)	12	(0.1)	12	(0.1)	9	(0.1)	3	(0.2)
2 and over	11798	2089	(11.3)	15	(0.1)	47	(0.2)	36	(0.1)	12	(0.1)	12	(0.1)	8	(0.1)		

Footnotes

- ¹ Percentages estimated as a ratio of each individual's energy intake of protein, carbohydrate, fat and alcohol divided by the individual's total food energy intake.
- ² Sample size and mean energy includes individuals (n = 3) with zero energy intake that are excluded from estimates of percentages of energy from protein, carbohydrate, fat, and alcohol.

Abbreviations

SE = standard error.

Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2017-March 2020 Prepandemic

The 2017—March 2020 files represent a unique public-use data release from NHANES. The coronavirus disease 2019 (COVID-19) pandemic required suspension of data collection in March 2020. As a result, the partially completed NHANES 2019–2020 cycle was not nationally representative. Therefore, the 2019—March 2020 data were combined with the data from the 2017–2018 cycle to create the nationally representative 2017—March 2020 prepandemic data files.

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2017-March 2020 Prepandemic. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2017-2018 and 2019-2020 as applicable www.ars.usda.gov/nea/bhnrc/fsrg.

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

Suggested Citation

U.S. Department of Agriculture, Agricultural Research Service. 2022. Energy Intakes: Percentages of Energy from Protein, Carbohydrate, Fat, and Alcohol, by Male/Female and Age, What We Eat in America, NHANES 2017-March 2020 Prepandemic.