

The effects of combining *Artemisia annua* and *Curcuma longa* ethanolic extracts in broilers challenged with infective oocysts of *Eimeria acervulina* and *E. maxima*

GUSTAVO F. D. ALMEIDA^{1*}, STIG M. THAMSBORG², ALDA M. B. N. MADEIRA³, JORGE F. S. FERREIRA⁴, PEDRO M. MAGALHÃES⁵, LUIZ C. DEMATTÊ FILHO⁶, KLAUS HORSTED¹ and JOHN E. HERMANSEN¹

¹ Department of Agroecology, Faculty of Sciences and Technology, Aarhus University, Research Centre Foulum, Blichers Allé 20, P.O. Box 50, DK-8830 Tjele, Denmark

² Danish Centre for Experimental Parasitology, Faculty of Life Sciences, University of Copenhagen, Dyrølægevej 100, DK-1870 Frederiksberg C, Denmark

³ Department of Parasitology, Institute of Biomedical Sciences, University of São Paulo, Av. Prof. Lineu Prestes, 1374. São Paulo, SP, Brazil

⁴ USDA-ARS, US Salinity Laboratory, 450 W. Big Springs Rd., Riverside, CA 92507-4617, USA

⁵ Chemical, Biological and Agricultural Pluridisciplinary Research Centre (CPQBA), University of Campinas – UNICAMP, P.O. Box 6171, BR-13081-970 Campinas, SP, Brazil

⁶ Department of Economy and Rural Sociology, Luiz de Queiroz College of Agriculture (ESALQ), University of São Paulo, Avenida Pádua Dias 11, Piracicaba, SP, Brazil

(Received 5 October 2012; revised 25 February and 28 April 2013; accepted 11 May 2013; first published online 22 October 2013)

SUMMARY

Due to an increasing demand for natural products to control coccidiosis in broilers, we investigated the effects of supplementing a combination of ethanolic extracts of *Artemisia annua* and *Curcuma longa* in drinking water. Three different dosages of this herbal mixture were compared with a negative control (uninfected), a positive control (infected and untreated), chemical coccidiostats (nicarbazin + narazin and, later, salinomycin), vaccination, and a product based on oregano. Differences in performance (weight gain, feed intake, and feed conversion rate), mortality, gross intestinal lesions and oocyst excretion were investigated. Broilers given chemical coccidiostats performed better than all other groups. Broilers given the two highest dosages of the herbal mixture had intermediate lesion scores caused by *Eimeria acervulina*, which was higher than in broilers given coccidiostats, but less than in broilers given vaccination, oregano and in negative controls. There was a trend for lower mortality ($P = 0.08$) in the later stage of the growing period (23–43 days) in broilers given the highest dosage of herbal mixture compared with broilers given chemical coccidiostats. In conclusion, the delivery strategy of the herbal extracts is easy to implement at farm level, but further studies on dose levels and modes of action are needed.

Key words: broiler, natural anti-protozoa drugs, coccidiostats, drug combination, plant extract, herbal medicine.

INTRODUCTION

Coccidiosis is a very serious parasitic disease in the broiler industry (Shirley *et al.* 2005) and is caused by highly host-specific protozoan parasites belonging to the genus *Eimeria* (Williams, 1999). Infections caused by *E. acervulina*, *E. maxima* and *E. tenella* are frequently diagnosed in intensive poultry systems (McDougald *et al.* 1997) and the control of these species is usually given higher priority than other *Eimeria* species affecting broilers (Shirley *et al.* 2005).

Coccidiosis is primarily controlled by medication and a range of limitations have been reported in the past few years (Martin *et al.* 1997). Underdosing of

anticoccidials in broiler feed could lead to parasitic resistance (Dauguschies *et al.* 1998) and the extensive use of drugs has led to the development of resistance to coccidiostat products (Chapman, 1993), giving rise to a need for alternative approaches in controlling coccidiosis.

The substitution of ionophore drugs is unlikely to be possible with a single-strand approach, but a comprehensive combination of strategies could help to reduce the prevalence of coccidiosis in intensive production systems (Pinard-van der Laan *et al.* 1998). Research has been carried out to find effective, but non-pathogenic vaccines (Lillehoj and Trout, 1993; Williams, 2002) to be used in the substitution of anticoccidial products. The use of plants and extracts as therapeutics might likewise be an option (Naidoo *et al.* 2008; Akhtara *et al.* 2012). Thus, natural ingredients, e.g. plants, extracts and their combinations, rather than synthetic drugs are

* Corresponding author: Department of Agroecology, Faculty of Sciences and Technology, Aarhus University, Research Centre Foulum, Blichers Allé 20, P.O. Box 50, DK-8830 Tjele, Denmark. E-mail: Gustavo.deAlmeida@agrsci.dk; gufoal@gmail.com

believed to have an important role to play in the near future because they are usually residue-free and well accepted by consumers (Orengo *et al.* 2012). For example, *Artemisia annua* has previously been suggested for control of *Eimeria acervulina* given as prophylactic supplementation before a coccidial challenge (Allen *et al.* 1997), and we found that a 3% content of dried *A. annua* leaves in the feed of broilers 3 weeks before subjecting the broilers to a coccidial challenge reduced oocyst excretion by 60–70% (Almeida *et al.* 2012). However, at the same time, feed intake was suppressed due to a larger content of unpalatable substances in the feed. Also, it has been shown that different levels of artemisinin (the main bioactive component of *A. annua*) in feed and oral supplementation had no effect against *E. maxima*, suggesting a species-specific mode of action (Allen *et al.* 1997; Arab *et al.* 2006).

Another relevant plant with known anti-protozoan properties is the spice turmeric, *Curcuma longa* (Shahiduzzaman *et al.* 2009), which has been suggested as a potential candidate for treating human malaria (Reddy *et al.* 2005; Cui *et al.* 2007). In Pakistan, smallholder farmers add turmeric powder to the feed to control coccidiosis in broilers (Abbas *et al.* 2012). Allen *et al.* (1998) reported a reduction in intestinal lesions and lower oocyst excretion with 1% dietary inclusion for broilers challenged with *E. maxima* oocysts. Here the effective bioactive ingredient is supposedly curcumin (diferuloylmethane), a phenolic compound with high antioxidative (Subramanian *et al.* 1994), anti-inflammatory (Huang *et al.* 1997) and anti-carcinogenic properties (Rao *et al.* 1995; Jung *et al.* 2004), and found in *C. longa* roots in concentrations ranging from 1 to 5% (Conney *et al.* 1991).

With the reported anti-protozoan activity of both artemisinin and curcumin, we hypothesized that a combination of these ingredients in *A. annua* and *C. longa* herbal extracts could provide suppressive effects and act against dual infections caused by *E. maxima* and *E. acervulina* oocysts in fast-growing broilers, similar to chemical coccidiostats.

On this background, this study was undertaken to evaluate the effect of three different dose levels of herbal extracts supplying artemisinin:curcumin for avian coccidiosis. The extracts were administered in the drinking water and compared with appropriate controls, with broilers receiving chemical products (representing the conventional production system), and with broilers vaccinated or given a product based on oregano, thus representing alternative strategies available in the market to control coccidiosis in organic production systems.

MATERIALS AND METHODS

Experimental design and broilers

The study was carried out at the experimental facilities of Korin Agricultural Ltd. (www.korin.com.br)

located in the municipality of Ipeúna, Sao Paulo, Brazil (22°24'S, 47°41'W). On 15 August 2011, 1-day-old male broilers (Cobb 500S, $n=1440$) previously vaccinated in hatchery against Marek's Disease, fowlpox and Gumboro Disease were randomly allocated to 48 pens distributed in eight treatment combinations with six replicates of 30 broilers, giving a total of 180 broilers per treatment. Three different dose levels of a combination of *A. annua* and *C. longa* ethanolic extracts (Low, Medium and High) supplemented in drinking water were compared with a negative control (uninfected, untreated), a positive control (infected and untreated), treatment with chemical coccidiostats (see next section), vaccination and a product based on oregano. Table 1 describes the treatments, the supplementation strategies, doses, age and duration of treatments. Each pen measured 3.0 m² (1.5 × 2 m) with food and water supplied *ad libitum*.

All pens were cleaned and disinfected 2 days before the beginning of the experiment and high standards of hygiene were followed during the trial. All practical handling was performed from outside the 48 pens in lateral corridors and the pens were isolated from each other by the use of PVC walls. When necessary, technicians used disposable plastic bags over their shoes to avoid contamination. Broilers were raised on concrete floors with wood shavings until they were 43 days of age, when the experiment was terminated. Broilers were raised in the comfort range of temperature and humidity recommended by the guidelines for the breed and the experimental protocols were in accordance with international guidelines for animal care and health and with Brazilian legislation.

Experimental diets, water supply and oral administration of herbal extracts

The feeding programme in our study followed the routine of farmers linked to the collaborative broiler company with diets adapted to the different development periods: initial (1–8 days), growth (9–22 days), fattening I (23–29 days), fattening II (30–36 days) and final (37–43 days). Broilers in the control groups (positive, T1 and negative, T4) were fed the basic diet without any other type of supplementation. Nicarbazin + narazin (Maxiban[®]) 50 ppm was supplemented from day 1 to 22 and salinomycin (Coxistac[®]) 66 ppm from day 22 to 39 in the diet of broilers receiving the chemical coccidiostats (T3). Livacox vaccine at recommended doses (Merial Ltd. Brazil) was administered to the broilers on the day they arrived from the hatchery by the use of the movable water dispensers situated in each replicated pen (T2). Broilers in group T5 were supplemented with a product based on oregano extract containing carvacrol + timol (Regano[®]) and broilers in T6, T7

Table 1. Description of treatments, acronyms, supplementation strategy and age of the broilers when treatments were supplemented

Treatment	Acronym	Delivering strategy ^a	Age of the broilers (days)
Negative control (untreated, uninfected)	T1	–	–
Vaccinated ^b , infected ^c	T2	Drinking water	1
Cocciostat ^d , infected	T3	Dietary inclusion	1–39
Positive control (untreated, infected)	T4	–	–
Regano ^{®c} , Infected	T5	Drinking water	1–39
Dose 1 of ethanolic herb extracts of <i>A. annua</i> + <i>C. longa</i> . (11 ppb artemisinin + 45 ppb curcumin), Infected	T6	Drinking water	1–42
Dose 2 of ethanolic herb extracts of <i>A. annua</i> + <i>C. longa</i> . (19 ppb artemisinin + 74 ppb curcumin), Infected	T7	Drinking water	1–42
Dose 3 of ethanolic herb extracts of <i>A. annua</i> + <i>C. longa</i> . (27 ppb artemisinin + 107 ppb curcumin), Infected	T8	Drinking water	1–42

^a Preparation of mixtures were performed every second day. 155 L of tap water were mixed with dosages of herbal ingredients in T5, T6, T7 and T8.

^b Livacox vaccine (Meril Ltd. Brazil) consisted of attenuated oocysts of *E. acervulina*, *E. maxima* and *E. tenella*.

^c Three (3) 'seeders' per pen (10%/pen) aged 8 days old where orally inoculated with 1 mL suspension containing 5×10^4 *E. maxima* and 2×10^5 *E. acervulina* sporulated oocysts. 'Seeders' were removed from the pens at 17 days old, 9 days post-inoculation.

^d 50 ppm Nicarbazin + Narazin (Maxiban[®]) supplemented from 1 to 22 days of age and 66 ppm Salinomycin 12% (Coxistac[®]) supplemented from 22 to 39 days of age.

^e 42 g Kg⁻¹ Carvacrol + 1.25 g kg⁻¹ Timol (Regano[®]).

and T8 received the herbal mixtures via drinking water from day 1 of their lives.

Eight water tanks (one tank per treatment) of 160 L capacity were raised above ground in the chicken house and water was distributed by gravity to each replicated pen ($n = 6$ per treatment) by the use of plastic hoses. Every other day from the beginning of the trial, the water in the tanks including the herbal extract mixture, also including T5, was replaced. Treatment supplements in water were prepared at 2-day intervals to avoid any confounding factor related to possible instability or oxidation of the herbal components.

The herbal extracts of *A. annua* and *C. longa* were prepared at CPQBA – University of Campinas, Brazil – using the same extraction procedure for both plant species. In summary, 100 g of dried leaves or 100 g curcuma powder were mixed with 1 L ethanol 70% (700 mL ethanol 97° + 300 mL distilled water) in two 1-L Erlenmeyer flasks wrapped with aluminium foil to prevent oxidation and degradation by light. The mixtures were kept in a dark, temperature-controlled room at 22 °C and homogenized with circular movements twice a day for 20 days. The contents were filtered by using paper filters and transferred to 100-mL flasks with dropping glass dispensers. Flasks were properly labelled and stored in the refrigerator at 5 °C until needed.

One week before the trial, the flasks containing the ethanolic herbal extracts of *A. annua* and *C. longa* were transported to the experimental station at Korin Agropecuária Ltd. and kept in the fridge before use. One day before allocating chicks to pens, the ethanolic concentrated solutions were mixed with

155 L tap water using the same volume of the herbal extracts for each dosage (1.5 mL of each extract in T6, 2.5 mL for T7 and 3.6 mL for T8). The content of active ingredients was estimated by HPLC-UV according to Ferreira and Gonzales (2009). The *A. annua* ethanolic extract was found to contain 1.15 mg artemisinin mL⁻¹, while from proximal calculations, the ethanolic extract of *C. longa* had an estimated concentration of 4.6 mg curcumin mL⁻¹. From relevant literature on curcumin and artemisinin contents in raw materials and in herbal ethanolic extracts (Souza and Glória, 1998; Ferreira *et al.* 2011 respectively) we estimated that the three dosages supplemented provided a 1:4 ratio of artemisinin:curcumin (Table 1).

Transmission by contact and monitoring strategy

At day 8, three randomly chosen broilers per pen ($n = 18$ per treatment) were artificially inoculated by gavage with a 1-mL suspension of distilled water with 5×10^4 *E. maxima* + 2×10^5 *E. acervulina* sporulated oocysts. Strains of intermediary virulence were obtained in the Laboratory of Molecular Biology of Coccidians, Department of Parasitology, University of Sao Paulo. These broilers were nominated 'seeders' and were raised with contemporary broilers with the aim of transmitting the disease by contact (Velkers *et al.* 2010). Seeders were removed from the pens 9 days post inoculation and harvested (cervical dislocation) for scoring of lesions ($n = 5$ per treatment) in a strategy adapted from Johnson and Reid (1970) and supported by illustrations provided by Anonymous (1990). Results (Table 3) of scoring

of lesions show the distribution of lesions caused by *E. acervulina* (duodenum) and *E. maxima* (middle section of small intestine) separately as proposed by Conway *et al.* (2003).

To assess the spread of infection by the seeders, five broilers were randomly selected in each pen and nominated as 'tracers'. Twice weekly starting 3 days post inoculation of seeders, the tracers were placed in communal cages (one cage per pen) until one drop per tracer was available for pooled samplings of fecal samples. In total, eight pooled fecal samples per pen were examined during the trial. On the same day as fecal collection, pooled samples ($n = 48$) were analysed by a modified McMaster technique using saturated NaCl solution with 50% glucose monohydrate as flotation fluid with a sensitivity of 20 oocysts per g of feces (OPG) (Almeida *et al.* 2012).

Performance attributes

Consumption of feed per pen was monitored and recorded every week. Individual body weight of the same 10 broilers in each pen (not seeders or tracers) ($n = 60$ per treatment) was measured every week. We estimated performance attributes by considering the mean body weight at 24 h post hatch, at 22 days of age, and when broilers were 43 days old before slaughter. With this strategy, we could estimate body weight gain, daily gain, feed intake and feed conversion rate for two main periods—before and after expected infection and for the total experiment (1–22 days; 23–43 days and 1–43 days of age). Results of performance attributes are presented as the mean \pm S.E.M. and averaged for each treatment combination. Performance attributes were the most important indicators of responses to different treatments, while lesion scores and infection dynamics provided by OPG were supportive indicators.

Statistical analysis

Broilers in our study were allocated to pens in a completely randomized design. Differences in performance attributes were assessed by analysis of variance with one-way ANOVA in SAS and the dependent variables were the mean values for each attribute observed for each pen (individual body weight gain, feed intake and feed conversion rate) and the independent variables were the treatments. Data on mortality were analysed by the Chi-square test. Categorical data on lesion score were analysed using the GENMOD procedure in SAS (DIST = MULTINOMIAL) with link functions for cumulative probabilities (LINK = CUMLOGIT). Differences between individual treatments were tested with Fisher's Exact Test for 2×2 tables for small samples (Quinn and Keough, 2002). The infection dynamics were analysed by a mixed procedure (PROC MIX) in SAS (SAS, 2000) where the

statistical unit was the logarithmically transformed value for each OPG observation according to the following model:

$$Y_{ijk} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + A_{k(ij)} + \varepsilon_{ijk}$$

where Y_{ijk} is the natural logarithm of the number of oocysts per gram of feces (OPG) for each pen in each sampling date; μ is the mean value; α_i is the treatment, i is 1, 2, 3, 4, 5, 6, 7, 8 (*Uninfected negative control; Vaccinated, infected; Coccidiostat, infected; Untreated, Infected; Regano[®], Infected; Dose 1 (LOW) of herbal extracts, Infected; Dose 2 (MEDIUM) of herbal extracts, Infected; Dose 3 (HIGH) of herbal extracts, Infected*); β_j is the OPG sampling date; j is (1 to 8); $(\alpha\beta)_{ij}$ is the interaction treatment \times OPG sampling date; A_k is the random effect of pen number (1 to 48) and ε_{ijk} is the residual. Sampling date was defined as repeated measurement and ε_{ijk} was assumed to have a multivariate distribution where observations from different treatments were uncorrelated, while observations from different weeks were assumed to have a Toeplitz (Type = TOEP) correlation structure. P -values less than or equal to 0.05 were considered statistically significant.

RESULTS

For the initial growing period (1–22 days), broilers supplemented with the botanical mixtures (T6, T7 and T8) and broilers vaccinated (T2) had lower feed intakes compared with broilers in the uninfected group (T1) (Table 2). In the final period investigated (22–43 days), broilers supplemented with Regano[®] (T5) and broilers vaccinated (T2) had the highest feed intakes compared with broilers in the negative control group (T1). Measured over the entire experimental period, feed intake was highest for the broilers consuming Regano[®] (T5) and lowest for the broilers consuming the High dose of the botanical mixture (T8).

In the initial period (1–22 days), body weight gain (BWG) was higher for the broilers supplemented with the chemical coccidiostats (T3) than in most other treatments, whereas broilers vaccinated (T2), supplemented with Regano[®] (T5), and given the High dose of the herbal mixture (T8) had the lowest BWG.

In the second growing period (22–43 days), after the dissemination of oocysts by the seeders, supplementation of chemical coccidiostats resulted in a higher BWG than supplementation with the herbal mixture, but with no difference from the vaccinated broilers (T2), the broilers in the negative control group (T1) and the broilers fed Regano[®] (T5). There were no differences in FCR in any of the periods investigated (Table 2). For the total duration of the experiment (1–43 days), BWG was higher for the broilers given chemical coccidiostats than in all other

Table 2. Performance attributes for each treatment* for two distinct periods under investigation and for the entire period investigated. Periods consisted of 1 to 22 days of age, 22 to 43 days of age and 1 to 43 days of age. Degree of infection is represented by the total sum of OPG over 8 consecutive samplings for each treatment combination. All results presented are in LS means \pm S.E.M.

Performance attributes		P values	Negative control (T1)	Vaccine (T2)	Chemical coccid. (T3)	Positive control (T4)	Regano (T5)	Low dose (T6)	Medium dose (T7)	High dose (T8)
First period 1–22 day	BWG ¹ (g)	0.01	838 \pm 15 ^{ab}	789 \pm 13 ^{cde}	873 \pm 09 ^a	818 \pm 14 ^{bcd}	779 \pm 15 ^{de}	836 \pm 12 ^{ab}	821 \pm 13 ^{bc}	768 \pm 16 ^e
	FI ² (g day ⁻¹)	0.01	68.9 \pm 2.3 ^a	62.6 \pm 0.5 ^{bc}	66.2 \pm 0.9 ^{ab}	65.1 \pm 1.1 ^{abc}	65.6 \pm 1.2 ^{abc}	63.7 \pm 0.4 ^{bc}	64.5 \pm 1.4 ^{bc}	61.6 \pm 0.9 ^c
	FCR ³	0.38	1.71 \pm 0.07	1.67 \pm 0.04	1.59 \pm 0.04	1.67 \pm 0.04	1.75 \pm 0.05	1.59 \pm 0.03	1.65 \pm 0.06	1.70 \pm 0.03
	Mort. ⁴ (%)	0.56	3.3 \pm 2.1	1.1 \pm 1.1	0.0 \pm 0.0	1.1 \pm 0.7	2.7 \pm 1.2	1.7 \pm 1.2	1.1 \pm 0.7	1.3 \pm 0.8
Second period 22–43 day	BWG (g)	0.05	2010 \pm 28 ^{ab}	2002 \pm 35 ^{ab}	2103 \pm 29 ^a	1993 \pm 44 ^b	2013 \pm 27 ^{ab}	1916 \pm 35 ^b	1946 \pm 37 ^b	1964 \pm 39 ^b
	FI (g day ⁻¹)	0.02	177.3 \pm 2.9 ^{bc}	187.7 \pm 5.8 ^{ab}	180.9 \pm 3.2 ^{bc}	180.1 \pm 2.0 ^{bc}	194.1 \pm 5.1 ^a	177.4 \pm 2.8 ^{bc}	183.0 \pm 2.8 ^{bc}	175.1 \pm 3.0 ^c
	FCR	0.08	1.80 \pm 0.03	1.88 \pm 0.07	1.71 \pm 0.03	1.85 \pm 0.05	1.91 \pm 0.05	1.87 \pm 0.05	1.93 \pm 0.06	1.87 \pm 0.04
	Mort. (%)	0.20	2.5 \pm 1.3	4.5 \pm 2.1	5.6 \pm 0.9	3.0 \pm 1.2	5.0 \pm 1.4	3.1 \pm 1.2	3.0 \pm 1.1	0.0 \pm 0.0
Total period 1–43 day	BWG (g)	0.01	2848 \pm 36 ^b	2791 \pm 41 ^b	2976 \pm 34 ^a	2811 \pm 46 ^b	2792 \pm 34 ^b	2787 \pm 39 ^b	2767 \pm 42 ^b	2732 \pm 47 ^b
	FI (g day ⁻¹)	0.02	124.4 \pm 1.9 ^{bc}	127.2 \pm 4.0 ^{ab}	125.5 \pm 1.7 ^{abc}	123.6 \pm 1.2 ^{bc}	132.0 \pm 3.1 ^a	121.4 \pm 1.6 ^{bc}	125.1 \pm 2.3 ^{abc}	118.4 \pm 1.9 ^c
	FCR	0.10	1.79 \pm 0.03	1.85 \pm 0.06	1.71 \pm 0.02	1.81 \pm 0.04	1.90 \pm 0.05	1.80 \pm 0.04	1.86 \pm 0.06	1.82 \pm 0.02
	Mort. (%)	0.59	5.9 \pm 3.2	5.6 \pm 2.6	5.6 \pm 0.9	4.1 \pm 1.5	7.7 \pm 1.3	4.8 \pm 1.7	4.1 \pm 1.4	1.3 \pm 0.8
Total OPG ⁵	0.05	7.4 \pm 0.9 ^{ab}	7.3 \pm 0.9 ^{ab}	5.3 \pm 0.7 ^b	22.1 \pm 2.7 ^a	15.5 \pm 1.9 ^a	24.6 \pm 3.4 ^a	8.8 \pm 1.1 ^{ab}	12.7 \pm 1.5 ^{ab}	

Number in a row marked with different letters are significantly different ($P < 0.05$).

* Treatments consisted of: T₁ = Uninfected negative control (untreated, uninfected); T₂ = Vaccinated, infected; T₃ = Coccidiostat, infected; T₄ = Untreated, Infected (positive control); T₅ = Regano[®], Infected; T₆ = Dose 1 (Low) of herbal extracts, Infected; T₇ = Dose 2 (Medium) of herbal extracts, Infected; T₈ = Dose 3 (High) of herbal extracts, Infected.

¹ BWG=Body weight gain for the period specified (g).

² FI=Feed intake for the period specified (g/broiler/day).

³ FCR=kg feed consumed/body weight gain in kg for the period specified.

⁴ Mort. (%)=Percentage mortality observed for the period specified.

⁵ Total OPG=Total sum over 8 sampling dates \times 100.




Fig. 1. Mean oocyst excretion over time for each treatment combination*. Samplings for OPG counts begun when broilers were 11 days old, the third day after the inoculation of the 'seeders' in respect to pre-patent periods of *Eimeria* spp. Treatments consisted of: T₁ = Negative control (untreated, uninfected); T₂ = Vaccinated, infected; T₃ = Coccidiostat, infected; T₄ = Untreated, Infected (Positive control); T₅ = Regano[®], Infected; T₆ = Dose 1 (Low) of herbal extracts, Infected; T₇ = Dose 2 (Medium) of herbal extracts, Infected; T₈ = Dose 3 (High) of herbal extracts, Infected.

treatments, but no significant differences between any of the other treatments were observed (Table 2).

Mortality was low in the first part of the experiment (1–22 days) with no deaths in the group given the chemical coccidiostat. However, in the second growing period (22–43 days), there was a trend for mortality to be lower ($P = 0.08$) for the broilers given the highest dosage of the herbal combination (T₈, 0.0%) compared with the birds given the chemical coccidiostat (T₃, 5.6%).

Table 3 illustrates the distribution of scores of lesions observed in the duodenum (*E. acervulina*) and in the small intestine (*E. maxima*), respectively, of the five seeders per treatment 9 days post inoculation. Overall, the treatment affected lesion score distribution significantly for both *E. acervulina* and *E. maxima* ($P < 0.01$ and $P < 0.05$, respectively). As expected, no lesions were observed for the broilers in the negative control group (T₁) and for broilers supplemented with the chemical coccidiostat (T₃) in the duodenum (*E. acervulina*). Vaccinated broilers (T₂) showed the highest frequency of lesions from *E. acervulina*, very similar to broilers in the positive control group (T₄).

Broilers supplemented with Regano[®] (T₅) did not differ from the positive control group (T₄) or vaccinated (T₂) in extent of lesions caused by *E. acervulina* (Table 3). Broilers supplemented with Low, Medium and High doses of the herbal mixture (T₆, T₇ and T₈ respectively) had an intermediary distribution of lesions, with response tending to be

dosage-dependent (Table 3). The two higher doses of the mixture *A. annua* and *C. longa* produced fewer lesions compared with the vaccinated broilers (T₂), broilers supplemented with Regano[®] (T₅) and broilers in the positive control group (T₄). However, higher scores were observed for treatments T₆, T₇ and T₈ when compared with broilers given the chemical coccidiostat (T₃) and broilers in the negative control group (T₁).

The scoring of lesions in the mid-intestinal mucosal surface (*E. maxima*) (Table 3) indicated that broilers receiving the two highest dosages of the herbal combination (T₇ and T₈) did not differ from broilers receiving the chemical coccidiostats (T₃). In addition, vaccinated broilers (T₂) had the lowest lesion scores in the middle section of the intestine, while broilers given Regano[®] (T₅) had the highest scores.

In terms of oocyst excretion, broilers in groups T₆ and T₄ had the highest infection peaks at 15 days post inoculation (PI) and a second peak at 22 days PI compared with broilers receiving the chemical coccidiostat (T₃), the uninfected group (T₁) and broilers receiving the Medium dose of the combined *A. annua* and *C. longa* extracts (T₇) (Fig. 1). Considering the total number of oocysts excreted, broilers supplemented with Medium (T₇) and High (T₈) levels of herbal ethanolic extracts showed intermediary numbers of OPG that were no different from those recorded for vaccinated broilers (T₂) and for broilers in the uninfected group (T₁), and not

Table 3. Distribution of lesion scores in broilers inoculated with *E. acervulina* or *E. maxima* supplemented with different treatments

Treatment	Number of birds/category of lesion score ¹									
	<i>Eimeria acervulina</i>					<i>Eimeria maxima</i>				
	0	1	2	3	4	0	1	2	3	4
T1	5 ^b	0	0	0	0	4 ^a	1	0	0	0
T2	0 ^a	0	3	2	0	3 ^{ab}	2	0	0	0
T3	5 ^b	0	0	0	0	2 ^a	3	0	0	0
T4	1 ^a	1	2	1	0	1 ^{ab}	3	1	0	0
T5	0 ^a	2	3	0	0	1 ^b	1	3	0	0
T6	0 ^{ac}	4	1	0	0	0 ^a	5	0	0	0
T7	1 ^c	4	0	0	0	1 ^{ac}	4	0	0	0
T8	2 ^{abc}	2	1	0	0	2 ^{ac}	3	0	0	0

¹ Distributions of scores in five categories based on severity of lesions (0 to 4). Different superscripts in the same column represent statistically different ($P \leq 0.05$) distributions after Fisher Exact Tests. Number of birds per treatment level equals 5.

Source: After Conway *et al.* 2003. Treatments consisted of: T₁ = Negative control (untreated, uninfected); T₂ = Vaccinated, infected; T₃ = Coccidiostat, infected; T₄ = Untreated, Infected (positive control); T₅ = Regano[®], Infected; T₆ = Dose 1 (Low) of herbal extracts, Infected; T₇ = Dose 2 (Medium) of herbal extracts, Infected; T₈ = Dose 3 (High) of herbal extracts, Infected.

statistically different from the broilers given chemical coccidiostats (Table 2).

DISCUSSION

To our knowledge, this is the first field study investigating the combined use of *A. annua* and *C. longa* (ethanolic extracts) against coccidiosis in broilers. The rationale for the study was the reporting of complementary effects of the artemisinin:curcumin combination against malaria (Nandakumar *et al.* 2006; Isacchi *et al.* 2012; Padmanaban *et al.* 2012). Our results showed a trend for a dose-dependent effect (based on a 1:4 ratio) against dual infections caused by *Eimeria acervulina* and *E. maxima* reflected by performance attributes and survival, reduced lesion score and excreted oocysts.

In general, broilers in the uninfected group (T1) showed no lesions in the mucosal surface caused by *E. acervulina* infection and a very low score attributed to infections caused by *E. maxima* (Table 3). However, when OPG counting was considered (Table 2 and Fig. 1), tracers located in two replicate pens in T1 started excreting oocysts between days 18 and 23, suggesting contamination of pen litter due to the opportunism of the *Eimeria* parasites.

Thus, when broilers were 25 to 30 days old, oocysts were found in two replicates of the negative group (T1). Even with the strict hygiene measures applied during the study, broilers in all six replicates of the uninfected group (T1) were contaminated and excreted oocysts in the final part of their growing period (22–43 days). Even though BWG for the second period did not differ from broilers fed chemical coccidiostats (T3), when the total period was taken into account, broilers in T1 had a smaller

BWG than broilers receiving chemical coccidiostats (T3) (Table 2). However, Rosen (1995) in a review of more than 1200 studies concluded that supplementation with ionophores improves performance attributes of fast-growing broilers, suggesting that the low infection experienced in tracer broilers for group T1 did not influence performance attributes and would be compatible with uninfected broilers. Nevertheless, flies, ants, other insects, and also the feet and hands of technicians may have served as vectors and highlight the fast dissemination of the *Eimeria* parasites (Henken *et al.* 1994).

In contrast, broilers in the positive group (T4) had the highest lesion scores for *E. acervulina* infections (Table 3) and a larger number of excreted oocysts, but with no increase in mortality compared with other groups (Table 2). Broiler seeders were artificially challenged with dual coccidial infections at 8 days of age and euthanized for lesion scoring 9 days post inoculation, allowing sufficient time to contaminate the litter in each pen in accordance with the pre-patent period of *E. acervulina* and *E. maxima* parasites (Eckert *et al.* 1995). From this strategy (even compromising the identification of lesions in broiler intestines in accordance with the pre-patent period of the parasites as proposed by Johnson and Reid (1970)), it allowed our group to assess the effects of treatments on intestinal lesions and, at the same time, allow seeders to infect fellow broilers by contact (Velkers *et al.* 2010) imitating the natural transmission pathway on commercial farms, which was the objective of the current study.

Vaccinated broilers had the highest lesion scores (three individuals scored 2 and two individuals scored 3) in the duodenum mucosal surface (Table 3). Perhaps the time of the *E. acervulina*

challenge, provoked by the artificial inoculation (2×10^5 oocysts), was too close to vaccination time, suggesting that the time lapse between vaccination and challenge was not sufficient to allow proper immunization, as also found by Oviedo-Rondón *et al.* (2006). This also harmonizes with the theory that antibodies released after an initial challenge take time to induce protection by blocking the development and replication of parasites (Crane *et al.* 1988; Hafeez *et al.* 2007; Anwar *et al.* 2008).

Vaccinated broilers (T2) had the lowest distribution of lesions in the middle section of the small intestine (three individuals scored 0 and two individuals scored 1), which is the site of *E. maxima* infections, the results being comparable to uninfected controls (T1). This could be either because vaccination provided sufficient immunity against *E. maxima*, quite different from what was observed with the absence of protection against *E. acervulina*, or because – according to Mathis (2005) – heavier challenges from *E. acervulina* may inhibit the development of *E. maxima*. The former hypothesis sounds more plausible since the lesion scores caused by *E. acervulina* were higher for broilers in T2 (Table 3).

We observed a trend in mortality of broilers and lesions caused by *E. acervulina* for dose-dependent response when supplementing ethanolic extracts of *A. annua* and *C. longa* equivalent to a artemisinin: curcumin ratio of 1.4. In addition, when assessing the scoring of lesions in the mid-intestinal mucosal surface caused by *E. maxima* infections (Table 3), broilers in T7 and T8 did not differ from broilers supplemented with the chemical coccidiostats (T3).

In terms of total numbers of oocysts excreted by tracers, broilers given Medium and High doses did not respond differently from broilers receiving the chemical coccidiostat supplement (Table 2). But on the other hand, the two higher dosages of the mixtures *A. annua* and *C. longa* (T7 and T8) provided some protection compared with the untreated group (T4), although not statistically different for the total OPG (Table 2). Also (as supported by Table 3), the lesion score caused by *E. acervulina* was significantly lower for group T7 than for group T4 (infected, untreated). For the total period, groups T7 and T8 had lower mortality rates (4.1 and 1.3%, respectively) than the group treated with the chemical coccidiostat (T3, 5.6%). Even though they were not different in terms of total number of oocysts excreted (Table 2), the infection dynamics (Fig. 1) illustrate that the broilers receiving the Medium (T7) dose had a smaller but not statistically different infection peak than broilers receiving the High dose (T8), comparable to broilers on the chemical coccidiostat.

Although one could argue that the interval between samplings for OPG did not capture the peak of oocyst excretion, in our opinion the combination of measured parameters (OPG and lesion score) helped explain the effects on the main indicators

(performance attributes), and thus captured the differences observed in tested treatments.

Strategies to control coccidiosis in conventional production systems are dependent on chemical drugs, while in organic systems live vaccines or commercial preparations of plant-based products are used (Abbas *et al.* 2012). Some studies associate vaccination with botanical products. For example, Waldenstedt (2003) suggested vaccination in combination with oregano-containing products to improve the intestinal health of chickens and thus reduce the effects caused by coccidiosis.

From the results achieved in this study, we conclude that broilers given a supplement of chemical coccidiostats showed superior performance attributes during the trial due to a better protection against dual infections caused by *E. acervulina* and *E. maxima* parasites. However, despite a lower BWG for the two highest doses of the herbal mixture, we suggest that the trend in reduced mortality, the protection against lesions and reduced OPG counts provided by the high dose of the combined *A. annua* and *C. longa* ethanolic extracts, compared with the positive control, are indications that the supplementation of herbal extracts in drinking water can be a feasible alternative method of coccidia control in intensive organic production systems. It is affordable, easy to implement at farm level and is residue-free. However, the suitability of the herbal extracts used in this study to control clinical coccidiosis resulting from exposure to high infection pressure, or more pathogenic species, remains to be investigated. Further investigations into dosages and modes of action of this combination would be useful. In addition, supplementation of herbal extracts in combination with other management practices may help farmers engaged in the production of broilers reduce the use of synthetic drugs.

ACKNOWLEDGEMENTS

The authors wish to thank Evandro Possamai, Dayana Pereira and Pedro Lorga from Korin agropecuaria Ltd. for the logistical and technical support during the study. We also wish to thank the anonymous reviewers who constructively contributed to the quality of the manuscript.

FINANCIAL SUPPORT

The Danish Ministry of Science, Technology and Innovation is thanked for financing collaboration between Brazilian and Danish scientists. Aarhus University, SOAR – Research School for Organic Agriculture and Food Systems and Korin Agropecuária Ltd. – are thanked for supporting our study.

REFERENCES

- Abbas, R. Z., Colwell, D. D. and Gilleard, J. (2012). Botanicals: an alternative approach for the control of avian coccidiosis. *World's Poultry Science Journal* **68**, 203–215. doi: 10.1017/S0043933912000268.

- Akhtara, M., Haia, A., Awaisa, M. M., Iqbal, Z., Muhammad, F., Haq, A. U. and Anwar, M. I.** (2012). Immunostimulatory and protective effects of *Aloe vera* against coccidiosis in industrial broiler chickens. *Veterinary Parasitology* **186**, 170–177.
- Allen, P. C., Lydon, J. and Danforth, H. D.** (1997). Effects of components of *Artemisia annua* on coccidia infections in chickens. *Poultry Science* **76**, 1156–1163.
- Allen, P. C., Danforth, H. D. and Augustine, P. C.** (1998). Dietary modulation of avian coccidiosis. *International Journal for Parasitology* **8**, 1131–1140.
- Almeida, G. F. d., Horsted, K., Thamsborg, S. M., Kyvsgaard, N. C., Ferreira, J. F. S. and Hermansen, J. E.** (2012). Use of *Artemisia annua* as a natural coccidiostat in free-range broilers and its effects on infection dynamics and performance. *Veterinary Parasitology* **186**, 178–187.
- Anonymous** (1990). *Diagnosis of Coccidiosis in Chickens*. Janssen Pharmaceutica, Beerse, Belgium.
- Anwar, M. I., Akhtar, M., Hussain, I., Muhammad, F. and Haq, A. U.** (2008). Effects of local gametocyte and livacox vaccines on live body weight gain and lymphoid organs in chickens. *Pakistan Veterinary Journal* **28**, 136–138.
- Arab, H. A., Rahbari, S., Rassouli, A., Moslemi, M. H. and Khosravirad, F. D. A.** (2006). Determination of artemisinin in *Artemisia sieberi* and anticoccidial effects of the plant extract in broiler chickens. *Tropical Animal Health and Production* **38**, 497–503.
- Chapman, H. D.** (1993). Resistance to anticoccidial drugs in fowl. *Parasitology Today* **9**, 159–162.
- Conney, A. H., Lysz, T., Ferraro, T., Abidi, T. F., Manchand, P. S., Laskin, J. D. and Huang, M. T.** (1991). Inhibitory effect of curcumin and some related dietary compounds on tumor promotion and arachidonic metabolism in mouse skin. *Advances in Enzyme Regulation* **31**, 385–396.
- Conway, D. P., Sasai, A. K., Gaafar, B. S. M. and Smothers, C. D.** (2003). Effects of different levels of oocyst inocula of *Eimeria acervulina*, *E. tenella*, and *E. maxima* on plasma constituents, packed cell volume, lesion score, and performance in chickens. *Avian Diseases* **37**, 118–123.
- Crane, M. S., Murray, P. K., Gnozzio, M. J. and MacDonald, T. T.** (1988). Passive protection of chickens against *Eimeria tenella* infection by monoclonal antibody. *Infection and Immunity* **56**, 972–976.
- Cui, L., Miao, J. and Cui, L.** (2007). Cytotoxic effect of curcumin on malaria parasite *Plasmodium falciparum*: inhibition of histone acetylation and generation of reactive oxygen species. *Antimicrobial Agents and Chemotherapy* **51**, 488–494.
- Dauguschies, A., Gässlein, U. and Rommel, M.** (1998). Comparative efficacy of anticoccidials under the conditions of commercial broiler production and in battery trials. *Veterinary Parasitology* **76**, 163–171.
- Eckert, J., Taylor, M., Catchpole, J., Licois, D., Coudert, P. and Bucklar, H.** (1995). Morphological characteristics of oocysts. In *Biotechnology. Guidelines of Techniques in Coccidiosis Research* (ed. Eckert, J., Braun, R., Shirley, M. and Coudert, P.), pp. 103–119. COST 89/820. Office for Official Publications of the European Communities, Luxembourg.
- Ferreira, J. F. S. and Gonzales, J. M.** (2009). Analysis of underivatized artemisinin and related sesquiterpene lactones by high-performance liquid chromatography with ultraviolet detection. *Phytochemical Analysis* **2**, 91–97.
- Ferreira, J. F. S., Peaden, P. and Keiser, J.** (2011). *In vitro* trematocidal effects of crude alcohol-ic extracts of *Artemisia annua*, *A. absinthium*, *Asimina triloba*, and *Fumaria officinalis*. *Parasitology Research* **109**, 1585–1592.
- Hafeez, M. A., Akhtar, M., Javed, M. T. and Haq, A. U.** (2007). Maternal immunization by egg propagated gametocyte vaccine to control *Eimeria tenella* infections in newly hatched chicks. *Parasitology Research* **100**, 1139–1141.
- Henken, A. M., Ploeger, H. W., Graat, E. A. M. and Carpenter, T. E.** (1994). Description of a simulation model for the population dynamics of *Eimeria acervulina* infection in broilers. *Parasitology* **108**, 503–512.
- Huang, M. T., Newmark, H. L. and Frenkel, K.** (1997). Inhibitory effects of curcumin on tumorigenesis in mice. *Journal of Cellular Biochemistry Supplement* **27**, 26–34.
- Isacchi, B., Bergonzi, M. C., Grazioso, M., Righeschi, C., Pietretti, A., Severini, C. and Bilia, A. R.** (2012). Artemisinin and artemisinin plus curcumin liposomal formulations: enhanced antimalarial efficacy against *Plasmodium berghei*-infected mice. *European Journal of Pharmaceutics and Biopharmaceutics* **80**, 528–534.
- Johnson, J. and Reid, W. M.** (1970). Anticoccidial drugs lesion scoring techniques in battery and floor-pen experiments with chickens. *Experimental Parasitology* **28**, 30–36.
- Jung, M., Lee, K., Kim, H. and Park, M.** (2004). Recent advances in artemisinin and its derivatives as antimalarial and antitumor agents. *Current Medical Chemistry* **11**, 1265–1284.
- Lillehoj, H. S. and Trout, J. M.** (1993). Coccidia: a review of recent advances on immunity and vaccine development. *Avian Pathology* **22**, 3–31.
- Martin, A., Danforth, H., Barta, J. and Fernando, M.** (1997). Analysis of immunological cross-protection and sensitivities to anticoccidial drugs among five geographical and temporal strains of *Eimeria maxima*. *International Journal for Parasitology* **27**, 527–533.
- Mathis, G. F.** (2005). Reasons for field problems with *E. maxima*: *E. acervulina* vs *E. maxima*. In *Proceeding of the IXth International Coccidiosis Conference*, Foz do Iguaçu, Brazil. 19–23 Sept 2005.
- McDougald, L. R., Fuller, L. and Mattiello, R.** (1997). A survey of coccidian on 43 poultry farms in Argentina. *Avian Diseases* **41**, 923–929.
- Naidoo, V., McGaw, L. J., Bisschop, S. P. R., Duncan, N. and Eloff, J. N.** (2008). The value of plant extracts with antioxidant activity in attenuating coccidiosis in broiler chickens. *Veterinary Parasitology* **153**, 214–219.
- Nandakumar, D. N., Nagaraj, V. A., Vathsala, P. G., Rangarajan, P. and Padmanaban, G.** (2006). Curcumin-artemisinin combination therapy for malaria. *Antimicrobial Agents and Chemotherapy* **50**, 1859–1860.
- Orengo, J., Buendía, A. J., Ruiz-Ibáñez, M. R., Madrid, J., Del Río, L., Catalá-Gregori, P., García, V. and Hernández, F.** (2012). Evaluating the efficacy of cinnamaldehyde and *Echinacea purpurea* plant extract in broilers against *Eimeria acervulina*. *Veterinary Parasitology* **185**, 158–163.
- Oviedo-Rondón, E. O., Clemente-Hernández, S., Salvador, F., Williams, P., Losa, R. and Stephen, F.** (2006). Essential oils on mixed coccidia vaccination and infection in broilers. *International Journal of Poultry Science* **5**, 723–730.
- Padmanaban, G., Nagaraj, V. A. and Rangarajan, P. N.** (2012). Artemisinin-based combination with curcumin adds a new dimension to malaria therapy. *Current Science* **102**, 704–711.
- Pinard-van der Laan, M.-H., Monvoisin, J.-L., Pery, P., Hamet, N. and Thomas, M.** (1998). Comparison of outbred lines of chickens for resistance to experimental infection with coccidiosis (*Eimeria tenella*). *Poultry Science* **77**, 185–191.
- Quinn, G. P. and Keough, M. J.** (2002). *Experimental Design and Data Analysis for Biologists*. 6th printing, 2007. Cambridge University Press, Cambridge, UK.
- Rao, C. V., Rivenson, A., Simi, B. and Reddy, B. S.** (1995). Chemoprevention of colon carcinogenesis by dietary curcumin, a naturally occurring plant phenolic compound. *Cancer Research* **55**, 259–266.
- Reddy, R. C., Vathsala, P. G., Keshamouni, V. G., Padmanaban, G. and Rangarajan, P. N.** (2005). Curcumin for malaria therapy. *Biochemical and Biophysical Research Communications* **326**, 472–474.
- Rosen, G. D.** (1995). Antibacterials in poultry and pig nutrition. In *Biotechnology in Animal Feeds and Animal Feeding* (ed. Wallace, R. J. and Chesson, A.), pp. 143–172. VCH Verlagsgesellschaft GmbH, Weinheim, Germany.
- SAS Institute** (2000). *SAS Online Doc[®], Version 9.2*. SAS Institute, Cary, NC.
- Shahiduzzaman, M., Dyachenko, V., Khalafalla, R. E., Desouky, A. Y. and Daugschies, A.** (2009). Effects of curcumin on *Cryptosporidium parvum* in vitro. *Parasitology Research* **105**, 1155–1161.
- Shirley, M. W., Smith, A. L. and Tomley, F. M.** (2005). The biology of avian *Eimeria* with an emphasis on their control by vaccination. *Advanced Parasitology* **60**, 285–330.
- Souza, C. R. A. and Glória, M. B. A.** (1998). Chemical analysis of turmeric from Minas Gerais, Brazil and comparison of methods for flavour free oleoresin. *Brazilian Archives of Biology and Technology* **41**, 218–224.
- Subramanian, M., Sreejayan Rao, M. N., Devasagayam, T. P. and Singh, B. B.** (1994). Diminution of singlet oxygen-induced DNA damage by curcumin and related antioxidants. *Mutation Research* **311**, 249–255.
- Velkers, F. C., Swinkels, W. J. C., Rebel, J. M. J., Bouma, A., Daemen, A. J. J. M., Klinkenberg, D., Boersma, W. J. A., Stegeman, J. A., de Jong, M. C. M. and Heesterbeek, J. A. P.** (2010). Effect of *Eimeria acervulina* infection history on the immune response and transmission in broilers. *Veterinary Parasitology* **173**, 184–192.
- Waldenstedt, L.** (2003). Effect of vaccination against coccidiosis in combination with an antibacterial oregano (*Origanum vulgare*) compound in organic broiler production. *Acta Agriculturae Scandinavica* **53**, 101–109.
- Williams, R. B.** (1999). A compartmentalized model for the estimation of the cost of coccidiosis to the world's chicken production industry. *International Journal for Parasitology* **8**, 1209–1229.
- Williams, R. B.** (2002). Fifty years of anticoccidial vaccines for poultry, 1952–2002. *Avian Disease* **46**, 755–802.